

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/112904>

Please be advised that this information was generated on 2017-12-06 and may be subject to change.

The economic analysis of visceral leishmaniasis control

Filip Meheus

The economic analysis of visceral leishmaniasis control

Proefschrift

ter verkrijging van de graad van doctor
aan de Radboud Universiteit Nijmegen
op gezag van de rector magnificus prof. mr. S.C.J.J. Kortmann,
volgens besluit van het college van decanen
in het openbaar te verdedigen op maandag 16 september 2013
om 13.30 uur precies

ISBN/EAN 978-94-6191-843-7

Design cover Ilse Stronks

Print Ipskamp Drukkers, Enschede

door

Filip André Magdalena Meheus
geboren op 7 september 1978
te Gent, België

© Copyright F Meheus, 2013. All rights reserved. No part of this book may be reproduced or transmitted in any form, by print or otherwise, without prior written permission of the holder of the copyright.

Promotoren: Prof. dr. Koos van der Velden
Prof. dr. Marleen Boelaert (Instituut voor Tropische Geneeskunde,
Antwerpen)

Copromotor: Dr. Rob Baltussen

**Manuscript-
commissie:** Prof. dr. André van der Ven (voorzitter)
Prof. dr. Piet Kager (Universiteit van Amsterdam)
Dr. Jeroen Smits

The economic analysis of visceral leishmaniasis control

Doctoral Thesis

to obtain the degree of doctor
from Radboud University Nijmegen
on the authority of the Rector Magnificus prof. dr. S.C.J.J. Kortmann,
according to the decision of the Council of Deans
to be defended in public on Monday, September 16, 2013
at 13.30 hours

by
Filip André Magdalena Meheus
Born on September 7, 1978
in Ghent, Belgium

| | |
|------------------------|-------------------------------------------------------------------------------------------------------|
| Supervisors: | Prof. dr. Koos van der Velden Prof. dr. Marleen Boelaert (Institute of Tropical Medicine, Antwerp) |
| Co-supervisor: | Dr. Rob Baltussen |
| Doctoral Thesis | Prof. dr. André van der Ven (chair) |
| Committee: | Prof. dr. Piet Kager (University of Amsterdam) Dr. Jeroen Smits |

CONTENTS

| | | |
|------------------|---------------------------------------------------------------------------------------------------------------|-----|
| Chapter 1 | General introduction | 9 |
| SECTION 1 | The burden of visceral leishmaniasis | 29 |
| Chapter 2 | Socio-economic aspects of neglected diseases: sleeping sickness and visceral leishmaniasis | 31 |
| Chapter 3 | The poorest of the poor: a poverty appraisal of households affected by visceral leishmaniasis in Bihar, India | 45 |
| SECTION 2 | The economic impact and costs of patient management of visceral leishmaniasis | 59 |
| Chapter 4 | The costs of patient management of visceral leishmaniasis in Muzaffarpur, Bihar, India | 61 |
| Chapter 5 | The household costs of visceral leishmaniasis care in south-eastern Nepal | 79 |
| Chapter 6 | The economic burden of visceral leishmaniasis in Sudan: an assessment of provider and household costs | 101 |
| SECTION 3 | The cost-effectiveness of treatment strategies for visceral leishmaniasis | 127 |
| Chapter 7 | Combination therapy for visceral leishmaniasis | 129 |
| Chapter 8 | Cost-effectiveness analysis of combination therapies for visceral leishmaniasis in the Indian subcontinent | 157 |
| Chapter 9 | General discussion | 179 |
| | Summary | 209 |
| | Samenvatting | 215 |
| | Acknowledgments | 221 |
| | Curriculum Vitae | 225 |
| | Publications | 229 |



CHAPTER 1

General introduction

This introduction is based in part on:

- Meheus F, Boelaert M (2010) The burden of visceral leishmaniasis in South Asia. *Trop Med Int Health* 15: 1-3.
- Meheus F, Rijal S, Lutumba P, Hendrickx D, Boelaert M (2012) NTD control and health system strengthening. *Lancet* 379: 2149-50 [letter].
- Uranw S, Hasker E, Roy L, Meheus F, Das ML, Bhattarai NR, et al. (2013) An outbreak investigation of visceral leishmaniasis among residents of Dharan town, eastern Nepal: evidence for urban transmission of *L. donovani*. *BMC Infect Dis* 13:21.
- Lutumba P, Makieya E, Shaw A, Meheus F, Boelaert M (2007) Human African trypanosomiasis in a rural community, Democratic Republic of Congo. *Emerg Infect Dis* 13: 248-254.

INTRODUCTION

Visceral leishmaniasis (VL) is a vector-borne disease caused by *Leishmania* parasites that are transmitted by the bite of an infected female sandfly. VL is an important public health problem and a significant cause of mortality and morbidity in the Indian subcontinent and East Africa. Since 2005 the governments of India, Nepal and Bangladesh are engaged in a collaborative effort to control and eliminate visceral leishmaniasis from the Indian subcontinent. Approaches to control aim on the one hand at improving the human recovery rate through early diagnosis and case management and on the other hand reducing vector density and biting rate through vector control interventions such as indoor residual spraying (IRS) and the use of bednets (WHO 2005). In East Africa such an integrated effort to control VL does not yet exist and control is mainly based on passive case detection and treatment, although bednets are distributed by non-governmental organizations (Ritmeijer *et al.* 2007).

However uptake of VL interventions remains low. VL occurs in regions with weak health systems and low quality public health services. The costs associated with an episode of VL are substantial and may drive already impoverished households further into poverty (Meheus *et al.* 2006; Uranw *et al.* 2013). In addition long delays in diagnosis are common since clinical symptoms often resemble that of other infections and households usually first visit informal or private health providers where health workers lack knowledge about appropriate treatments, and do not have the tools to diagnose VL. As a result VL affected households incur substantial financial and economics costs in seeking appropriate care.

Over the past years various new technologies have become available that may improve VL case management. Newer drugs provide opportunities for shorter, less expensive treatments that can be provided on an ambulatory basis, including the first oral VL drug, Miltefosine, adopted by the VL elimination initiative as first line treatment, and paromomycin which is very cheap (US\$15 for an adult full course) and administered intramuscularly. However both drugs are vulnerable for development of parasite resistance (Bryceson 2001; Seifert *et al.* 2003). More recently, a short course of liposomal amphotericin B showed to be very efficacious (Sundar *et al.* 2010), although the cost of the drug is still a barrier for use as a first line. Combination therapy for the treatment of visceral leishmaniasis has increasingly been advocated as a way to increase treatment efficacy and tolerance, reduce treatment duration and cost, and limit the emergence of drug resistance (Alvar *et al.* 2006; Croft *et al.* 2006). Once definite results from clinical trials are available combination therapies can be considered as first line treatment. The development of the rK39 immunochromatographic test, a rapid diagnostic test with high sensitivity and specificity, and the Direct Agglutination Test (DAT) make the rapid diagnosis

of VL possible and are less invasive than previous procedures (Sundar *et al.* 2002b). As these several options became available, the debate surrounding the best policy for case detection and treatment flared up.

Economic analysis provides an important set of tools to underpin the choice for evidence-based policy. Given the limited financial resources available to VL control and the increasing number of possible interventions, especially regarding drug treatment, economic evaluation provides us with methods to systematically compare all relevant options in terms of their costs and consequences and thereby contribute to more rational decision-making and efficient allocation of scarce resources. Moreover it can improve our understanding of factors influencing both the supply of VL interventions and factors related to the behaviour of VL affected households. While economic analysis is increasingly recognized by both national and international policy makers as an important tool in the development of VL control policies, the scarcity of reliable data, in particular on costs, severely hampers efforts to control the disease.

Against this background, I focus in this thesis on different aspects of economic analysis for VL control. The studies were carried out in three VL endemic countries – Nepal, India and Sudan. In the following paragraphs an overview is provided on the epidemiological and public health aspects of VL followed by the objectives and outline of the thesis.

LEISHMANIASIS

There are about twenty different species of leishmania that cause disease in humans. They cause a spectrum of diseases broadly categorized as cutaneous, mucocutaneous (MCL) and visceral leishmaniasis. Cutaneous leishmaniasis (CL) is the most common form of leishmaniasis and is endemic in 82 countries across all continents except Australia. The disease causes ulcers or nodules on exposed parts of the body (face, arms or legs) where the sand fly has bitten. CL is an important cause of disability, stigma and social isolation especially among women (Reithinger *et al.* 2005). MCL is a severe form of CL with progressive destructive ulcerations developing at the mucosal regions of nose and mouth and spreading to the throat (Chappuis *et al.* 2007). MCL is highly disfiguring and primarily occurs in Brazil, Bolivia & Peru.

The most severe form of leishmaniasis is visceral leishmaniasis (VL) and is fatal if untreated. VL is often referred to as *kala-azar*, which means black fever in Hindi due to the hyperpigmentation of the skin that may occur during the course of the disease (Sanyal 1985; Pearson & de Queiroz Sousa 1996). VL is mainly caused by the parasite species *L. donovani* (Chappuis *et al.* 2007) and transmitted by the bite of an infected female sandfly from infected to uninfected humans (i.e. anthroponotic VL; AVL). Although in East Africa, the presence of parasites has been shown in

rodents (Elnaïem *et al.* 2001) and dogs (Dereure *et al.* 2003), it is unclear whether there is also zoonotic transmission (i.e. from animals to humans) (Elnaïem *et al.* 1998a). On rare occasions VL transmission by blood transfusion has also been documented (Singh *et al.* 1996; Dey & Singh 2006).

Clinical signs and symptoms of VL include prolonged fever, fatigue, weakness, loss of appetite, weight loss and enlargement of the spleen and liver. After several months, untreated VL will result in the death of the infected individual mainly due to cachexia, bleeding and/or from secondary infections (Berman 1988). About 5-10% and 50% of patients that recover from VL after treatment in the Indian subcontinent and East Africa respectively, develop a skin condition named post kala-azar dermal leishmaniasis (PKDL) (Zijlstra 2001; Zijlstra *et al.* 2003). Depending on the grade of severity, PKDL develops as a rash, usually around the mouth and spreading to other parts of the body such as the arms and the upper torso (Zijlstra *et al.* 2003). Because parasites can be found within the lesions, PKDL is considered an important reservoir for the human transmission of the disease between epidemic cycles (Addy & Nandy 1992).

BURDEN OF VL: INCIDENCE AND MORTALITY

VL is reported in 62 countries (figure 2) with 200 million people at risk (Desjeux 1996). Recently the WHO estimated that worldwide about 200,000 to 400,000 new cases of VL and 50,000 to 60,000 deaths occur annually with the majority of VL cases (>90%) concentrated in only four countries: Bangladesh, India, Brazil and Sudan (WHO 2010; Alvar *et al.* 2012).

However because of a lack of reliable empirical data there is much uncertainty on figures of VL incidence. In the Indian subcontinent, the current officially reported figures are obtained through passive case detection in government health services and usually do not include cases detected by the private for-profit sector, which constitute a majority of the VL care providers in India. These figures therefore largely underestimate the actual number of cases. Extrapolation from official data sources is also difficult due to the focal distribution of VL (Bern *et al.* 2008). A number of studies have estimated the degree of underreporting, although these studies were limited to specific geographical areas. For instance, Desjeux (1992) found a 1:5 ratio of reported to unreported VL cases in community surveys in India. Singh *et al.* (2006) documented underreporting by a factor of eight in a community development block in Muzaffarpur district in Bihar, India in 2001–2003 and more recently Singh *et al.* (2010) estimated underreporting by a factor of four in Vaishali district, also in Bihar. In south Sudan, Collins *et al.* (2006) estimated that 91% of VL deaths were undetected.



Figure I: Geographical distribution of visceral leishmaniasis

Source: WHO 2010

The annual reported and estimated number of VL cases in the three study countries are shown in table 1. In Nepal, VL is reported in 12 districts situated in south-eastern Nepal in the Terai region bordering the highly VL-endemic northern state of Bihar in India. More than 90% of all cases in India are reported from Bihar State. In Sudan, Gedaref State is the epicentre of VL in East Africa (Burki 2009). The region is prone to epidemics and frequent outbreaks of VL with high case-fatality rates caused among other by malnutrition as well as mass population displacements due to war and famine. Between 1984 and 1994, a major epidemic of VL resulted in an estimated 100,000 VL deaths in the Western Upper Nile province of Sudan (Seaman *et al.* 1996). In 2010, Médecins sans Frontières reported an outbreak of VL in South Sudan leading to an eight-fold increase in the number of treated patients compared to the previous year (Médecins sans Frontières 2010).

Table I: Reported and estimated annual incidence of visceral leishmaniasis in India, Nepal and Sudan

| | Reported VL cases/year | Estimated annual VL incidence | | |
|--------------------------------------------------|------------------------|-------------------------------|----|---------|
| India | 34,918* | 146,700 | to | 282,300 |
| Nepal | 1,477* | 3,000 | to | 5,900 |
| Subtotal Indian subcontinent and South East Asia | 42,623 | 162,100 | to | 313,600 |
| Sudan | 3742* | 15,700 | to | 30,300 |
| Subtotal East African region | 8,569 | 29,400 | to | 56,700 |
| Elsewhere | 7,034 | 6,200 | to | 18,800 |
| Total worldwide | 58,227 | 202,200 | to | 389,100 |

* Years of report: Sudan 2005-2009; India & Nepal: 2004-2008

Source: Compiled from Alvar *et al.* 2012

ECONOMIC BURDEN OF VL

The economic burden of VL is measured in terms of two broad categories of costs. These include the direct costs associated with accessing treatment (e.g. drugs, laboratory tests, transportation) and the indirect costs associated with for example absenteeism or reduced productivity due to VL illness, which translates into an income loss to the household. For a review on the economic burden of VL we refer to chapter 2.

CONTROL OF VL

In the absence of a vaccine and in areas with *L. Donovanii* transmission, the control of VL is mainly based on accurate diagnosis and treatment, combined with vector control (Boelaert *et al.* 2000).

Vector control

Indoor residual spraying (IRS) is the most widely used intervention and is implemented routinely in India and Nepal by spraying houses twice yearly in villages where a case of VL had been reported in the year before. IRS has been shown to be effective in reducing sandfly density in India and Nepal where the sandfly (*Phlebotomus argentipes*) is endophilic and rests inside cracks and crevices of mud plastered walls (Kaul *et al.* 1994; Kumar *et al.* 2009; Das *et al.* 2008). Although there are no studies that looked at the effects of IRS on VL transmission and incidence, the large scale spraying campaigns organized in the 1950s in an effort to eradicate malaria also reduced kala-azar incidence in the region (Birley *et al.* 1993). VL subsequently re-emerged in the early eighties with the discontinuation of the spraying activities (Bern &

Chowdhury 2006). In Sudan, and eastern Africa, the sandfly species is different (*Phlebotomus orientalis*) and transmission occurs both in villages (Hassan *et al.* 2004; Elnaiem & Osman 1998) and outside in *Acacia/Balanites* forests (Elnaiem *et al.* 1998b; Thomson *et al.* 1999). Because individuals are exposed differently to sandflies, e.g. by sleeping outside during the dry season (Ritmeijer *et al.* 2007) or through activities such as shepherding (Hoogstraal & Heyneman 1969; Chappuis *et al.* 2007) IRS is thought to be less efficacious and no spraying activities are currently taking place. Insecticide treated nets (ITN) have been distributed in Sudan by Médecins sans Frontières. While ITNs are also distributed in India and Nepal, the evidence on the effectiveness of insecticide treated nets (ITN) is mixed. Although some studies showed sleeping under a bednet to be a protective factor against VL (e.g. Bern *et al.* 2000; Bern *et al.* 2005), recent evidence from a community intervention trial of LLIN in the Indian subcontinent suggests a 25% reduction in vector density but not leading to reduced VL incidence compared to control areas where untreated bednet use was high (Picado *et al.* 2010).

Case detection and management

Diagnosis

The clinical diagnosis of VL is difficult because the signs and symptoms are not characteristic to VL only and may be confused with other diseases such as malaria, schistosomiasis or typhoid fever (Singh 2006). Microscopic confirmation of parasites in aspirates of lymph nodes, bone marrow or spleen is the recommended method of diagnosis for VL (Chappuis *et al.* 2007; WHO 2010). However, the sensitivity of microscopic examination varies. Splenic aspiration has the highest sensitivity, but is an invasive and complex technique and life-threatening bleeding can occur (Boelaert *et al.* 2007). The rK39 immunochromatographic test (rK39) and the Direct Agglutination Test (DAT) have now largely replaced these tests (Sundar *et al.* 2002b). Both the rK39 and DAT are recommended by the VL elimination initiative in India and Nepal (WHO 2005) and national guidelines in Sudan. However, the rK39 seems to be less sensitive in Sudan compared to India and Nepal (Ritmeijer *et al.* 2006) and in case of a negative test result, a second test is still required.

Treatment strategies

Treatment options for VL are limited and treatment policies vary across countries and regions. There are currently five drugs available for the treatment of VL. Pentavalent antimonials (sodium stibogluconate and meglumine antimoniate) have been the first line treatment for VL for over 70 years. They exist in both branded (Pentostam®, Glucantime®) and generic forms (produced by Albert David Ltd. in India). Generic antimonials are as effective as their branded counterparts but are up to 4 times cheaper (Veeken *et al.* 2000; Moore *et al.* 2001; Ritmeijer *et al.* 2001). The recommended treatment schedule for antimonials is 20 mg/kg/day for 20

to 30 days and is usually administered intramuscularly on a daily basis. Antimonials are toxic drugs with frequent adverse side effects such as cardio-, hepato- and nephrotoxicity (Ballou *et al.* 1987). In eastern Africa, antimonials are still the first line treatment for VL and patients are usually admitted for the full duration of treatment to monitor side effects and ensure adherence. In India and Nepal, antimonials are no longer recommended due to increasing reports of treatment failure (Lira *et al.* 1999; Rijal *et al.* 2003; Sundar 2001; Croft *et al.* 2006). Nonetheless a recent study showed they continued to be used in peripheral health facilities in Bihar (Hasker *et al.* 2010).

Conventional amphotericin B deoxycholate has still very high cure rates in the Indian subcontinent and is now most often used as second-line treatment (Sundar *et al.* 2004; Thakur *et al.* 1999). The drug is administered intravenously with 15–20 slow infusions of 1 mg/kg given either on a daily or alternate basis (Sundar *et al.* 2004). Amphotericin B shares drawbacks with antimonials such as the length of therapy (28–30 days) and parental administration. Because of its potential toxicity and mode of administration, hospitalization is required for the full duration of treatment (Murray 2000).

Three other drugs have been approved for the treatment of VL, but the efficacy and required dosage have been shown to differ between the Indian subcontinent and East Africa (WHO 2010). Liposomal Amphotericin B (Ambisome; Gilead), a lipid formulation of amphotericin B, is considered the best monotherapy for anthroponotic VL (WHO 2010). A single dose of liposomal amphotericin B delivered intravenously, was shown to be very effective in the Indian subcontinent and resulted in less side-effects compared to other anti-leishmanial treatments (Sundar *et al.* 2010; Thakur 2001). The very high price has prevented its widespread use as first line treatment but preferential pricing agreements negotiated by the WHO (Matlashewski *et al.* 2011) and a recent donation of 445,000 vials to treat 50,000 patients in the coming 5 years may change this (WHO 2011). Miltefosine is the first oral drug for VL and was originally developed as an anti-cancer drug. The drug is administered orally for 28 days, but is possibly teratogenic and cannot be administered to pregnant women or women in childbearing age that refuse the use of contraceptives (Sundar *et al.* 2002a). Despite these limitations, it was adopted as first line treatment by the VL elimination initiative in the Indian subcontinent. Aminosidine (paromomycin), an antibiotic with good anti-leishmanial activity has been registered in India since 2006. It is administered intramuscularly for 21 days, has few side effects and is very cheap. At a dosage of 15mg per kg per day it has been shown to be effective in India (Sundar *et al.* 2007), but not in East Africa in a phase II trial (Musa *et al.* 2010). Both miltefosine and paromomycin have a long half-life and are suspected to rapidly generate resistance (Sundar & Murray 2005; Perez-Victoria *et al.* 2006; Seifert *et al.* 2007). An overview of the current

monotherapies, their clinical efficacy, toxicity and disadvantages are presented in table 1 of Chapter 7.

As there are no new compounds for VL expected to come to the market in the near future, combination therapy, as used successfully against malaria and tuberculosis, is increasingly being considered as a possibility to delay or prevent the emergence of resistance to the currently available VL drugs. Combination therapies may also increase tolerability, reduce treatment duration and possibly (direct and indirect) costs. Médecins sans Frontières in southern Sudan already used a combination of sodium stibogluconate with paromomycin during an epidemic (Ritmeijer and Davidson 2003). For a complete overview and discussion of combination therapies we refer to Chapter 7.

VL: A NEGLECTED DISEASE

The World Health Organization (WHO) considers VL as a neglected tropical disease (NTD). The NTDs are a group of chronic vector-borne protozoan, helminthic and bacterial infections including VL and other diseases such as hookworm infections, schistosomiasis, lymphatic filariasis, African trypanosomiasis, buruli ulcer or leprosy (WHO 2012). These diseases affect almost exclusively poor rural populations and are considered “neglected” as they are under-researched and lack adequate funding for their control, although recently a number of diseases have been given higher priority by the international donor community. Together, the NTDs affect more than a billion people worldwide and cause the loss of 57 million disability adjusted life-years (DALYs), a burden higher than tuberculosis or malaria (Hotez *et al.* 2006b). They are an important cause of disability (e.g. lymphatic filariasis), disfigurement and stigma (e.g. cutaneous leishmaniasis) (Beyrer *et al.* 2007; Perera *et al.* 2007), blindness (onchocerciasis) (Boatin & Richard 2006), anaemia and malnutrition (e.g. schistosomiasis, hookworm infections) (King *et al.* 2005) and death in developing countries (e.g. schistosomiasis, ascariasis, visceral leishmaniasis) (Hotez *et al.* 2006a; Alvar *et al.* 2012). A common feature of the NTDs is their inextricable link with poverty. On the one hand NTDs often cluster in the same geographical areas, primarily in low-income countries, where the poor and disadvantaged are most at risk of infection with one or more NTDs due to poor living and working conditions, inadequate sanitation or lack of access to safe drinking water and health care. On the other hand, the NTDs are also an important determinant of poverty through their effect on child development, the reduced productivity of infected individuals and the costs related to accessing treatment services. For example children infected with schistosomiasis or hookworms are more absent from school, perform less well and have diminished cognitive abilities (De Clercq *et al.* 1998; Sakti *et al.* 1999; Jukes *et al.* 2002), which may impact their educational achievements and lead to

lower wage-earning potentials in the future (Hotez *et al.* 2009). The physical impairments, such as the swelling of the legs caused by lymphatic filariasis (i.e. elephantiasis), prevents infected individuals from working in the fields or performing other productive activities (Ramaiah *et al.* 2000; Gyapong *et al.* 2000) which has a negative impact on the income of the household. The stigma caused by diseases such as lymphatic filariasis, or the disfigurement from lesions to the face and visible parts of the body in the case of cutaneous leishmaniasis often leads to physical and emotional isolation (Reitinger *et al.* 2005) or even exclusion from community life (Wiess 2008). Other NTDs, like visceral leishmaniasis, have a direct effect on household wealth through catastrophic health expenditures on treatment and sales of livestock or assets to pay for the costs of care (Adhikary *et al.* 2009; Sharma *et al.* 2006; Meheus *et al.* 2006).

OBJECTIVES AND OUTLINE OF THE THESIS

Objectives

Visceral leishmaniasis disproportionally affects the poorest population groups in low-income countries. For the control, or even elimination, of VL to be successful, it will require a great deal of effort that needs to be sustained over the longer term. Ultimately, effective control of VL will depend on a mix of interventions, consisting of early and efficient diagnosis and treatment, with drug regimens that put as little burden as possible on both the patient and the health system. However, the scarcity of reliable data, both epidemiological and economic, contributes to the low visibility and priority given to VL by national control programmes and international donors.

The general objective of this thesis was to improve our understanding on the economic aspects of visceral leishmaniasis, provide evidence for more rational decision making on diagnosis, case management and VL control and justify increased investment in VL research and control in the Indian subcontinent and East Africa.

More specifically, the studies included in this thesis were framed around the following specific objectives (figure 2):

- a. To understand the burden of visceral leishmaniasis in India, Nepal and Sudan;
- b. To provide a better understanding of the health seeking behaviour and costs of VL illness from the provider and household perspective;
- c. To examine the cost and cost-effectiveness of VL treatment alternatives with a focus on combination therapies.

The studies were carried out in collaboration with research partners in the three countries, in particular the BP Koirala Institute of Health Sciences (Dharan, Nepal), the Institute of Medical Sciences, Banaras Hindu University (Varanasi, India) and the Institute of Endemic Diseases, University of Khartoum (Khartoum, Sudan). International partners involved in the studies include the Drugs for Neglected Diseases Initiative and the World Health Organization (Geneva, Switzerland).




Figure 2: Overview of the thesis by research questions (RQ)

Outline of the thesis

The thesis is divided into three sections structured around the specific objectives described above that examine different aspects in the field of health economics applied to VL. In the first section on the burden of VL we present two studies. *Chapter 2* provides an overview of studies that investigated the socio-economic aspects of two neglected tropical diseases: visceral leishmaniasis and sleeping sickness. Both of these diseases affect the poorest of the poor in endemic countries, cause considerable direct and indirect costs (even though the national control programmes tend to provide free care) and push affected households deeper into poverty. In *chapter 3* we compared the socio-economic profile of households in areas with high VL endemicity to the

general population in Bihar State in India. The aim was to assess whether VL truly affects the poorest of the poor, a statement that is often made but had never been substantiated before. The data for this comparison were obtained from a community intervention trial in VL-endemic areas and the National Family Health Survey of India. In the second section of the thesis we estimated the cost-of-illness of VL In India, Nepal and Sudan. In India (*chapter 4*) we examined the economic costs (direct and indirect) of VL treatment from the societal and household perspective at a charitable hospital in Bihar, India. At the time, it was the first economic study on VL to be conducted in India. In Nepal (*chapter 5*) we carried out a household survey in 5 districts in Nepal to assess the economic impact of VL on households and examine whether the intensified VL control efforts induced by the VL elimination initiative resulted in a decrease in household costs. It was the largest study of its kind in Nepal to systematically estimate the cost incurred by households at all health providers, both public and private, when seeking VL care. In Sudan (*chapter 6*) we estimated the cost of providing VL diagnosis and treatment services (i.e. health-care provider perspective) and the cost to patients and their family to access these services (i.e. household perspective) at three health care facilities in Gedaref State. Besides costs, these studies also examined health seeking behaviour, strategies to cope with the costs of illness and the extent of catastrophic health expenditures households incurred. In the third and last section of the thesis we carried out a review on the current evidence and potential of combination therapies (*chapter 7*), while in *chapter 8* we analysed the cost and cost-effectiveness of treatment strategies for VL in the Indian subcontinent with a focus on combination therapies. The final *chapter 9* discusses our findings.

REFERENCES

- Addy M, Nandy A (1992) Ten years of kala-azar in west Bengal, Part I. Did post-kala-azar dermal leishmaniasis initiate the outbreak in 24-Parganas? *Bull World Health Organ* 70: 341–46.
- Adhikari SR, Maskay NM, Sharma BP (2009) Paying for hospital-based care of Kala-azar in Nepal: assessing catastrophic, impoverishment and economic consequences. *Health Policy Plan* 24: 129–39.
- Alvar J, Croft S, Olliaro P (2006) Chemotherapy in the treatment and control of leishmaniasis. *Adv Parasitol* 61: 223–74.
- Alvar J, Vélez ID, Bern C, et al. (2012) Leishmaniasis Worldwide and Global Estimates of Its Incidence. *PLoS ONE* 7: e35671.
- Ballou WR, McClain JB, Gordon DM, et al. (1987) Safety and efficacy of high-dose sodium stibogluconate therapy of American cutaneous leishmaniasis. *Lancet* 2: 13–16.
- Berman JD (1988) Chemotherapy for visceral leishmaniasis: biochemical mechanisms, clinical efficacy, and future strategies. *Rev Infect Dis* 10: 560–86.
- Bern C, Chowdhury R (2006) The epidemiology of visceral leishmaniasis in Bangladesh: prospects for improved control. *Indian J Med Res* 123: 275–88.
- Bern C, Hightower AW, Chowdhury R, et al. (2005) Risk factors for kala-azar in Bangladesh. *Emerg Infect Dis* 11: 655–62.
- Bern C, Joshi AB, Jha SN, et al. (2000) Factors associated with visceral leishmaniasis in Nepal: bed-net use is strongly protective. *Am J Trop Med Hyg* 63: 184–88.
- Bern C, Maguire JH, Alvar J (2008) Complexities of Assessing the Disease Burden Attributable to Leishmaniasis. *PLoS Negl Trop Dis* 2: e313.
- Beyrer C, Villar JC, Suwanvanichkij V, Singh S, Baral SD, Mills EJ (2007) Health and human rights 3: neglected diseases, civil conflicts, and the right to health. *Lancet* 370: 619–27.
- Birley MH (1993) An historical review of malaria, kala-azar and filariasis in Bangladesh in relation to the Flood Action Plan. *Ann Trop Med Parasitol* 87: 319–34.
- Boatin BA, Richards FO Jr (2006) Control of onchocerciasis. *Adv Parasitol* 61: 349–94.
- Boelaert M, Bhattacharya S, Chappuis F, et al. (2007) Evaluation of rapid diagnostic tests: visceral leishmaniasis. *Nat Rev Microbiol* S30–S39.
- Boelaert M, Criel B, Leeuwenburg J, Van Damme W, Le Ray D, Van der Stuyt P (2000) Visceral leishmaniasis control: a public health perspective. *Trans R Soc Trop Med Hyg* 94: 465–71.
- Bryceson A (2001) A policy for leishmaniasis with respect to the prevention and control of drug resistance. *Trop Med Int Health* 6: 928–34.
- Burki T (2009) East African countries struggle with visceral leishmaniasis. *Lancet* 374: 371–72.
- Chappuis F, Rijal S, Soto A, Menten J, Boelaert M (2006) A meta-analysis of the diagnostic performance of the direct agglutination test and rK39 dipstick for visceral leishmaniasis. *BMJ* 333: 723.

- Chappuis F, Sundar S, Hailu A, et al. (2007) Visceral leishmaniasis: what are the needs for diagnosis, treatment and control? *Nat Rev Microbiol* 5: 873–82.
- Collin SM, Coleman PG, Ritmeijer K, Davidson RN (2006) Unseen Kala-azar deaths in south Sudan (1999–2002). *Trop Med Int Health* 11:509–12.
- Croft SL, Sundar S, Fairlamb AH (2006) Drug resistance in leishmaniasis. *Clin Microbiol Rev* 19: 111–26.
- Das M, Banjara M, Chowdhury R, et al. (2008) Visceral leishmaniasis on the Indian sub-continent: a multi-centre study of the costs of three interventions for the control of the sandfly vector, *Phlebotomus argentipes*. *Ann Trop Med Parasitol* 102: 729–41.
- de Clercq D, Sacko M, Behnke J, Gilbert F, Vercruysse J (1998) The relationship between *Schistosoma haematobium* infection and school performance and attendance in Bamako, Mali. *Ann Trop Med Parasitol* 92: 851–58.
- Dereure J, El-Safi SH, Bucheton B, et al. (2003) Visceral leishmaniasis in eastern Sudan: parasite identification in humans and dogs; host-parasite relationships. *Microbes Infect* 5: 1103–08.
- Desjeux P (1992) Human leishmaniasis: epidemiology and public health aspects. *World Health Stat Q* 45: 267–75.
- Desjeux P (1996) Leishmaniasis. Public health aspects and control. *Clin Dermatol* 14: 417–23.
- Desjeux P (2001) The increase in risk factors for leishmaniasis worldwide. *Trans R Soc Trop Med Hyg* 95: 239–43.
- Desjeux P (2004) Focus: Leishmaniasis. *Nat Rev Microbiol* 2: 692–93.
- Dey A, Singh S (2006) Transfusion transmitted leishmaniasis: a case report and review of literature. *Indian J Med Microbiol* 24:165–70.
- Elnaiem DA, Ward RD, Hassan HK, Miles MA, Frame IA (1998a) Infection rates of *Leishmania donovani* in *Phlebotomus orientalis* from a focus of visceral leishmaniasis in eastern Sudan. *Ann Trop Med Parasitol* 92: 229–32.
- Elnaiem DA, Connor SJ, Thomson MC, et al. (1998b) Environmental determinants of the distribution of *Phlebotomus orientalis* in Sudan. *Ann Trop Med Parasitol* 92: 877–87.
- Elnaiem DA, Hassan MM, Maingon R, et al. (2001) The Egyptian mongoose, *Herpestes ichneumon*, is a possible reservoir host of visceral leishmaniasis in eastern Sudan. *Parasitology* 122: 531–536.
- Elnaiem DA, Osman OF (1998) Evidence for active transmission of visceral leishmaniasis within a village in eastern Sudan. *Acta Trop* 71: 305–09.
- Gyapong M, Gyapong J, Weiss M, Tanner M (2000) The burden of hydrocele on men in Northern Ghana. *Acta Trop* 77: 287–94.
- Hasker E, Singh SP, Malaviya P, et al. (2010) Management of visceral leishmaniasis in rural primary health care services in Bihar India. *Trop Med Int Health* 15 (S2): 55–62.
- Hassan MM, Elraba'a FM, Ward RD, Maingon RD, Elnaiem DA (2004) Detection of high rates of in-village transmission of *Leishmania donovani* in eastern Sudan. *Acta Trop* 92: 77–82.
- Hoogstraal H, Heyneman D (1969) Leishmaniasis in the Sudan Republic: 30. Final Epidemiologic Report. *Am J Trop Med Hyg* 18: 1091–210.
- Hotez P, Ottesen E, Fenwick A, Molyneux D (2006a) The neglected tropical diseases: the ancient afflictions of stigma and poverty and the prospects for their control and elimination. *Adv Exp Med Biol* 582: 23–33.
- Hotez PJ, Molyneux DH, Fenwick A, Ottesen E, Sachs SE, Sachs JD (2006b) Incorporating a rapid-impact package for neglected tropical diseases with programs for HIV/AIDS, tuberculosis, and malaria. *PLoS Med* 3: e102.
- Hotez PJ, Fenwick A, Savioli L, Molyneux DH (2009) Rescuing the bottom billion through control of neglected tropical diseases. *Lancet* 373: 1570–05.
- Jukes MC, Nokes CA, Alcock KJ, et al. (2002) Partnership for child development. Heavy schistosomiasis associated with poor short-term memory and slower reaction times in Tanzanian schoolchildren. *Trop Med Int Health* 7: 104–17.
- Kaul SM, Sharma RS, Dey KP, Rai RN, Verghese T (1994) Impact of DDT indoor residual spraying on *Phlebotomus argentipes* in a kala-azar endemic village in eastern Uttar Pradesh. *Bull World Health Organ* 72: 79–81.
- King CH, Dickman K, Tisch DJ (2005) Reassessment of the cost of chronic helminthic infection: a meta-analysis of disability-related outcomes in endemic schistosomiasis. *Lancet* 365: 1561–69.
- Kumar V, Kesari S, Dinesh DS, et al. (2009) A report on the indoor residual spraying (IRS) in the control of *Phlebotomus argentipes*, the vector of visceral leishmaniasis in Bihar (India): an initiative towards total elimination targeting 2015 (Series-1). *J Vector Borne Dis* 46: 225–29.
- Lira R, Sundar S, Makharia A, et al. (1999) Evidence that the high incidence of treatment failures in Indian kala-azar is due to the emergence of antimony-resistant strains of *Leishmania donovani*. *J Infect Dis* 180: 564–67.
- Matlashewski G, Arana B, Kroeger A, et al. (2011) Visceral leishmaniasis: elimination with existing interventions. *Lancet Infect Dis* 11: 322–25.
- Médecins sans Frontières (2010) Southern Sudan in grips of worst disease outbreak in eight years. Press release Juba, Dec. 16 2010. Available at: <http://www.doctorswithoutborders.org/press/release.cfm?id=4924&cat=press-release> (Accessed on June 15, 2012)
- Meheus F, Boelaert M, Baltussen R, Sundar S (2006) Costs of patient management of visceral leishmaniasis in Muzaffarpur Bihar, India. *Trop Med Int Health* 11: 1715–24.
- Moore E, O'Flaherty D, Heuvelmans H, et al. (2001) Comparison of generic and proprietary sodium stibogluconate for the treatment of visceral leishmaniasis in Kenya. *Bull World Health Organ* 79: 388–93.
- Murray HW (2000) Treatment of visceral leishmaniasis (Kala-Azar): a decade of progress and future approaches. *Int J Infect Dis* 4: 158–77.
- Musa AM, Younis B, Fadlalla A, et al. (2010) Paromomycin for the Treatment of Visceral Leishmaniasis in Sudan: A Randomized, Open-Label, Dose-Finding Study. *PLoS Negl Trop Dis* 4: e855.
- Pearson RD, de Queiroz Sousa A (1996) Clinical spectrum of Leishmaniasis. *Clin Infect Dis* 22: 1–13.

- Perera M, Whitehead M, Molyneux D, Weerasooriya M, Gunatilleke G (2007) Neglected patients in neglected disease? A qualitative study of lymphatic filariasis. *PLoS Negl Trop Dis* 1: e128.
- Perez-Victoria FJ, Sánchez-Cañete MP, Seifert K, et al. (2006) Mechanisms of experimental resistance of *Leishmania* to miltefosine: implications for clinical use. *Drug Resist Updat* 9: 26–39
- Picado A, Singh SP, Rijal S, et al. (2010) Long-lasting insecticidal nets for prevention of *Leishmania donovani* infection in India and Nepal: paired cluster randomised trial. *BMJ* 341: c6760
- Ramaiah KD, Radhamani MP, John KR, et al. (2000) The impact of lymphatic filariasis on labour inputs in southern India: results of a multi-site study. *Ann Trop Med Parasitol* 94: 353–63.
- Reithinger R, Aadil K, Kolaczinski J, Mohsen M, Hami S (2005) Social impact of leishmaniasis, Afghanistan. *Emerg Infect Dis* 11: 634–36.
- Rijal S, Chappuis F, Singh R, et al. (2003) Treatment of visceral leishmaniasis in south-eastern Nepal: decreasing efficacy of sodium stibogluconate and need for a policy to limit further decline. *Trans R Soc Trop Med Hyg* 97: 350–354.
- Ritmeijer K, Davidson RN (2003) Royal Society of Tropical Medicine and Hygiene joint meeting with Médecins Sans Frontières at Manson House, London, 20 March 2003: field research in humanitarian medical programmes. Médecins Sans Frontières interventions against kala-azar in the Sudan, 1989–2003. *Trans R Soc Trop Med Hyg* 97: 609–13.
- Ritmeijer K, Davies C, van Zorge R, et al. (2007) Evaluation of a mass distribution programme for fine-mesh impregnated bednets against visceral leishmaniasis in eastern Sudan. *Trop Med Int Health* 12: 404–14.
- Ritmeijer K, Melaku Y, Mueller M, Kipnetich S, O'keeffe C, Davidson RN (2006) Evaluation of a new recombinant K39 rapid diagnostic test for Sudanese visceral leishmaniasis. *Am J Trop Med Hyg* 74: 76–80.
- Ritmeijer K, Veeken H, Melaku Y, et al. (2001) Ethiopian visceral leishmaniasis: generic and proprietary sodium stibogluconate are equivalent; HIV co-infected patients have a poor outcome. *Trans R Soc Trop Med Hyg* 95: 668–72.
- Sakti H, Nokes C, Hertanto WS, et al. (1999) Evidence for an association between hookworm infection and cognitive function in Indonesia school children. *Trop Med Int Health* 4: 322–34.
- Sanyal RK (1985) *Leishmaniasis in the Indian subcontinent*. In: Chang KP, Bray RS (eds.) Leishmaniasis. Amsterdam: Elsevier Science Publishers. pp. 443–467.
- Seaman J, Mercer AJ, Sondorp E (1996) The epidemic of visceral leishmaniasis in Western Upper Nile, Southern Sudan: course and impact from 1984 to 1994. *Int J Epidemiol* 25: 862–71.
- Seifert K, Matu S, Perez-Victoria J, Castanys S, Gamarro F, Croft S (2003) Characterisation of *Leishmania donovani* promastigotes resistant to hexadecylphosphocholine (miltefosine). *Int J Antimicrob Agents* 22: 380–87.
- Seifert K, Perez-Victoria FJ, Stettler M, et al. (2007) Inactivation of the miltefosine transporter, LdMT, causes miltefosine resistance that is conferred to the amastigote stage of *Leishmania donovani* and persists in vivo. *Int J Antimicrob Agents* 30: 229–35.
- Sharma A, Bern C, Varghese B, et al. (2006) The economic impact of visceral leishmaniasis on households in Bangladesh. *Trop Med Int Health* 11: 757–64.
- Singh P, Reddy DC, Rai M, Sundar S (2006) Serious underreporting of visceral leishmaniasis through passive case reporting in Bihar, India. *Trop Med Int Health* 11: 899–905
- Singh S (2006) New developments in diagnosis of leishmaniasis. *Indian J Med Res* 123: 311–30.
- Singh S, Chaudhry VP, Wali JP (1996) Transfusion-transmitted kala-azar in India. *Transfusion* 36: 848–49.
- Singh VP, Ranjan A, Topno RK, et al. (2010) Estimation of under-reporting of visceral leishmaniasis cases in Bihar, India. *Am J Trop Med Hyg* 82: 9–11
- Sundar S (2001) Drug resistance in Indian visceral leishmaniasis. *Trop Med Int Health* 6: 849–54.
- Sundar S, Chakravarty J, Agarwal D, Rai M, Murray HW (2010) Single-dose liposomal amphotericin B for visceral leishmaniasis in India. *N Engl J Med* 362: 504–12.
- Sundar S, Jha TK, Thakur CP, et al. (2002a) Oral miltefosine for Indian visceral leishmaniasis. *N Engl J Med* 347: 1739–46.
- Sundar S, Sahu M, Mehta H, et al. (2002b) Non-invasive management of Indian visceral leishmaniasis: clinical application of diagnosis by K39 antigen strip testing at a kala-azar referral unit. *Clin Infect Dis* 35: 581–86.
- Sundar S, Jha TK, Thakur CP, Sinha PK, Bhattacharya SK (2007) Injectable paromomycin for Visceral leishmaniasis in India. *N Engl J Med* 356: 2571–81.
- Sundar S, Mehta H, Suresh AV, Singh SP, Rai M, Murray HW (2004) Amphotericin B treatment for Indian visceral leishmaniasis: conventional versus lipid formulations. *Clin Infect Dis* 38: 377–83.
- Sundar S, Murray HW (2005) Availability of miltefosine for the treatment of kala-azar in India. *Bull World Health Organ* 83: 394–95.
- Thakur CP (2001) A single high dose treatment of kala-azar with Ambisome (amphotericin B lipid complex): a pilot study. *Int J Antimicrob Agents* 17: 67–70.
- Thakur CP, Singh RK, Hassan SM, Kumar R, Narain S, Kumar A (1999) Amphotericin B deoxycholate treatment of visceral leishmaniasis with newer modes of administration and precautions: a study of 938 cases. *Trans R Soc Trop Med Hyg* 93: 319–23.
- Thomson MC, Elnaïem DA, Ashford RW, Connor SJ (1999) Towards a kala-azar risk map for Sudan: mapping the potential distribution of *Phlebotomus orientalis* using digital data of environmental variables. *Trop Med Int Health* 4: 105–13.
- Uranw S, Meheus F, Baltussen R, Rijal S, Boelaert M (2013) The economic burden of visceral leishmaniasis on households in eastern Nepal. *PLoS Negl Trop Dis* 7: e2062
- Veeken H, Ritmeijer K, Seaman J, Davidson R (2000) A randomized comparison of branded sodium stibogluconate and generic sodium stibogluconate for the treatment of visceral leishmaniasis under field conditions in Sudan. *Trop Med Int Health* 5: 312–17.
- Weiss MG (2008) Stigma and the Social Burden of Neglected Tropical Diseases. *PLoS Negl Trop Dis* 2: e237
- World Health Organization (2011) WHO and Gilead sign agreement for enhanced access to visceral leishmaniasis treatment. Dec 8. Available: http://www.who.int/neglected_diseases/Gilead_donation_2011/en/index.html (Accessed on 14/06/2012).

World Health Organization (2005) *Regional strategic framework for elimination of kala-azar from the South-East Asia region (2005–2015)*. New Delhi: WHO Regional Office for South-East Asia.

World Health Organization (2010) *Control of the Leishmaniasis. Report of a meeting of the WHO Expert Committee on the Control of the Leishmaniasis*. Geneva: World Health Organization.

World Health Organization (2012) *Neglected Tropical Diseases*. Available online: http://www.who.int/neglected_diseases/diseases/en/. Accessed online on July 2, 2012.


Zijlstra EE, el-Hassan AM (2001) Leishmaniasis in Sudan. Post kala-azar dermal leishmaniasis. *Trans R Soc Trop Med Hyg* 95(S1): S59–76.

Zijlstra EE, Musa AM, Khalil EA, el-Hassan IM, el-Hassan AM (2003) Post kala-azar dermal leishmaniasis. *Lancet Infect Dis* 3: 87–98.



PART 1

The burden of visceral leishmaniasis



CHAPTER 2

Socio-economic aspects of neglected diseases: sleeping sickness and visceral leishmaniasis

Boelaert M,
Meheus F
Robays J
Lutumba P

ABSTRACT

Several tropical diseases that are essentially poverty-related have recently gained more attention under the label of ‘neglected tropical diseases’ or NTD. It is estimated that over 1000 million people currently suffer from one or more NTD. Here, the socio-economic aspects of two NTD— human African trypanosomiasis and human visceral leishmaniasis — are reviewed. Both of these diseases affect the poorest of the poor in endemic countries, cause considerable direct and indirect costs (even though the national control programmes tend to provide free care) and push affected households deeper into poverty.

The neglected tropical diseases of humans (NTD) include a wide range of mainly, but not exclusively, parasitic diseases that can be usefully considered as a group because they are concentrated almost entirely among the poorest populations in the least-developed countries. Recent advocacy campaigns led the international donor community to give higher priority to these neglected health issues that, taken together, contribute a substantial part of the burden of disease in the tropics (Holveck *et al.* 2007). As NTD are so strongly related to poverty, an understanding of the socio-economic context is essential for their control. The available information on the socio-economic aspects of sleeping sickness and visceral leishmaniasis, two vector-borne diseases that are intrinsically poverty-related, is reviewed below.

SLEEPING SICKNESS

In the fields of research and control, sleeping sickness or human African trypanosomiasis (HAT) is a typical example of a ‘neglected disease’ (Trouiller *et al.* 2001, 2002). The numerous scientific articles on the unique immunological properties of trypanosomes contrast with the few papers that have been published on improvements in the diagnosis, treatment and control of HAT.

The large HAT outbreaks observed in colonial times have been attributed to forced displacement and deterioration in living conditions (Welburn *et al.* 2004). As HAT posed a threat to the economic development of large areas, the colonial authorities did invest in research on the diseases and the development of control tools. Large campaigns for the (mandatory) screening of populations were set up, and by the 1960s the disease had almost disappeared from Central Africa. Control methods have not fundamentally changed since this colonial period, although screening is no longer mandatory. The 1990s brought the progressive introduction of a serological test for mass screening —the card agglutination test for trypanosomiasis (CAT; Magnus *et al.* 1978) — and, more recently, eflornithine (Nightingale 1991) or a nifurtimox-eflornithine combination have been proposed, as alternatives to melarsoprol, for the treatment of stage-2 sleeping sickness (Priotto *et al.* 2009). The case-management strategy still remains complex, however, and much uncertainty surrounds the effectiveness of tsetse control.

The funding of HAT control currently depends nearly entirely on international aid. Interruptions in control efforts, linked to civil unrest in the affected areas, have led to spectacular re-emerging epidemics in recent years, in Sudan (Moore *et al.* 1999), Angola (Stanghellini & Josenando 2001) and the Democratic Republic of Congo (DRC; Lutumba *et al.* 2005). Although the global incidence of HAT seems to be declining, several non-governmental organizations (NGO) have warned that claims that HAT could be eliminated in the near future are probably premature,

given the 'hidden epidemic' in regions affected by civil war, in north-eastern DRC, southern Sudan and the Central African Republic (Chappuis *et al.* 2010).

The high burden that HAT places on affected households and communities is often not very visible in national or regional health data because of its focal nature. Within HAT foci, prevalence is often around 1% and, in the absence of control, it can rise relatively rapidly, sometimes to 50% in certain villages. Thus, much of the burden of this disease often falls, very heavily, on a few locations (Lutumba *et al.* 2007). The direct effects of sleeping sickness are mainly felt by economically active adults, and the cases require a great deal of intensive care, often from the other adults in their families. The seeking of a diagnosis and treatment are costly and time-consuming, and the economic burden on the household of an HAT case is high — between 1.5 and 10 months of household income, even when the diagnostics and anti-trypanosomal drugs are provided free of charge (Lutumba *et al.* 2007).

There have been few attempts to quantify the full costs borne by households with HAT cases, which include the costs of care at home and during hospital treatment, the costs of seeking a diagnosis, income lost by the HAT cases and their carers, medical fees, and the costs of transport. In the Republic of Congo, a case of HAT who was correctly diagnosed and treated was found to cost his or her household the equivalent of 2.6–5 months of household income from agriculture (Gouteux *et al.* 1987). In their more recent study in the DRC, where the direct costs of screening and treatment are fully subsidized by a national HAT programme, Lutumba *et al.* (2007) estimated the mean household costs for a diagnosed and treated case of HAT to be 5 months of household income, with such costs rising to over 10 months of household income for a case with complications. The costs of HAT treatment that are borne by households may be so high that they inhibit the timely uptake of treatment. When they discover that they have cases of HAT, households often take their time to prepare themselves financially and mobilize the necessary resources (often relying on the solidarity of the extended family), before the cases present themselves for treatment. High household costs (or, rather, the fear of being identified as an HAT case and then having to bear such expense) may partly explain the low levels of participation often seen at active-screening sessions organized by mobile teams (Robays *et al.* 2004).

There have been few published studies on the economics of controlling HAT, and fewer still on the comparative cost-effectiveness of vector control, at least in the context of dealing with HAT rather than trypanosomiasis in livestock. The costs of vector control to reduce the transmission of the parasites causing HAT have been estimated as U.S.\$17,400/25,000 inhabitants over 3

years (Gouteux & Sinda 1990), U.S.\$0.90/person-year (Lancien 1991), or 2000 CFA francs/person in year 1 and 250 CFA francs/person-year subsequently (Laveissière & Couret 1986). The costs of various approaches to active case-finding and treatment have been studied in several contexts and areas (Shaw & Cattand 2001), including the DRC (Lutumba *et al.* 2007). When Shaw (1989) compared the costs of vector control with the finding and treatment of human cases, she found active case-finding to be the more cost-effective option. The assumptions that Shaw (1989) made on the effectiveness of such case-finding were, however, probably optimistic compared with the trends observed in real-life settings (Robays *et al.* 2004).

VISCERAL LEISHMANIASIS

The World Health Organization estimates the incidence of human visceral leishmaniasis (VL) at 500,000 new cases/year (Murray & Lopez 1996¹), with 90% of cases occurring in just four countries: India, Bangladesh, Brazil and Sudan (Desjeux 1996).

Desjeux (1996) characterised VL patients as '*those of lowest socio-economic status, who have minimal political power to influence the decisions of the government and a very limited capacity to assume the costs of the disease.*' At country level, leishmaniasis is often a hidden problem, as cases usually live in remote areas with poor access to services, and case loads are poorly documented, with five to eight unreported cases for every case that is reported (Singh *et al.* 2006, 2010b; Das *et al.* 2010). As VL, like HAT, often has a very focal distribution, the national (country aggregated) figures on VL incidence often fail to reflect the heavy burden on the worst affected communities. The 'hyper-clustering' of VL cases has recently been described in Nepal (Rijal *et al.* 2010) and India (Singh *et al.* 2010a), where the incidences in endemic clusters have been found to be 10 times higher than those reported at the corresponding district level.

In industrialized countries, the leishmaniasis of humans were not perceived as a direct threat until their emergence as a co-infection in HIV/AIDS, and, since the number of cases seen annually in the Mediterranean region has declined (as the result of widespread use of highly active antiretroviral therapy), such diseases seem to have disappeared again from the local public-health agenda (Dujardin *et al.* 2008).

The importance of VL as a hindrance to economic development is still not fully recognised although there is little doubt that the disease is poverty-related (Alvar *et al.* 2006). In Bihar state, one of the economically most deprived states in India, 80% of the families in VL-affected villages were found to belong to the poorest two quintiles in terms of wealth distribution

¹ www.who.int/leishmaniasis/burden/magnitude/burden_magnitude/en/index.html

(Boelaert *et al.* 2009). The characterisation of a typical VL case as the ‘poorest of the poor’ should therefore be taken fairly literally.

VL is a deadly disease if left untreated and one that can have a disastrous impact when it strikes a non-immune population. In Sudan, a VL epidemic caused major disruption to a famine-stricken society when, between 1984 and 1994, an estimated 100,000 cases died in the Western Upper Nile province (Griekspoor *et al.* 1999). VL often has profound effects even when endemic. The reported incidences of VL in endemic communities vary between two cases/1000 person-years in Kenya (Schaefer *et al.* 1995) to 14 cases/1000 person-years in Ethiopia (Ali & Ashford 1994) and 40 cases/1000 person years in an endemic community in eastern Sudan (Zijlstra *et al.*, 1994). Similar incidences have been observed in endemic areas of Asia (Rijal *et al.* 2010; Singh *et al.* 2010a). VL can suddenly ‘unfold’ without being noticed for years, as was recently documented in the Bakool region of Somalia, where VL had never previously been reported (Marlet *et al.* 2003).

Despite the considerable health burden posed by VL, there have been few attempts to quantify the economic impacts of the disease in affected communities. Every member of a household where a VL case occurs may lose many days of productive life. This loss, when added to the costs of treatment, may push an already poor household further into poverty (Adhikari *et al.* 2009), reduce food security and keep children from school. As with HAT, the high expected costs associated with a case of VL may deter the search for treatment. In 1997, the socio-economic characteristics of a cohort of 938 parasitologically-confirmed cases of VL from Bihar were investigated (Thakur 2000). Most (75%) of these cases were classified as poor (with daily incomes below U.S.\$1.00), most (82%) were engaged in agriculture and/or animal husbandry, and most (77%) lived in mud or grass-covered houses. Delays in the seeking of medical care were considerable, with only 33% of the cases attending a healthcare facility within a month of the onset of their symptoms.

Over the last decade there have been at least seven attempts to estimate the financial and economic costs incurred by VL cases in India (Meheus *et al.* 2006; Sarnoff *et al.* 2010; Sundar *et al.* 2010a), Nepal (Sharma *et al.* 2004; Rijal *et al.* 2006; Adhikari *et al.* 2009) or Bangladesh (Sharma *et al.* 2006). The results of these studies unequivocally show that VL puts a considerable economic burden on the household of a case, with the cost of just one episode of VL exceeding the annual per-capita income. The estimates of the total cost of care incurred by the household of a case, including the direct and indirect costs, varied from U.S.\$113.60 (Rijal *et al.* 2006) to U.S.\$232.10 (Adhikari *et al.* 2009). The largest cost driver was found to be the income lost because of illness (i.e. an indirect cost), which may represent up to 60% of the total

household cost (Meheus *et al.* 2006). The relative importance of income loss is explained by several factors. The mean time period between the onset of symptoms and the first presentation by the case at a formal health facility is often very long — Hasker *et al.* (2010) found it to be about 40 days — and the case is usually unable to maintain their normal productive activities until treatment is completed. In addition, the household members who care for the case are also diverted away from their usual means of income generation. The direct costs of drugs or medical supplies are often another important contributor to total costs. Although treatment is generally provided free of charge at public health facilities, such facilities (particularly those in India and Bangladesh) may not have the necessary drugs in stock, forcing households to buy drugs from private pharmacies, where the drugs may cost more than three times their official price (Sharma *et al.* 2006). Sometimes households may also need to purchase the medical supplies needed for the administration of the drugs, such as intravenous sets. As described by Adhikari *et al.* (2009), household expenditure on VL may be catastrophic. Coping strategies including the use of savings and the sale of assets and livestock, although the latter tends to reduce further the livelihood of the VL-affected households. More than 80% of such households investigated by Sarnoff *et al.* (2010) took a loan to pay for the costs of care, even though the interest payable on such a loan may exceed the amount loaned (Sharma *et al.* 2006; Adhikari *et al.* 2009).

There appears to be no published information on the economic impact of VL in East Africa or Latin America, although one relevant study is currently in progress in Sudan (A.A. Abuzaid, unpubl. obs.).

A few studies have addressed the choice of the most cost-effective strategy for VL control. In the first published cost-effectiveness analysis for VL, Griekspoor *et al.* (1999) tried to answer the question whether VL control was cost-effective in an emergency setting, in comparison to other relief measures (such as diarrhoea control). They estimated that, in a relief programme in Sudan, VL treatment cost U.S.\$18.40 for each disability-adjusted life-year (DALY) averted (with an uncertainty range of U.S.\$13.53–U.S.\$27.63) and U.S.\$595 for each life saved. The mean cost of the intervention was U.S.\$394/treated VL case. As any health intervention costing less than U.S.\$25/DALY averted is considered ‘very good value for money’ by the World Bank, Griekspoor *et al.* (1999) concluded that, at least in the local conditions, their treatment programme was cost-effective. Even more encouragingly, these estimates of cost-effectiveness were based on the use of branded sodium stibogluconate (i.e. Pentostam™; GlaxoSmithKline, Brentford, U.K.), at a mean cost of U.S.\$100 per patient. The control programme has since switched to the use of generic antimonial drugs, which reduce drug costs by a factor of 5. Although generic antimonials have been shown to be effective in East Africa (Veeken *et al.* 2000) and to be cost-effective in

areas where the frequency of cure following antimonial treatment exceeds 94% (Vanlerberghe *et al.* 2007), they are, however, no longer recommended in India and Nepal because of high frequencies of failure (Sundar *et al.* 2000; Rijal *et al.* 2003).

Recently a number of therapeutic breakthroughs have occurred, including the clinical development of miltefosine as the first oral drug for VL (Sundar *et al.* 2002), the evaluation of paromomycin in combination therapy (Thakur *et al.* 2000; Melaku *et al.* 2007) and the development of a single-dose regimen of liposomal amphotericin B (Sundar *et al.* 2010b). Although a single dose of liposomal amphotericin B has been shown to be very effective, the cost of the drug still poses a barrier to its routine use as a first-line treatment (unless the international community is willing to finance the drug). While the VL-elimination initiative has adopted miltefosine as a first-line treatment, parasite resistance to both this drug and paromomycin may develop, as shown experimentally (Seifert *et al.* 2003). To safeguard the few effective drugs that currently exist from the emergence of resistance, combination therapies are increasingly being advocated, with the results from phase-III trials of combinations eagerly awaited. The additional advantages of combination therapies include shorter treatment durations and lower costs to both the public-health system and affected households. Olliaro *et al.* (2009) compared the cost-effectiveness of a combination of liposomal amphotericin B and miltefosine with those of various monotherapies and found that, at a cost of U.S.\$124–U.S.\$160/death averted, the combination therapy was a ‘competitive’ alternative. Another cost-effectiveness study for the Indian subcontinent, conducted from a societal perspective (including both health system and patient costs), was recently completed by Meheus *et al.* (2010). This study compared three plausible combination therapies with all possible monotherapies and showed that 10 days of treatment with a miltefosine–paromomycin combination was the most cost-effective treatment strategy (costing U.S.\$92/death averted), followed by a combination of liposomal amphotericin B with either paromomycin or miltefosine.

CONCLUSIONS

The fact that sleeping sickness and human visceral leishmaniasis have such low profiles, at both the national and international level, has prevented health systems in affected countries from allocating the appropriate resources for the control of these diseases. The Millennium Development Goals, with their emphasis on poverty reduction, provide a unique ‘window of opportunity’ to raise awareness about these diseases, and other NTD, at international level. A supranational approach to surveillance and control — as in the VL-elimination initiative in South Asia — might well be justified, as foci are often spread across borders. The results of

studies on health economics can support disease-control initiatives and help strengthen the case for greater investment in research on, and control of, VL, HAT and other poverty-related diseases.

REFERENCES

- Adhikari SR, Maskay NM, Sharma BP (2009) Paying for hospital-based care of kala-azar in Nepal: assessing catastrophic impoverishment and economic consequences. *Health Policy Plann* 24: 129–39.
- Ali A, Ashford RW (1994) Visceral leishmaniasis in Ethiopia IV Prevalence incidence and relation of infection to disease in an endemic area. *Ann Trop Med Parasitol* 88: 289–93.
- Alvar J, Yactayo S, Bern C (2006) Leishmaniasis and poverty. *Trends Parasitol* 22: 552–57.
- Boelaert M, Meheus F, Sanchez A, et al. (2009) The poorest of the poor: a poverty appraisal of households affected by visceral leishmaniasis in Bihar India. *Trop Med Int Health* 14: 639–44.
- Chappuis F, Lima MA, Flevaud L, Ritmeijer K (2010) Human African trypanosomiasis in areas without surveillance. *Emerg Infect Dis* 16: 354–56.
- Das P, Samuels S, Desjeux P, et al. (2010) Annual incidence of visceral leishmaniasis in an endemic area of Bihar India. *Trop Med Int Health* 15 (S2): 4–11.
- Desjeux P (1996) Leishmaniasis Public health aspects and control. *Clin Dermatol* 14: 417–23.
- Dujardin JC, Campino L, Canavate MC, et al. (2008) Spread of vectorborne diseases and neglect of leishmaniasis Europe. *Emerg Infect Dis* 14: 1013–18.
- Gouteux JP, Sinda D (1990) Community participation in the control of tsetse flies Large scale trials using the pyramid trap in the Congo. *Trop Med Parasitol* 41: 49–55.
- Gouteux JP, Bansimba P, Noireau F, Frezil JL (1987) Le coût du traitement individuel de la trypanosomiase à T.b. gambiense dans le foyer du Niari (Congo). *Med Trop* 47: 61–63.
- Griekspoor A, Sondorp E, Vos T (1999) Cost-effectiveness analysis of humanitarian relief interventions: visceral leishmaniasis treatment in the Sudan. *Health Policy Plann* 14: 70–76.
- Hasker E, Singh SP, Malaviya P, et al. (2010) Management of visceral leishmaniasis in rural primary health care services in Bihar India. *Trop Med Int Health* 15 (S2): 55–62.
- Holveck JC, Ehrenberg JP, Ault SK, et al. (2007) Prevention control and elimination of neglected diseases in the Americas: pathways to integrated inter-programmatic inter-sectoral action for health and development. *BMC Public Health* 7: 6.
- Lancien J (1991) Lutte contre la maladie du sommeil dans le sud-est Ouganda par piégeage des glossines. *Ann Soc Belg Med Tr* 71(S1): 35–47.
- Laveissière C, Couret D (1986) La campagne pilote de lutte contre la trypanosomiase humaine dans le foyer de Vavoua (Côte d'Ivoire) 5 Bilan financier Cahiers ORSTOM *Série Entomologie Médicale et Parasitologie* 24: 149–53.
- Lutumba P, Robays J, Miaka mia Bilenge C, et al. (2005) Trypanosomiasis control Democratic Republic of Congo 1993–2003. *Emerg Infect Dis* 11: 1382–88.
- Lutumba P, Makieya E, Shaw A, Meheus F, Boelaert M (2007) Human African trypanosomiasis in a rural community Democratic Republic of Congo. *Emerg Infect Dis* 13: 248–54.
- Magnus E, Vervoort T, van Meirvenne N (1978) A card-agglutination test with stained trypanosomes (CATT) for the serological diagnosis of *Trypanosoma brucei gambiense* trypanosomiasis. *Ann Soc Belg Med Tr* 58: 169–76.
- Marlet MV, Wuillaume F, Jacquet D, Quispe KW, Dujardin JC, Boelaert M (2003) A neglected disease of humans: a new focus of visceral leishmaniasis in Bakool Somalia. *Trans R Soc Trop Med Hyg* 97: 667–71.
- Meheus F, Boelaert M, Baltussen R, Sundar S (2006) Costs of patient management of visceral leishmaniasis in Muzaffarpur Bihar, India. *Trop Med Int Health* 11: 1715–24.
- Meheus F, Balasegaram M, Oliario P, et al. (2010) Cost-effectiveness analysis of combination therapies for visceral leishmaniasis in the Indian subcontinent. *PLoS Negl Trop Dis* 4: e818.
- Melaku Y, Collin SM, Keus K, Gatluak F, Ritmeijer K, Davidson RN (2007) Treatment of kala-azar in southern Sudan using a 17-day regimen of sodium stibogluconate combined with paromomycin: a retrospective comparison with 30- day sodium stibogluconate monotherapy. *Am J Trop Med Hyg* 77: 89–94.
- Moore A, Richer M, Enrile M, Losio E, Roberts J, Levy D (1999) Resurgence of sleeping sickness in Tambura county Sudan. *Am J Trop Med Hyg* 61: 315–18.
- Murray CJL, Lopez AD (Eds) (1996) *The Global Burden of Disease: a comprehensive assessment of mortality and disability from diseases injuries and risk factors in 1990 and projected to 2020*. Cambridge MA: Harvard School of Public Health.
- Nightingale SL (1991) Drug for sleeping sickness approved. *JAMA* 265: 1229.
- Oliario P, Darley S, Laxminarayan R, Sundar S (2009) Cost-effectiveness projections of single and combination therapies for visceral leishmaniasis in Bihar, India. *Trop Med Int Health* 14: 918–25.
- Priotto G, Kasparian S, Mutombo W, et al. (2009) Nifurtimox–eflornithine combination therapy for second-stage African *Trypanosoma brucei gambiense* trypanosomiasis: a multicentre randomised phase III non-inferiority trial. *Lancet* 374: 56–64.
- Rijal S, Chappuis F, Singh R, et al. (2003) Treatment of visceral leishmaniasis in south-eastern Nepal: decreasing efficacy of sodium stibogluconate and need for a policy to limit further decline. *Trans R Soc Trop Med Hyg* 97: 350–54.
- Rijal S, Koirala S, van der Stuyft P, Boelaert M (2006) The economic burden of visceral leishmaniasis for households in Nepal. *Trans R Soc Trop Med Hyg* 100: 838–84.
- Rijal S, Uranw S, Chappuis F, et al. (2010) Epidemiology of *Leishmania donovani* infection in high-transmission foci in Nepal. *Trop Med Int Health* 15(S2): 21–28.
- Robays J, Bilengue MM, van der Stuyft P, Boelaert M (2004) The effectiveness of active population screening and treatment for sleeping sickness control in the Democratic Republic of Congo. *Trop Med Int Health* 9: 542–50.
- Sarnoff R, Desai J, Desjeux P, et al. (2010) The economic impact of visceral leishmaniasis on rural households in one endemic district of Bihar India. *Trop Med Int Health* 15(S2): 42–49.
- Schaefer KU, Kurtzhals JA, Gachihi GS, Muller AS, Kager PA (1995) A prospective sero-epidemiological study of visceral leishmaniasis in Baringo district Rift Valley province Kenya. *Trans R Soc Trop Med Hyg* 89: 471–75.

- Seifert K, Matu S, Perez-Victoria FJ, Castanys S, Gamarro F, Croft SL (2003) Characterisation of *Leishmania donovani* promastigotes resistant to hexadecylphosphocholine (miltefosine). *Int J Antimicrob Agents* 22: 380–87.
- Sharma AD, Bern C, Varghese B, et al. (2006) The economic impact of visceral leishmaniasis on households in Bangladesh. *Trop Med Int Health* 11: 757–64.
- Sharma BP, Maskay NM, Adhikari SR, et al. (2004) Socio-economic determinants of kala-azar in Nepal. *J Nepal Health Res Counc* 2: 35–42.
- Shaw AP (1989) Comparative analysis of the costs and benefits of alternative disease control strategies: vector control versus human case finding and treatment. *Ann Soc Belg Med Tr* 69(S1): 237–53.
- Shaw AP, Cattand P (2001) Analytical tools for planning cost-effective surveillance in gambiense sleeping sickness. *Med Trop* 61: 412–21.
- Singh P, Reddy DC, Rai M, Sundar S (2006) Serious underreporting of visceral leishmaniasis through passive case reporting in Bihar, India. *Trop Med Int Health* 11: 899–905.
- Singh SP, Picado A, Boelaert M, et al. (2010a) The epidemiology of *Leishmania donovani* infection in high transmission foci in India. *Trop Med Int Health* 15(S2): 12–20.
- Singh VP, Ranjan A, Topno RK, et al. (2010b) Estimation of under-reporting of visceral leishmaniasis cases in Bihar, India. *Am J Trop Med Hyg* 82: 9–11.
- Stanghellini A, Josenando T (2001) The situation of sleeping sickness in Angola: a calamity. *Trop Med Int Health* 6: 330–34.
- Sundar S, More DK, Singh MK, et al. (2000) Failure of pentavalent antimony in visceral leishmaniasis in India: report from the center of the Indian epidemic. *Clin Infect Dis* 31: 1104–07.
- Sundar S, Jha TK, Thakur CP, et al. (2002) Oral miltefosine for Indian visceral leishmaniasis. *N Engl J Med* 347: 1739–46.
- Sundar S, Arora R, Singh SP, Boelaert M, Varghese B (2010a) Household cost-of-illness of visceral leishmaniasis in Bihar India. *Trop Med Int Health* 15(S2): 50–54.
- Sundar S, Chakravarty J, Agarwal D, Rai M, Murray HW (2010b) Single-dose liposomal amphotericin B for visceral leishmaniasis in India. *N Engl J Med* 362: 504–12.
- Thakur CP (2000) Socioeconomic of visceral leishmaniasis in Bihar (India). *Trans R Soc Trop Med Hyg* 94: 156–57.
- Thakur CP, Kanyok TP, Pandey AK, et al. (2000) A prospective randomized comparative open-label trial of the safety and efficacy of paromomycin (aminosidine) plus sodium stibogluconate versus sodium stibogluconate alone for the treatment of visceral leishmaniasis. *Trans R Soc Trop Med Hyg* 94: 429–31.
- Trouiller P, Torreele E, Olliaro P, et al. (2001) Drugs for neglected diseases: a failure of the market and a public health failure? *Trop Med Int Health* 6: 945–51.
- Trouiller P, Olliaro P, Torreele E, Orbinski J, Laing R, Ford N (2002) Drug development for neglected diseases: a deficient market and a public health policy failure. *Lancet* 359: 2188–94.
- Vanlerberghe V, Diap G, Guerin PJ, et al. (2007) Drug policy for visceral leishmaniasis: a cost-effectiveness analysis. *Trop Med Int Health* 12: 274–83.
- Veeken H, Ritmeijer K, Seaman J, Davidson R (2000) A randomized comparison of branded sodium stibogluconate and generic sodium stibogluconate for the treatment of visceral leishmaniasis under field conditions in Sudan. *Trop Med Int Health* 5: 312–17.
- Welburn SC, Fèvre EM, Coleman PG, Maudlin I (2004) *Epidemiology of human African trypanosomiasis*. In: Maudlin I, Holmes PH, Miles MA (eds.) *The Trypanosomiasis*. Wallingford UK: CABI. pp. 219–232.
- Zijlstra EE, el-Hassan AM, Ismael A, Ghalib HW (1994) Endemic kala-azar in eastern Sudan: a longitudinal study on the incidence of clinical and subclinical infection and post-kala-azar dermal leishmaniasis. *Am J Trop Med Hyg* 51: 826–36.



CHAPTER 3

The poorest of the poor: a poverty appraisal of households affected by visceral leishmaniasis in Bihar, India

Boelaert M
Meheus F
Sanchez A
Singh SP
Vanlerberghe V
Picado A
Meessen B
Sundar S

ABSTRACT

Objective

To provide data about wealth distribution in visceral leishmaniasis (VL)-affected communities compared to that of the general population of Bihar State, India.

Methods

After extensive disease risk mapping, 16 clusters with high VL transmission were selected in Bihar. An exhaustive census of all households in the clusters was conducted and socio-economic household characteristics were documented by questionnaire. Data on the general Bihar population taken from the National Family Health Survey of India were used for comparison. An asset index was developed based on Principal Components Analysis and the distribution of this asset index for the VL communities was compared with that of the general population of Bihar.

Results

83% of households in communities with high VL attack rates belonged to the two lowest quintiles of the Bihar wealth distribution. All socio-economic indicators showed significantly lower wealth for those households.

Conclusion

Visceral leishmaniasis clearly affects the poorest of the poor in India. They are most vulnerable, as this vector-borne disease is linked to poor housing and unhealthy habitats. The disease leads the affected households to more destitution because of its impact on household income and wealth. Support for the present VL elimination initiative is important in the fight against poverty.

INTRODUCTION

Visceral leishmaniasis (VL) or kala-azar is a parasitic disease affecting an estimated 500,000 new cases per year, mostly in the Indian subcontinent and East Africa, though this figure is subject to substantial uncertainty. VL usually occurs in small clusters in remote areas with poor access to health services and exact case numbers are poorly documented. In prospective studies of such clusters in East-Africa, incidence rates of VL varied between 2 / 1000–40 / 1000 person-years (Ali & Ashford 1994; Zijlstra *et al.* 1994; Schaefer *et al.* 1995). Incidence rates reported from the Indian subcontinent in 2001–2003 were in similar ranges (Singh *et al.* 2006). These rates are comparable to or higher than the incidence rates of tuberculosis at community level, but cannot be extrapolated to the population at large beyond the boundaries of the clusters where active VL transmission occurs, whereas tuberculosis is more homogeneously distributed. Nonetheless, VL can have a disastrous impact when an epidemic strikes a non-immune population. In famine-affected Southern Sudan, a VL epidemic caused the death of an estimated 100 000 people between 1984 and 1994 in Western Upper Nile Province (Seaman *et al.* 1996). Visceral leishmaniasis is nowadays considered as one of the ‘most neglected diseases’, a term coined to highlight the lack of innovation in both its clinical management and control (Trouiller *et al.* 2001). Several authors also concur that VL is a poverty-related disease, though the importance of VL as a hindrance for economic development is still not fully recognized (Wijeyaratne *et al.* 1994).

Advocates for the control of the disease will often claim that VL disproportionately affects ‘the poorest of the poor’ (Alvar *et al.* 2006), but this statement was not substantiated so far, apart from showing that VL patients live below the poverty line (Desjeux 1996), but so do many of their non-affected fellow citizens. In the context of a community intervention trial on the effect of long-lasting impregnated bed nets in the prevention of VL conducted in Bihar, India (the KALANET trial, see <http://www.kalanetproject.org>), we studied the socio-economic profile of the households living in areas with high VL endemicity and compared it to that of the general population of Bihar.

METHODS

Study area

Within the framework of the KALANET community intervention trial (Clinicaltrials.gov CT-2005-015374), we started in February 2006 to identify eligible VL clusters (i.e. well-defined communities with 350–1500 inhabitants) with a high incidence of VL in Muzaffarpur district (3.7 million inhabitants) in the northern part of Bihar State, India. Preliminary information

about villages with large numbers of VL cases was obtained from the public health system (Primary Health Centres, District Hospital and Medical College) and private and charitable hospitals. By the end of March 2006, we had identified 35 clusters with the highest reported number of VL cases in Muzaffarpur district. This was followed by a house to house survey of these clusters in May 2006 to evaluate the incidence rate of VL during the preceding three years (3.5 years in some clusters; 2003–2005 / 06). From these, 16 study clusters were selected (Table 1), based on five characteristics: (i) Reporting VL cases in each year of the 3-year-period, thus showing continuous transmission of the disease; (ii) an average of at least 0.8% VL incidence rate over the past 3 or 3.5 years; (iii) highest incidence rates; (iv) population between 350 and 1500; (v) minimum distance between any two clusters of 1 km.

Table 1: Selected clusters with high visceral leishmaniasis (VL) incidence in Muzaffarpur district, Bihar

| Cluster | Population | VL incidence | | | Annual incidence rate per 100 person/years |
|---------|------------|--------------|------|------------|-----------------------------------------------|
| | | 2003 | 2004 | 2005/2006* | |
| 1* | 779 | 20 | 1 | 6 | 0.99 |
| 2 | 1210 | 7 | 12 | 27 | 1.27 |
| 3 | 716 | 6 | 9 | 5 | 0.93 |
| 4 | 601 | 6 | 6 | 9 | 1.16 |
| 5* | 605 | 6 | 7 | 11 | 1.13 |
| 6 | 517 | 2 | 8 | 11 | 1.35 |
| 7* | 648 | 6 | 3 | 37 | 2.02 |
| 8 | 630 | 7 | 8 | 12 | 1.43 |
| 9* | 850 | 1 | 4 | 31 | 1.21 |
| 10* | 559 | 6 | 10 | 24 | 2.04 |
| 11 | 464 | 11 | 2 | 4 | .122 |
| 12* | 728 | 5 | 10 | 21 | 1.41 |
| 13* | 459 | 5 | 6 | 11 | 1.37 |
| 14* | 1150 | 11 | 13 | 32 | 1.39 |
| 15 | 350 | 8 | 5 | 4 | 1.62 |
| 16* | 1051 | 4 | 15 | 65 | 2.28 |

* Figure represents incidence calculated over 3.5 years; these are clusters where the number of VL cases during the first 6 months in 2006 was also collected

Socio-economic data

In the 16 study clusters, extensive household (HH) and individual data were collected in August–September 2006 on pre-designed questionnaires that included standardized questions on socio-economic items in an exhaustive census of the households. Information was collected from 2013 households and double-entered into an Access database.

As comparison data, we used data from the third National Family Health Survey (NFHS-3) of India (authorized on 26/11/2007) to document the socioeconomic status of the households in the state of Bihar, to which the above clusters belonged. The Indian NFHS are nationwide surveys conducted with a representative sample of households throughout the country, which provide data on population and health indicators such as fertility, family planning, infant and child mortality, maternal and child health, nutrition or morbidity and health care. These surveys are implemented by the International Institute for Population Sciences of Mumbai, and belong to the Monitoring and Evaluation to Assess and Use Results (MEASURE), Demographic and Health Surveys (DHS) project, composed of several partners who provide technical assistance and funding to more than 200 surveys in 75 countries since 1984. The Indian NFHS-3 was conducted between November 2005 and August 2006, with an overall sample size of 109 041 households (International Institute for Population Sciences & Macro International 2007) and 2997 households from the state of Bihar. We used the latter data.

Asset index

To allow poverty ranking of the Bihar as well as the households living in areas with a high VL endemicity, we chose an approach based on an asset index, which has been widely used as a proxy to measure poverty when income and expenditure data are of poor quality or not available. Data on ownership of durable assets in the households, characteristics of the habitat and access to basic services are the indicators most commonly collected and used to construct a single ‘asset index’. This measure reflects the long-run household wealth or living standard, so unequal measures of the asset index can be considered as proxies for inequalities in long-run wealth (Filmer & Pritchett 2001; Falkingham & Namazie 2002; McKenzie 2005). We selected nine indicators, reflecting four dimensions of longrun wealth, to be included in the asset index (Table 2).

Table 2: Indicators included in the asset index

| Category | Asset indicator |
|--------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------|
| Housing structure | Type of house: Kachha; Semi-pucca; Pucca |
| Household access to utilities and infrastructure | Main type of fuel used for cooking : traditional (animal dung / agricultural crop waste / straw, shrubs, grass) or modern (gas, electricity or petrol) |
| Productive assets | Whether the household has electricity or not |
| | Whether the household has a bicycle or not |
| | Whether the household has a motorcycle / scooter or not |
| Non-productive assets | Whether the household has a car / tractor or not |
| | Whether the household has a radio or not |
| | Whether the household has a television or not |
| | Whether the household has a cot / bed or not |

We used Principal Components Analysis (PCA) to aggregate the information from the different indicators in order to create an asset index that explained 35.4% of variance. The analysis was performed with spss v.15.0 software. PCA is a multivariate statistical technique that assigns weights to the indicators so to create the weighted linear combination of the variables that accounts for the largest amount of the total variation in the data (Kleinbaum *et al.* 1988; Vyas & Kumaranayake 2006). PCA has been applied by several authors to create indices of relative poverty (Filmer & Pritchett 2001; McKenzie 2005; Vyas & Kumaranayake 2006; Zeller *et al.* 2006; Morris *et al.* 2007) with the underlying assumption that relative poverty explains the maximum variance in the variables.

When the indicators included in the PCA differ in their measurement scale, they should be converted into standardized variables, so that the resulting index can be represented by the formula:

$$A_i = f_1 \frac{(a_{i1} - \bar{a})}{s_1} + \dots + f_n \frac{(a_{in} - \bar{a}_n)}{s_n}$$

where A_i is the asset index of the i th household, f_1 is the weight for the first indicator, a_{i1} is the i th household’s value for the first indicator and \bar{a}_1 and s_1 are the mean and standard deviation of the first indicator over all households (Filmer & Pritchett 2001).

We used the asset index score to classify the households from Bihar state in wealth quintiles. Using the formula mentioned above, we applied the weights obtained with PCA to the indicator values of the households living in areas with a high VL endemicity (KALANET project clusters),

in order to place them within the wealth quintiles of the state of Bihar. Student’s t-test was used to compare asset ownership between Bihar and highly endemic VL clusters.

Ethical approval

The study was conducted as part of the KALANET community intervention trial (Clinicaltrials.gov CT-2005-015374), for which ethical clearance was obtained from the Institutional Review Boards at Banaras Hindu University and the Institute of Tropical Medicine.

RESULTS

The distribution of the asset index score for the general population of the State of Bihar is shown in Figure 1. The asset index is a measure of relative poverty, so the lower the score, the poorer the household relative to all others with higher scores.




Figure 1: Histogram of the Bihar Asset Index

Figure 2 shows how households from highly endemic VL areas fit within the wealth distribution of the households in Bihar based on NFHS-3 data. The number of households in the top two quintiles (8.3%) is very low, and the vast majority of households in VL endemic areas are situated in the two poorest quintiles of the wealth distribution of Bihar (83.3%).




Figure 2: Distribution of wealth of households living in highly endemic VL clusters within the socioeconomic quintiles of all Bihar households

Table 3 shows the distribution of the socio-economic indicators included in the asset index for the Bihar and the populations living in areas with a high VL endemicity. There are significant differences between both populations in all the indicators, pointing out the extremely low numbers of households with appropriate housing and access to basic services and assets in the communities affected by kala-azar.

Table 3: Socio-economic indicators for Bihar and highly endemic visceral leishmaniasis (VL) cluster populations

| Asset indicator | Bihar | | Highly endemic VL cluster | | P-value |
|---------------------------|-------|---------|---------------------------|--------|----------|
| | n | (%) | n | (%) | |
| Type of house | | | | | |
| Kachha | 794 | (26.5) | 1418 | (70.4) | < 0.0001 |
| Semi-Pucca | 1248 | (41.6) | 477 | (23.7) | < 0.0001 |
| Pucca | 955 | (31.9) | 118 | (5.9) | < 0.0001 |
| Traditional* cooking fuel | 1639 | (54.7) | 1845 | (91.7) | < 0.0001 |
| Electricity | 1212 | (40.4) | 193 | (9.6) | < 0.0001 |
| Bicycle | 1646 | (54.9) | 839 | (41.7) | < 0.0001 |
| Motorcycle/scooter | 338 | (11.3) | 46 | (2.3) | < 0.0001 |
| Car/tractor | 96 | (3.2) | 10 | (0.5) | < 0.0001 |
| Radio | 983 | (32.8) | 197 | (9.8) | < 0.0001 |
| Television | 864 | (28.8) | 136 | (6.8) | < 0.0001 |
| Cot/bed | 2722 | (90.8) | 1556 | (77.3) | < 0.0001 |
| Total | 2997 | (100.0) | 2013 | 100.0) | |

* Traditional = Animal dung, agricultural crop waste, straw, shrubs or grass

Other socio-economic indicators (not included in the asset index) such as education, caste or religion of the head of the household also show significant differences (Table 4). Higher percentages of illiteracy are found amongst the heads of the households living in areas with a high VL endemicity. There was a stronger presence of Scheduled Castes (SC), Scheduled Tribes (ST) and Other Backward Castes (OBC), the lower castes in India, in the clusters affected by high VL transmission.

Table 4: Socio-economic characteristics of heads of households

| Variable | Bihar | | Highly endemic VL cluster | | P-value |
|------------|-------|---------|---------------------------|---------|----------|
| | n | (%) | n | (%) | |
| Caste* | | | | | |
| SC/ST | 503 | (16.8) | 440 | (21.9) | < 0.0001 |
| OBC | 1792 | (59.8) | 1381 | (68.6) | < 0.0001 |
| Other | 697 | (23.3) | 175 | (8.7) | < 0.0001 |
| Illiterate | 1588 | (53.0) | 1354 | (67.3) | < 0.0001 |
| Religion | | | | | |
| Hindu | 2477 | (82.6) | 1801 | (89.5) | < 0.0001 |
| Muslim | 513 | (17.1) | 212 | (10.5) | < 0.0001 |
| Total | 2997 | (100.0) | 2013 | (100.0) | |

* SC/ST: scheduled caste/scheduled tribe; OBC: other backward caste

DISCUSSION

This study is the first to show clearly that VL affects the poorest of the poor. Bihar state is known to be one of the poorer states of India, with 40% of its population living below the poverty line, while the national average is 29% (World Bank 2008). Our data indicate that the communities with high and active VL transmission over the past three years are situated at the lower end of the wealth distribution of Bihar state.

The fact that data from two different surveys were used to construct the asset index can be considered a limitation, though both were carried out contemporaneously and worked with an almost similar set of standard questions. PCA was used to derive the weights assigned to the assets included in the index. Filmer and Pritchett (2001) showed that a PCA-based index performed as well as consumption expenditure, and more recently a study comparing different methods to assign weights to the indicators concluded that PCA was a suitable method (Howe *et al.* 2008) although the weighing method might be of less importance than the data used.

There exists ample evidence that poverty and ill-health are intertwined, with poor people suffering worse health and having more limited access to health care. Likewise, ill-health may lead to a loss of income through absence from work and high health care costs (Wagstaff 2002), driving poor households even further into poverty (Bloom & Canning 2003). As described by Alvar *et al.* (2006), the relationship of VL to poverty is complex and additional research that leads to a better understanding on the interaction between VL and poverty is needed. In the Indian subcontinent, infection risk is clearly related to poor housing conditions and sanitation, as the sand fly vector *Phlebotomus argentipes* breeds in cracks of mud-plastered houses and moist soils. The disease has a chronic course and is fatal if left untreated; therefore households often sell their assets and take loans to pay for health care and expensive drugs which may result in significant impoverishment (Desjeux 1996). Moreover in every family with a VL case, the severely debilitating disease has a significant impact on earnings and consumption with many days of productive life lost due to illness. Two studies from Nepal and a study from India showed how the total cost of a VL episode to affected families exceeded their average annual per capita income and how households either sold part of their livestock or took loans to cover the costs (Adhikari & Maskay 2003; Rijal *et al.* 2006; Meheus *et al.* 2006).

Although our results have shown that in Bihar communities with high VL incidence are considerably poorer than the rest of the state, this does not necessarily imply a causal relationship between VL and poverty. Wealth differences between VL affected clusters and the rest of Bihar could also be explained by other factors such as differences in ecological conditions and economic growth within Bihar or individual differences such as caste, literacy rates and land ownership. Comparing VL endemic clusters with for example North Bihar only, which would have taken into account regional wealth differences to a certain extent, was not possible since the NFHS-3 does not include district level identifiers for reasons of confidentiality (the survey also measures HIV-status). Even so, this does not alter the findings presented in this paper.

Though global case numbers of VL might rank lower than those of HIV/AIDS, malaria and tuberculosis, the social and economic implications of VL in affected communities are profound. VL clearly affects the poorest of the poor in India and adds to their destitution. This study suggests that current and future preventive measures for such a deadly disease need to be subsidized or provided free to the majority of households living in VL affected areas of India, or else they might prove futile. Support for the present VL elimination initiative is important in the fight against poverty.

ACKNOWLEDGEMENTS

This study received financial support from the KALANET project funded by the European Union DG Research Sixth Framework Programme / INCODEV. (Proposal Contract No.: 015374). The funding source had no role in the study design; collection, analysis, and interpretation of data; in the writing of the report; and in the decision to submit the article for publication. The author's work was independent of the funders (the funding source had no involvement).

REFERENCES

Adhikari SR, Maskay NM (2003) The economic burden of Kala-azar in households of the Danusha and Mahottari districts of Nepal. *Acta Trop* 88: 1–2.

Ali A, Ashford RW (1994) Visceral leishmaniasis in Ethiopia. IV. Prevalence, incidence and relation of infection to disease in an endemic area. *Ann Trop Med Parasitol* 88: 289–93.

Alvar J, Yactayo S, Bern C (2006) Leishmaniasis and poverty. *Trends Parasitol* 22, 552–57.

Bloom D, Canning D (2003) The health and poverty of nations: from theory to practice. *J Hum Dev* 4: 47–71.

Desjeux P (1996) Leishmaniasis. Public health aspects and control. *Clin Dermatol* 14: 417–423.

Falkingham J, Namazie C (2002) Measuring Health and Poverty: A Review of Approaches to Identifying the Poor. London: DFID Health Systems Resource Centre.

Filmer D, Pritchett LH (2001) Estimating wealth effects without expenditure data—or tears: an application to educational enrolments in states of India. *Demography* 38: 115–32.

Howe LD, Hargreaves JR, Huttly SRA (2008) Issues in the construction of wealth indices for the measurement of socioeconomic position in low-income countries. *Emerg Themes Epidemiol* 5: 3.

International Institute for Population Sciences and Macro International (2007) National Family Health Survey (NFHS-3), 2005–06. India, Volume I. Mumbai: International Institute for Population Sciences.

Kleinbaum DG, Kupper LL, Muller KE (1988) *Applied Regression Analysis and Other Multivariable Methods*, 2nd edn. Boston: PWS-Kent Publishing Company.

McKenzie DJ (2005) Measuring inequality with asset indicators. *J Popul Econ* 18: 229–60.

Meheus F, Boelaert M, Baltussen R, Sundar S (2006) Costs of patient management of visceral leishmaniasis in Muzaffarpur, Bihar, India. *Trop Med Int Health* 11: 1715–24.

Morris SS, Ranson MK, Sinha T, Mills AJ (2007) Measuring improved targeting of health interventions to the poor in the context of a community-randomised trial in rural India. *Contemp Clin Trials* 283: 382–90.

Rijal S, Koirala S, Van der Stuyft P, Boelaert M (2006) The economic burden of visceral leishmaniasis for households in Nepal. *Trans R Soc Trop Med Hyg* 100: 838–41.

Schaefer KU, Kurtzhals JA, Gachihi GS, Muller AS, Kager PA (1995) A prospective sero epidemiological study of visceral leishmaniasis in Baringo District, Rift Valley Province, Kenya. *Trans R Soc Trop Med Hyg* 89: 471–75.

Seaman J, Mercer AJ, Sondorp E (1996) The epidemic of visceral leishmaniasis in western Upper Nile, southern Sudan: course and impact from 1984 to 1994. *Int J Epidemiol* 25: 862–71.

Singh SP, Reddy DCS, Rai M, Sundar S (2006) Serious underreporting of visceral leishmaniasis through passive case reporting in Bihar, India. *Trop Med Int Health* 11: 899–905.

Trouiller P, Olliaro P, Torreele E, Orbinski J, Laing R, Ford N (2001) Drugs for neglected diseases: a failure of the market and a public health failure? *Trop Med Int Health* 6: 945–51.

Vyas S, Kumaranayake L (2006) Constructing socio-economic status indices: how to use principal components analysis. *Health Policy Plann* 21: 459–68.

Wagstaff A (2002) Poverty and health sector inequalities. *Bull World Health Organ* 80: 97–105.

Wijeyaratne PM, Arsenault LK, Murphy CJ (1994) Endemic disease and development: the leishmaniasis. *Acta Trop* 56: 349–64.

Zeller M, Sharma M, Henry C, Lapenu C (2006) An operational method for assessing the poverty outreach performance of development policies and projects: results of case studies in Africa, Asia, and Latin America. *World Dev* 34: 446–64.

Zijlstra EE, el-Hassan AM, Ismael A, Ghalib HW (1994) Endemic kala-azar in eastern Sudan: a longitudinal study on the incidence of clinical and subclinical infection and post-kala-azar dermal leishmaniasis. *Am J Trop Med Hyg* 51: 826–36.



PART II

The economic impact and costs of patient management of visceral leishmaniasis



CHAPTER 4

The costs of patient management of visceral leishmaniasis in Muzaffarpur, Bihar, India

Meheus F
Boelaert M
Baltussen R
Sundar S

ABSTRACT

Objective

To identify and quantify the direct and indirect economic cost of treatment for visceral leishmaniasis (VL) with conventional Amphotericin B deoxycholate, currently the first-line treatment in Muzaffarpur.

Methods

Costs of patient management for VL were estimated from a societal and household perspective by means of a questionnaire designed for this study, interviews and financial reports.

Results

The total cost of care per episode of VL from the societal perspective was estimated at US\$355, equivalent to 58% of annual household income. The largest cost category was medical costs (55%), followed by indirect costs (36%) and non-medical costs (9%). The cost from the household perspective was equivalent to US\$217. The largest cost category was indirect costs (59%), followed by medical costs (27%) and non-medical costs (15%). Loss of income because of illness and hospitalization and expenses for drugs were the largest cost components.

Conclusions

The economic costs related to VL are substantial, both to society and the patient. Public health authorities in Bihar should focus on policies that detect VL in the early stage and implement interventions that minimize the burden to households affected by VL.

INTRODUCTION

Visceral leishmaniasis (VL) or Kala-azar is a life-threatening disease if left untreated. Ninety per cent of all cases occur in only five countries: Bangladesh, Brazil, Nepal, Sudan and India (WHO 2000); with India accounting for half of the cases and about 90% of Indian patients living in the state of Bihar (Sundar *et al.* 2001).

Visceral leishmaniasis in India is caused by *Leishmania donovani* and is transmitted by the bite of an infected phlebotomus female sandfly. Patients infected with VL usually present prolonged fever, diarrhoea, cough, abdominal pain, enlarged liver and/or spleen, nose bleeds and severe loss of weight. Complications of VL include post-kala-azar dermal leishmaniasis (Guerin *et al.* 2002).

Pentavalent antimonial compounds (Sbv) is the recommended treatment in all parts of the world except for Bihar where 64% of previously untreated patients showed unresponsiveness to Sbv, even with the WHO recommended treatment of 20 mg/kg/day for 30 days (Murray 2000; Sundar 2001). Conventional Amphotericin B deoxycholate is now the drug of choice in Bihar, with 15–20 infusions of 1 mg/kg given either on a daily or alternate basis enforcing prolonged hospitalization (Sundar *et al.* 2004). Amphotericin B shares drawbacks with Sbv such as length of therapy and parental administration; it is more expensive and requirements of infusions and close monitoring because of its potential toxicity necessitate hospitalization for the duration of treatment (Murray 2000). Over the past years, tangible progress has been made and has offered some promising alternatives to current treatments: Miltefosine has recently been registered in India and is the first oral drug for VL and Paromomycin, which is currently being tested in India, seems a promising low-cost alternative to current treatment with Amphotericin B in Bihar (Murray 2004).

Whilst the literature on clinical studies of treatment options for VL steadily increased in the 1990s (Olliaro *et al.* 2005), there has been little effort to quantify the economic consequences of the disease on the affected population. This paper presents the results from a costing analysis in Muzaffarpur, Bihar (India). The objectives of the study were to identify/quantify the economic cost of treatment for VL with conventional Amphotericin B deoxycholate, currently the first-line treatment in Muzaffarpur. Costs in this study are defined in terms of opportunity cost (or economic cost), which is the value of resources foregone that could have been used elsewhere. When opportunity costs are estimated, all inputs are valued, even those for which there was no monetary cost, such as donations or inputs that have been paid for below market price. Costs associated with patient management of VL include direct medical costs borne by the provider, the patient and its family, direct non-medical costs borne by patients and family and

indirect costs related to time lost from productive work borne by patient and family and were estimated from the perspective of society as a whole and that of the household. The analysis of household costs is important because we believe that as VL mainly affects individuals from the lowest socioeconomic class, treatment of the disease can play an important role in poverty reduction and should therefore try to minimize the cost to patients. Furthermore, over the past years there have been major advances in the development of anti-leishmanial drugs that have a different impact on costs (including costs to the patient). It is therefore important to analyse all costs associated with VL patient management from different perspectives. The findings from this study can be useful for further costing studies and for evaluating the recently developed alternatives with the current treatment for VL in Bihar. The paper is organized as follows: first we present the conceptual framework used for the analysis and the process of patient management, data collection and cost estimation. The conceptual framework draws on guidelines for cost and cost-effectiveness analysis in general (Creese & Parker 1994; Tan-Torres Edejer *et al.* 2003) and for particular diseases (Asenso-Okyere & Dzator 1997; WHO 2005). The subsequent section presents the results of our cost analysis from the societal as well as from the patient perspective, and discusses the implications of our findings for VL control.




Figure 1. Map showing endemicity of visceral leishmaniasis in Bihar.
Source: Sundar (2001)

METHODS

Study site

Data were collected in July and August 2005 at the Kala-azar Medical Research Centre (KAMRC). The KAMRC is a hospital that only provides care for VL and is run by a non-governmental charitable Sitaram Memorial Trust supported by grants and private donations. The hospital is located in Muzaffarpur 75 km away from the state capital of Patna. Muzaffarpur is the second largest city in Bihar (India) and is considered a highly endemic area of VL with a high degree of antimony resistance (Sundar 2001).

The region is the third most densely populated area of India and performs considerably worse with regard to poverty indicators compared to the rest of India. An estimated 40% of the population lives below the poverty line of 1 US\$ per day. The vast majority of the workforce (80%) is employed in agriculture with a predominance of subsistence agriculture (World Bank 2005). The total population in the district of Muzaffarpur is around 3.75 million inhabitants of whom the vast majority (90%) lives in a rural setting (Figure 1).

Process of patient management

In order to understand the process of data collection and correctly interpret the estimated costs a description of a standard patient management follows. First a patient suspected of having VL presents spontaneously or is being referred to the outpatient section of the KAMRC. Confirmation of Kala-azar is based on typical clinical and laboratory features like fever, splenomegaly, pancytopenia and diagnostic tests such as presence of parasites (asmatigotes) in spleen aspirate smears and/or positive rK39 rapid strip test. If the patient has Kala-azar, he/she is admitted to the hospital for the full duration of treatment. Up to this point, all costs (travel, food, diagnostic tests) are paid out-of-pocket by the patient. Once admitted, the patient receives conventional Amphotericin B deoxycholate on a daily or alternate (every 2 days) basis depending on the physiological characteristics of the patient. The patient is also regularly submitted to laboratory investigations to test for toxicity and side-effects related to the drug. On admission the patient only pays a nominal admission fee to the hospital and other services (e.g. tests) are provided free of charge. The drugs for VL (Amphotericin B) have to be retrieved by the patient or a family member outside the hospital at a local private pharmacy at a pre-negotiated subsidized cost and brought back to the hospital where it is administered intravenously by medical personnel. The i.v. sets and related equipment for the i.v. injections as well as the laboratory tests and the daily bed fee are provided free of charge by the hospital to the patient. Similarly medical supervision, nursing are also provided free of charge. During hospitalization, in most cases one or more family members stay with the patient on a permanent

basis. Once discharged, the patients comes back to the hospital after the 1st, 3rd, 6th and 12th month to check for side-effects and relapse. In this study, we only analysed the costs related to diagnosis and treatment of VL, and not the costs incurred by the patient after discharge.

Cost categories

The costs associated with patient management of VL can be direct (medical and non-medical) and indirect (time losses).¹




Figure 2. Costs of patient management of Visceral Leishmaniasis
Source: Adapted from Brouwer et al (2001)

Direct costs

Direct medical costs associated with VL include costs borne by the provider to deliver the intervention (i.e. provider costs) and costs borne by the patient and family to receive treatment (i.e. household costs). The provider costs are the resources used to implement and make the treatment for VL available. These include personnel costs (doctors, nurses), overhead costs (administration, maintenance and electricity), infrastructure (buildings), medical equipment (i.v. sets, syringes and needles), medication, laboratory tests and hotel costs (e.g. bed linen).

Costs borne by the household are the resources used to access the treatment for VL and can be medical or nonmedical. Direct medical costs include out-of-pocket expenditures on consultation fees, diagnostic tests and payment for VL drugs. The direct non-medical costs borne by the patient/family are travel costs to and from the health facility and drug stores and food expenses (Sauerborn *et al.* 1991; Barnum & Kutzin 1993; Asenso-Okyere & Dzator 1997; Duraisamy *et al.* 2006).

¹ A third category that is sometimes mentioned in the literature are the intangible costs. These are costs related to anxiety and pain to the patient and relatives and can be captured with non-monetary measures such as QALYs. These were not included in this study because of the difficulty to estimate these.

Indirect costs

Indirect costs refer to the value of time losses because the patient is unable to carry out his/her normal productive activities because of ill-health associated with VL. Time losses include travel time to and from the hospital and drugs stores, time lost at the hospital (e.g. consultation) and the time of hospitalization. The valuation of these time losses is a controversial issue. It is generally recommended not to include them in the analysis unless their exclusion may introduce bias into the estimates (Hutton & Baltussen 2005). As patients and their relatives require prolonged and repeated contact with the system, we valued the time of lost work with the human capital method where earnings (i.e. income) are used as a proxy of the opportunity cost of lost time². These time losses can be incurred by the patient and by the family members caring for the patient at the hospital (Figure 2).

Data collection and cost estimation

Information on provider’s cost was derived from annual financial reports for the year 2003–2004 of the medical centre and from the hospital administrator. These financial reports only present aggregated financial expenditure lines. This was not considered a problem as the KAMRC is fully dedicated to VL care, and all expenditures by the hospital can be attributed to the VL programme. Methods to allocate the overhead costs to different department or programmes were therefore not necessary. Items purchased by the hospital below market rates were corrected using market prices to reflect economic costs. For example the cost of the hospital building was valued with the rental value of a similar space in the same location (Hutton & Baltussen 2005).

Data on the patient’s direct and indirect costs were obtained from medical records (e.g. to retrieve the number of lab tests undergone), medical personnel and from patients and their relatives with a structured questionnaire designed for this study. The patient, the head of the household or a principal respondent was interviewed. The questionnaire collected data on demographic characteristics (age, relation to head of household, household size, education and literacy) and socioeconomic characteristics (occupation, monthly income of patient and of family members staying with the patient at the hospital as well as household income), on direct medical costs (consultation fees, laboratory tests and drugs) and direct non-medical costs (travel expenses, food costs) incurred by the patient and accompanying relatives. The questionnaire also included questions on health-seeking behaviour and costs prior to admission at the KAMRC and on additional expenditures to meet the costs of treatment: loans taken and

² The main approaches to value the time of lost work are the human capital method, the friction cost method and willingness to pay methods such as revealed preference studies and stated preference studies. For a discussion of these and other methods see Koopmanschap *et al.* (1995), Drummond *et al.* (1997) and Sculpher (2001).

interest charged, origin of the loan (moneylender, relative, village person, etc.) and assets sold (livestock, land, durables).

Direct medical costs to the patient were calculated by taking an ingredient approach (Creese & Parker 1994). This involves asking the patient or the principal respondent the quantities consumed and price paid for each cost item (mainly drugs and laboratory tests). Patients or relatives purchase Amphotericin B at a subsidized price at the local private pharmacy. Because this situation is very specific to the KAMRC and evaluating costs from a societal perspective, this subsidized price was replaced by the market price (Hutton & Baltussen 2005). An ingredient approach was also taken when calculating travel costs. Travel costs consist of visits to and from the health facility and drug stores by the patient and/or relatives as well as visits received by the patient from family members during his/her hospitalization. The travel costs were assessed by asking the patient or principal respondent the cost for a one-way journey to the health centre. This amount was then doubled to obtain the total travel cost for one trip (Cho-Min-Naing & Gatton 2004). The travel costs are for the patient and accompanying relatives together as it was difficult for the respondent to separate them (e.g. rented car for unlimited number of passengers, young patients not paying for rickshaw). Food costs were assessed by asking the daily food cost for the patient and all family members staying permanently with the patient at the hospital and for family members from the same household visiting the patient. This amount does not include food brought from the household by visiting members as it was difficult to estimate this cost.

Indirect costs were calculated by asking for the monthly income of the patient, family members staying with the patient at the hospital and the rest of the household. In many cases, respondents could only give a daily income. This daily income was then multiplied by 30 (days) to calculate the monthly income. It was also assumed that this daily income was stable over the entire month. Because many households were employed in agriculture and thus earned a very small monetary income, the questionnaire included questions on the amount of agricultural produce and the market price of these crops to allow for estimation of a monthly income if these were to be sold. Same was carried out for output of livestock (e.g. milk from cattle). The sum of the patient's earnings and those of its family members constituted the average monthly household income. In the interpretation of costs, care was taken not to double count certain cost items such as hospital costs.

Study population and sample

Patients were selected from admission records. All patients admitted/or presenting for follow-up at the time of the study (between 23 July 2005 and 4 August 2005) were approached. Patients on follow-up visit were only included if they had been discharged no longer than 6 months to minimize recall bias. All patients received treatment with conventional Amphotericin B deoxycholate. Seventy-seven patients were approached. After explaining the objective of the study and the type of data required, all patients gave their consent to participate with the study. Of the seventy-seven patients, fifty patients belonged to the group of inpatients and 27 were follow-up patients. All costs were identified in terms of the local currency (1US\$ is equivalent to INR 43.40 for the month July 2005 at the official exchange rate during the period of the study).

RESULTS

Demographic and socioeconomic profile of study participants

The surveyed patients were on average 16 years old (range 3–60 years). A majority of the patients were male (64%) and 18% of the patients were head of their household. The average size of the household was eight persons. Although, a majority of the patients (56%) reported having followed formal schooling with more educated men (33%) than women (23%), the average years of schooling was only 3 years. Twenty-six per cent of interviewed patients were engaged in agriculture and/or animal husbandry, 13% were working in salaried employment, 3% in business, 25% of the patients were students, 17% were engaged in domestic duties (cooking, cleaning, fetching water, wood, etc.). The remaining patients (16%) were not engaged in any productive activity because of their young age (median age 5 years.).

Of the patients earning a monetary income (n=30), 43% earned less than a dollar per day, 43% had an income between \$1 and \$2 and 14% earned more than \$2 on a daily basis. The median household income on a monthly basis was 2200 rupees (range 500–9000).

Clinical and treatment characteristics of study participants

The median duration of hospitalization was 18 days which is lower than the standard length of hospitalization of 28 days reported in the literature because most patients (88%) in this study were treated with conventional Amphotericin B deoxycholate with infusions on a daily basis and were admitted to the hospital for the full duration of treatment. (Table 1).

Table 1: Clinical and treatment characteristics (n=77)

| Variables | Median | (IQR ^a) |
|--------------------------------------|--------|---------------------|
| Duration of hospitalization (# days) | 18 | (18-19) |
| Duration of illness (# days) | 70 | (24-90) |
| Number of vials per patient | 8 | (5-11) |
| Body weight (kg) | 36 | (20-45) |

^a Interquartile range

Costs from the societal perspective

Medical, non-medical and indirect costs associated with VL patient management are presented in Table 2. The total cost of care per episode of VL was estimated at 15 400 rupees and is equivalent to 58% of annual household income. The total *medical* cost over the period of hospitalization amounted to 8490 rupees. The ‘accommodation’ cost for hospitalization per patient was 2736 rupees and is an aggregate of personnel costs, overhead costs, infrastructure. The cost of investigations (diagnostic and laboratory tests) per patient was 2700 rupees. The median cost of drugs is 2334 rupees and consists of Amphotericin B (92%) and miscellaneous drugs (8%). The medical supplies amount to 720 rupees per patient and include i.v. sets and 5% dextrose water necessary for the intravenous injections of Amphotericin B. *Non-medical* costs amounted in total to 1410 rupees and consists of transportation costs (33%) and food cost (67%). The patient received on average four visits from members of the same household. In all cases a relative stayed permanently with the patient at the hospital: in 69% of the cases this was one relative, in 26% of the cases this were two relatives and in the remaining 5% three relatives stayed with the patient at the hospital.

The median loss of income to the patient and attendants was 5300 rupees. Other costs include the monthly interest payment on loans taken to meet the costs of VL treatment and amounted to 200 rupees.

Costs from the household perspective

Costs from the perspective of the household include out-of pocket expenditures and indirect costs incurred by the patient and their relatives (Table 2). All patients incurred substantial out-of-pocket expenditures to acquire treatment for VL. The main cost item for the patient is the drug cost which amounts to 2160 rupees. Consultation fees amounted to 170 rupees. The median cost of investigations purchased by the patients amounted to 180 rupees and consists of diagnostic tests only. The median transportation cost was 460 rupees. The median daily food cost for the patient and relative(s) staying permanently at the hospital was 50 rupees. The total amount of out-of-pocket payments by the patient (and relatives) over the entire length of hospitalization was 3920 rupees.

Table 2. Costs of patient management from societal and household perspective (rupees)

| | Costs from the societal perspective | | | Costs from the household perspective | | |
|---------------------------------|-------------------------------------|---------------------|-----|--------------------------------------|---------------------|-----|
| | Median | (IQR ^a) | % | Median | (IQR ^a) | % |
| Medical costs | | | | | | |
| Consultation fees | - | | - | 170 ^b | | 2 |
| Accommodation cost | 2,736 | (2,736-2,888) | 18 | - | | |
| Investigations | 2,700 | (2,650-2,735) | 18 | 180 | (130-215) | 2 |
| Medicines | | | | | | |
| Amphotericin B | 2,160 | (1,350-2,970) | 14 | 2,160 | (1,350-2,970) | 23 |
| Other drugs ^c | 174 | (109-239) | 1 | - | | |
| Medical supplies | 720 | (450-990) | 5 | - | | |
| Total medical costs | 8,490 | | 55 | 2,510 | | 27 |
| Non-medical costs | | | | | | |
| Transportation costs | 460 | (340-660) | 3 | 460 | (340-660) | 5 |
| Food costs | 950 | (810-1,240) | 6 | 950 | (810-1,240) | 10 |
| Total non-medical costs | 1,410 | | 9 | 1,410 | | 15 |
| Indirect costs | | | | | | |
| Loss of income to the patient | 4,400 | (1,691-6,555) | 29 | 4,400 | (1,691-6,555) | 46 |
| Loss of income to the attendant | 900 | (600-1,305) | 5 | 900 | (600-1,305) | 10 |
| Monthly interest on loans | 200 | (95-250) | 1 | 200 | (95-250) | 2 |
| Total indirect costs | 5,500 | | 36 | 5,500 | | 59 |
| Total (rupees) | 15,400 | | 100 | 9,420 | | 100 |
| Total (US\$) | 355 | | | 217 | | |

^a Interquartile range

^b Is the same for all

^c These are miscellaneous drugs such as aspirin, antibiotics,...

The total loss of income to the patient (income loss because of illness and hospitalization) was 4400 rupees, while the income loss for the relative(s) staying permanently with the patient was 900 rupees. School-going patients were absent from school on average 67 days. Households used one or more strategies to cover the costs of treatment: (1) using available cash and savings, (2) taking loans, (3) the sale of assets and/or rental of land and (4) gifts.

Forty-nine per cent of patients covered expenditures by using available cash and savings. However, the amount of money available was insufficient to cover all expenditures and a vast majority of patients (81%) had taken a loan. The median amount borrowed was 4000 rupees

with a reported monthly interest of 5% for most respondents (range 0–10%). The monthly interest payment was 200 rupees. Ninety-one per cent borrowed the money from someone from the same village as the patient, either a neighbour or a moneylender, while 9% got a loan from a friend or a relative. Only five patients (6%) covered all expenditures with available cash and savings. Nine per cent of patients temporarily rented a part of their land in exchange for cash and could regain ownership of their land once they had repaid the loan. One patient sold a parcel of land which resulted in a permanent loss of income. Four per cent sold livestock or poultry while 13% mentioned they would be unable to pay all costs.

Health seeking behaviour and costs prior to the KAMRC

The survey also collected information on household costs and health seeking behaviour prior to admission at the KAMRC (Table 3). Patients visited on average two different providers prior to seeking/receiving care the KAMRC. Only in 24% of cases, these providers suspected/diagnosed the patient to be infected with VL and referred the patient to the KAMRC for specialised treatment. In all other cases, the patient was referred to the centre by either a relative, a member of the same village or the patient knew about the centre him/herself. The total (direct) cost of seeking/receiving care for VL prior to the KAMRC was estimated at 1220 rupees (range 75–10 530 rupees). Medical costs (consultation fees, drugs and tests) constituted 89% and non-medical costs (transportation costs) 11% of total costs.

Table 3: Patients costs and health seeking behaviour prior to the Kala-azar Medican Research Centre

| Cost items | Median (INR) | (IQR ^a) |
|-------------------|--------------|---------------------|
| Medical costs | 1090 | (650-2450) |
| Non-medical costs | 120 | (620-2300) |
| Total (rupees) | 1220 | |
| Total (US\$) | 28 | |

| Type of provider consulted | No. of patients | (%) |
|----------------------------|-----------------|------|
| Private doctor/hospital | 49 | (64) |
| Compounder (i.e. quack) | 22 | (28) |
| Public health facility | 5 | (7) |
| Traditional healer | 1 | (1) |

^a Interquartile range

DISCUSSION

This study is a first attempt to estimate the direct and indirect costs associated with patient management of VL from a societal and household perspective in Bihar. Data were collected on provider’s costs and a structured questionnaire was administered to 77 patients to estimate their direct and indirect costs to receive treatment for VL and was supplemented with information from patient records to improve reliability. The indirect costs were calculated using the human capital method and were reported separately as recommended by the WHO (Tan-Torres Edejer *et al.* 2003).

From our findings, the indirect cost, mainly the loss of income to the patient and relatives, of seeking/receiving care represents the highest cost item. This high indirect cost follows from the long duration of illness and is explained by the fact that patients visited on average two different providers prior to seeking/receiving care at the KAMRC. The importance of indirect costs has been reported in various studies from Africa (Sauerborn *et al.* 1991; Ettling *et al.* 1994).

The results on the health seeking behaviour and the costs incurred by patients prior to the KAMRC should be interpreted with care and should be considered as indicative results rather than robust findings because of recall bias. Nevertheless it shows that a considerable amount of time elapses between the first signs of VL and the time patients receive treatment for VL. Moreover, in only 24% of cases previous providers suspected/diagnosed the patient to be infected with VL and referred the patient to the KAMRC for specialised treatment. Awareness on VL among health professionals should therefore be increased to ensure a fast and accurate diagnosis of VL in the early stage of disease. This will decrease the risk of misdiagnosis and mistreatment and therefore result in a decreased absence from productive activities and reduce the loss of income to the patient.

A few limitations to this study should be mentioned when attempting to generalise our findings. First, the study is based on only one health care facility in an urban centre and costs were collected in a short period of evaluation with a relatively small sample size. The choice of the study site can lead to selection bias as all patients were selected from the same hospital. As a result, the patients described in this study may not be representative for the entire group affected by VL. Although some studies (Desjeux 1996; Guerin *et al.* 2002) suggest an occupational hazard explaining the high number of infected males, there may well be a bias at the level of the household as described by Thakur (2000) where women and young children are not sent to the hospital for care because of the high cost of treatment and the financial burden it

imposes on households. Second, the costs collected at the KAMRC, a charitable hospital, might not be representative for other hospitals in the region or for other regions. The treatment costs in a private clinic (providing health care to an estimated 80% of all detected VL cases in Bihar state) are likely to be substantially higher because actual cost for medications, tubing's, infusions, laboratory investigations and hospital beds will be significantly higher. Similarly, medical care fees like consultation and visiting fees, fees for infusion of the drug will further add to the cost of treatment. An important variable determining costs is the type and duration of treatment and the process of patient management which can be different over hospitals. In this study patients were treated with conventional Amphotericin B deoxycholate where a majority of patients (88%) received infusions on a daily basis and were admitted to the hospital for the full duration of treatment. In other settings, such as in the BPKIHS hospital in Dharan (Nepal), where antimonial treatment is still in use, patients are admitted for the first week and thereafter managed on an outpatient basis (Rijal *et al.* 2003) while in other centres patients are even treated on a complete outpatient basis.

Despite the limitations mentioned above, this study provides valuable information on the total cost of VL and shows that the economic costs related to VL are substantial.

The study shows that patients incur substantial costs to seek and receive treatment for VL. Patients visit several health care providers, receive and pay for inefficient treatments, subsequently depleting the little savings they have, before being diagnosed with VL and receiving appropriate treatment. Most households are forced to take on a loan with high interest rates or sell assets such as livestock to cover the costs of VL treatment. This pattern has also been observed in recent studies on the economic impact of VL in Bangladesh (Ahluwalia *et al.* 2003; Sharma *et al.* 2006) and Nepal (Rijal *et al.* 2006) as well as studies analysing out-of-pocket expenditures and debt in other parts of Asia (Van Damme *et al.* 2004) and in Africa (Sauerborn *et al.* 1996; Mugisha *et al.* 2002). VL leads to catastrophic health expenditures and further impoverishes households but also keeps them into poverty because of the heavy financial commitments on the long term because of indebtedness (Meessen *et al.* 2003).

As VL mainly occurs in poor rural households with weak medical facilities (Bryceson 2001), public health authorities in Bihar should focus on policies that detect VL in the early stage of disease and implement interventions that minimize the burden to households affected by VL. We see a role for the government to ensure access of care for poor patients (Baltussen 2005) because of the high (economic and financial) costs to the patient; we also recommend that public hospitals subsidise treatment of VL to the patient.


More research is required to gain a better understanding of the health seeking behaviour of patients, as well as the socioeconomic impact of VL on households. Globally, despite the development of new drugs such as Paromomycin and Miltefosine, the challenge remains formidable and requires increased and continued investment from the international community to encourage the development of new and improved tools for control of VL, and neglected diseases in general.

Acknowledgements

Financial support for this study was received from the Institute of Tropical Medicine, Antwerp (Belgium) and from the Drugs for Neglected Diseases Initiative (DNDi). The authors would like to thank Kamlesh Gidwani for the provided assistance during the patient surveys and also the health and administrative personnel at the KAMRC for their willingness to provide the necessary inputs for the study.

REFERENCES

- Ahluwalia IB, Bern C, Costa C, et al. (2003) Visceral leishmaniasis: consequences of a neglected disease in a Bangladeshi community. *Am J Trop Med Hyg* 69: 624–28.
- Asenso-Okyere WK & Dzator JA (1997) Household cost of seeking malaria care: a retrospective study of two districts in Ghana. *Soc Sci Med* 45: 659–67.
- Baltussen R (2005) Priority setting of public spending in developing countries: do not try to do everything for everybody. *Health Policy* 78: 149–56.
- Barnum H & Kutzin J (1993) *Public hospitals in developing countries: resource use, cost, financing*. Baltimore, MD: John Hopkins University Press.
- Brouwer W, Rutten F & Koopmanschap M (2001) *Costing in economic evaluations*. In: Drummond M, McGuire A (eds.) *Economic evaluation in health care: merging theory with practice*, 2nd edn. Oxford: Oxford University Press. pp. 68–93.
- Bryceson A (2001) A policy for leishmaniasis with respect to the prevention and control of drug resistance. *Trop Med Int Health* 6: 928–34.
- Cho-Min-Naing ML, Gatton ML (2004) Costs to the patient for seeking malaria care in Myanmar. *Acta Trop* 92: 173–77.
- Creese A, Parker D (1994) *Cost analysis in primary health care: a training manual for programme managers*. Geneva: World Health Organization.
- Desjeux Ph (1996) Leishmaniasis: public health aspects and control. *Clin Dermatol* 14: 417–23.
- Drummond MF, O'Brien B, Stoddart GL, Torrance GW (1997) *Methods for the Economic Evaluation of Health Care Programmes*. Oxford University Press, Oxford.
- Duraisamy P, Ganesh AK, Homan R, et al. (2006) Costs and financial burden of care and support services to PLHA and households in South India. *AIDS care* 18: 121–27.
- Ettling M, McFarland DA, Schultz LJ, Chitsulo L (1994) Economic impact of malaria in Malawian households. *Trop Med Parasitol* 45: 74–79.
- Guerin PJ, Olliaro P, Sundar S, et al. (2002) Visceral leishmaniasis: current status of control, diagnosis, and treatment, and a proposed research and development agenda. *Lancet Infect Dis* 2: 494–501.
- Hutton G, Baltussen R (2005) Cost valuation in resource poor settings. *Health Policy Plann* 20: 252–59.
- Koopmanschap MA, Rutten FFH, Ineveld BM, van Roijen L (1995) The friction cost method of measuring the indirect costs of disease. *J Health Econ* 14: 171–89.
- Meessen B, Zhenzhong Z, Van Damme W, Devadasan N, Criel B, Bloom G (2003) Iatrogenic poverty. *Trop Med Int Health* 8: 581–84.
- Mugisha F, Kouyate B, Gbangou A, Sauerborn R (2002) Examining out-of-pocket expenditure on health care in Nouna, Burkina Faso: implications for health policy. *Trop Med Int Health* 7: 187–96.
- Murray HW (2000) Treatment of visceral leishmaniasis (Kala-Azar): a decade of progress and future approaches. *Int J Infect Dis* 4: 158–77.
- Murray HW (2004) Treatment of visceral leishmaniasis in 2004. *Am J Trop Med Hyg* 71: 787–94.
- Olliaro P, Guerin PJ, Gerstl S, Haaskjold AA, Rottingen JA, Sundar S (2005) Treatment options for visceral leishmaniasis: a systematic review of clinical studies done in India 1980–2004. *Lancet Infect Dis* 5: 763–74.
- Rijal S, Chappuis F, Singh R, et al. (2003) Treatment of visceral leishmaniasis in south-eastern Nepal: decreasing efficacy of sodium stibogluconate and need for a policy to limit further decline. *Trans R Soc Trop Med Hyg* 97: 350–54.
- Rijal S, Koirala S, Van der Stuyft P, Boelaert M (2006) The economic burden of visceral leishmaniasis for households in Nepal. *Trans R Soc Trop Med Hyg* 100: 838–41.
- Sauerborn R, Shepard DS, Ettling MB, Brinkmann U, Nougara A, Diesfeld HJ (1991) Estimating the direct and indirect economic costs of malaria in a rural district of Burkina Faso. *Trop Med Parasitol* 42: 219–23.
- Sauerborn R, Adams A & Hien M (1996) Household strategies to cope with the economic costs of illness. *Soc Sci Med* 43: 291–301.
- Sharma A, Bern C, Varghese B, Chowdhury R, Haque R, Ali M (2006) The economic impact of visceral leishmaniasis on households in Bangladesh. *Trop Med Int Health* 11: 757–65.
- Sculpher M (2001) *The role and estimation of productivity costs in economic evaluation*. In: Drummond M, McGuire A (eds.) *Economic evaluation in health care: merging theory with practice*, 2nd edn. Oxford: Oxford University Press. pp. 94–112.
- Sundar S (2001) Drug resistance in Indian visceral leishmaniasis. *Trop Med Int Health* 6: 849–54.
- Sundar S, Agrawal G, Rai M, Makharia MK, Murray HW (2001) Treatment of Indian visceral leishmaniasis with single or daily infusions of low dose liposomal amphotericin B: randomized trial. *BMJ* 323: 419–22.
- Sundar S, Mehta H, Suresh AV, Singh SP, Rai M, Murray HW (2004) Amphotericin B treatment for visceral leishmaniasis: conventional versus lipid formulation. *Clin Infect Dis* 38: 377–83.
- Tan-Torres Edejer T, Baltussen R, Adam T, et al. (2003) *Making choices in health: WHO guide to cost-effectiveness analysis*. Geneva: World Health Organization.
- Thakur CP (2000) Socio-economics of visceral leishmaniasis in Bihar (India). *Trans R Soc Trop Med Hyg* 94: 156–57.
- Van Damme W, Van Leemput L, Por I, Hardeman W, Meessen B (2004) Out-of-pocket expenditure and debt in poor households: evidence from Cambodia. *Trop Med Int Health* 9: 273–80.
- WHO (2000) *The leishmaniasis and leishmania/HIV co-infections*. Fact sheet n° 116, May 2000. Geneva: World Health Organization. [online] <http://www.who.int/mediacentre/factsheets/fs116/en/>.
- WHO (2005) *Guidelines for estimating the economic burden of diarrhoeal disease, with focus on assessing the costs of rotavirus diarrhoea*. Document WHO/IVB/05.10. Geneva: World Health Organization.
- World Bank (2005) Bihar: towards a development strategy. World Bank Report 2005.



CHAPTER 5

The household costs of visceral leishmaniasis care in south-eastern Nepal

Uranw S[§]
Meheus F[§]
Baltussen R
Rijal S
Boelaert M

Shared first authorship [§]

ABSTRACT

Background and objectives

Visceral leishmaniasis (VL) is an important public health problem in south-eastern Nepal affecting very poor rural communities. Since 2005, Nepal is involved in a regional initiative to eliminate VL. This study assessed the economic impact of VL on households and examined whether the intensified VL control efforts induced by the government resulted in a decrease in household costs.

Methods

Between August and September 2010, a household survey was conducted among 168 patients that had been treated for VL within 12 months prior to the survey in 5 districts in south-eastern Nepal. We collected data on health seeking behaviour, direct and indirect costs and coping strategies.

Results

The median total cost of one episode of VL was US\$ 165 or 11% of annual household income. The median delay between the onset of symptoms and presentation to a qualified provider was 25 days. Once the patient presented to a qualified provider, the delay to correct diagnosis was minimal (median 3 days). Direct and indirect costs (income losses) represented 47% and 53% of total costs respectively. Households used multiple strategies to cope with the cost of illness, mainly mobilizing cash/savings (71%) or taking a loan (56%).

Conclusions

The provision of free VL diagnosis and drugs by the Nepalese control programme has been an important policy measure to reduce the cost of VL to households. But despite the free VL drugs, the economic burden is still important for households. More efforts should be put in reducing indirect costs, in particular the length of treatment, and prevent the transmission of VL through vector control.

INTRODUCTION

Since 2005, the Government of Nepal has been involved in a Visceral Leishmaniasis (VL) elimination programme, alongside the governments of India and Bangladesh, to reduce annual VL incidence to less than one case per 10,000 population by 2015 (WHO 2005). VL, also known as kala-azar, is a parasitic disease that is fatal if treatment is not provided timely. The disease is transmitted from humans-to-humans through the bite of a female sandfly.

Geographically, VL occurs in the alluvial plains of the river Ganges, in districts bordering the frontiers between Bangladesh, India and Nepal. The cases in this region account for 60% of the global burden of VL. In Nepal, 8 million people are at risk of acquiring VL in 12 districts in the central and eastern regions of the country (Joshi et al. 2006). Between 2000 and 2010, 17,462 cases and 244 deaths were reported in Nepal although these figures, obtained through passive case surveillance at government health facilities, are likely to be underestimations since many cases are not reported or remain undetected (Epidemiology and Disease Control Division 2010; Singh et al. 2006; Singh et al. 2010).

The cornerstones of the VL elimination initiative are early detection and appropriate treatment, in an attempt to curtail transmission of the disease. A standardized clinical case definition was adopted, the rK39 rapid diagnostic test was introduced to enable faster detection of suspected cases and miltefosine, the first oral drug for VL, replaced sodium stibogluconate (SSG) to which growing failure rates had been reported in Nepal (Rijal et al. 2003). In addition, both VL diagnosis and drugs are now provided free of charge at public health care facilities. These measures along with enhanced vector control resulted in a steady decrease in the annual number of reported VL cases in Nepal from 2,229 in 2003 to 900 in 2010. Another important outcome of the improved case management strategy may be its effect on the household costs of seeking and obtaining appropriate VL care. VL is a disease of poverty affecting the poorest of the poor (Alvar et al. 2006; Boelaert et al. 2009). Households with low incomes and living in precarious housing conditions such as mud -or grass covered houses are most at risk of acquiring VL disease (Singh et al. 2010; Uranw et al. unpublished data). The few studies in Nepal quantifying the economic burden of VL on households and conducted prior to the elimination initiative showed a VL episode to profoundly impact the socio-economic status of the household. Adhikari et al. (2009) reported that up to 26% of previously non-poor households were pushed into poverty as a direct result of out-of-pocket expenditures on VL care while Rijal et al. 2006 showed that the (median) direct and indirect costs of a VL episode were equal to one year of median per capita income. Usually an expenditure exceeding 10% of annual household income is defined as catastrophic, meaning it drives households into destitution (O'Donnell et al. 2008; Ranson

2002). These high costs are caused, amongst others, by long delays, up to 2 months, before correct diagnosis whereby households consulted traditional and private-for-profit providers (Rijal et al. 2006) as well as the long hospitalization due to the use of SSG. Faster screening of suspected cases, enhanced access to free treatment, and a different treatment regimen may reduce these household costs. We examined whether the intensified VL control efforts induced by the elimination initiative resulted in a demonstrable impact at household level in terms of health seeking behaviour, costs and coping strategies.

METHODS

Nepal is administratively divided into 14 zones and 75 districts. In 2010, VL was reported in 12 districts situated in south-eastern Nepal in the Terai region bordering the highly VL-endemic northern state of Bihar in India. The study was conducted between August and September 2010 in five of these highly endemic districts, namely Siraha, Saptari, Sunsari, Morang and Jhapa. VL incidence rates in the study districts varied from 0.52 cases per 10,000 persons per year (Sunsari district) to 2.03 cases per 10,000 persons per year (Saptari district) in 2010 (table 1).

Table 1: Characteristics of districts included in the study

| | Siraha | Saptari | Sunsari | Morang | Jhapa |
|-----------------------------------------------|---------------|------------|------------|------------|------------|
| Location | Central Terai | East Terai | East Terai | East Terai | East Terai |
| Population, 2006 | 638,375 | 633,965 | 710,842 | 941,614 | 755,494 |
| VL cases, 2006* | 142 | 255 | 117 | 113 | 52 |
| Case detection rate# | 22.2 | 40.2 | 16.5 | 12 | 6.9 |
| Cases in survey (% of notified cases in 2006) | 8 (5.6) | 54 (21.2) | 12 (10.3) | 78 (69.0) | 16 (30.8) |

Source: National Population & Housing Census 2011; adapted from Mills 1993

*Notified to Epidemiology and Diseases Control Division, Nepal

Number of new cases reported per 100 000 person-years

We searched the medical records of the District Public Health Office in each district and the database of the B.P. Koirala Institute of Health Sciences (BPKIHS) to identify all households in the five districts with a household member treated for kala-azar within 12 months prior to the survey. BPKIHS is a tertiary level hospital situated in Sunsari district and draws many patients from the surrounding areas due to its widespread reputation as a VL treatment and research centre. Patients treated at BPKIHS are not included in the medical records of the District Public Health Office. Furthermore, to minimize recall bias, we only considered the most recent case of VL in the household.

Organization of VL care services

Health care services at the district level are provided by sub-health posts, health posts, primary health care centres and district hospitals (i.e. primary care level) (WHO 2007). A network of female community health volunteers at the village level refer patients to health -and sub-health posts. Patients suspected of VL (defined as individuals with a history of fever of more than 2 weeks with a palpable spleen) seeking care from female community health volunteers, sub-health posts and health posts are referred to primary health care centres or district hospitals for diagnosis by a rapid diagnostic test (rK39 immunochromatographic strip test). If positive, the patient is treated at the PHC or referred to a district hospital (or higher level) if the PHC does not have a medical doctor, which is often observed. All VL drugs are provided free of costs. Diagnosis through parasitology (bone marrow or splenic aspiration) can only be done at district hospitals or above. While private formal providers such as private clinics also provide diagnosis and treatment of VL, free treatment is only available at public facilities. Various anti-leishmanial drugs are available in Nepal: since 2006 SSG, administrated intramuscularly for 30 days, was replaced as first line treatment by miltefosine, an oral drug given for 28 days. Due to its possible teratogenic effect, miltefosine is not given to pregnant women. The second line treatment is amphotericin B deoxycholate given every day for 14 doses.

Data collection & analysis

Information for the study was obtained from patient medical records and a household questionnaire. Medical records at the District Public Health Offices and the BPKIHS were consulted to retrieve data on the type of VL drugs received and the length of treatment. Subsequently households were visited at their homestead by a team of trained field workers who had previously been involved in other kala-azar related community and household surveys in the area. The field workers were supervised on a daily basis by the first author (S. Uranw). They used a pre-tested structured questionnaire administered to the head of the household or the most knowledgeable person. The survey collected data on treatment seeking behaviour (health providers visited, mode of travel, delay to presentation to first qualified health professional, etc.), direct and indirect costs and the coping strategies to meet the health seeking and treatment costs.

Direct medical and non-medical cost data were gathered for each provider visited. Direct medical costs included all out-of-pocket expenditures by the household on consultation, medicines and laboratory tests. Direct non-medical costs included expenditure on transportation to and from the health facility, food costs and other daily expenditures for the patient and accompanying family members. The indirect cost of a VL episode represented the loss of productivity within the household due to illness and was estimated using the human capital approach. The loss of

productivity was valued in terms of the loss of earnings of the patient and household members caring for the patient (either at home or hospital). For patients and attendants, the daily wage rate was estimated and multiplied by the number of work days lost to obtain the indirect cost of a VL episode. The daily wage rate was determined by asking a series of questions on the daily monetary income (the main source of income to most household members was daily labour). For patients and attendants reporting farming as their main source of income, the survey collected data on the yearly production of each produce which was then valued with local market prices and divided by the number of agriculturally active household members. We also estimated total household income as the sum of monthly cash income from daily labour for each economically active household member, the income from agriculture, income from sales of animals and animal products (e.g. milk) and remittances from family members.

Analysis

The data is described using descriptive statistics showing proportions, means and standard deviations. We also presented medians and interquartile ranges (25th and 75th percentile) because of skewed distributions in the cost data; many households reported zero out-of-pocket expenditure for some cost categories and providers, in particular informal health care providers. Costs were defined as catastrophic if they exceeded 10% of annual household income (O'Donnell et al. 2008; Ranson 2002). All costs were converted from Nepalese rupees (Rs.) to US dollars using the exchange rate prevailing at the time of the study (1 USD = Rs. 74.8; OANDA August 2010). Data entry and cleaning were done in Microsoft Excel and analysis in STATA v10.1 (Stata Corp., College Station Tx, USA).

Ethical considerations

Ethical clearance was obtained from the ethics committee of the BP. Koirala Institute of Health Sciences, Nepal and the ethics committee of the University of Antwerp, Belgium. Patient's medical records were reviewed retrospectively and all information retrieved from medical records was anonymized. Signed informed consent was obtained from all adult patients and from a parent or guardian of participating minors. All households that were approached for the study, whether they accepted to participate or not, received a free long-lasting insecticide treated net (Vestergaard Frandsen A/S, Denmark) as a compensation for their time spent with the survey team.

RESULTS

Characteristics of study participants and the household

We randomly retrieved a total of 200 households where a case of VL occurred in the past 12 months in the five districts, of which 168 households were located by field workers and accepted to be interviewed. The majority of patients were male (60%) and 41 of them were head of the household (24%). Most patients were over 14 years of age (68%); the median age was 19 (IQR 12.5-35). Male patients were significantly older than female (median age 22.5 versus 17.0; $p < 0.05$). The percentage of women of childbearing age (15-49 years) among patients was 19%. The median household size was 5.7 persons.

Out of 168 patients, sixty-one (36%) were economically active at the time of illness, most of them day labourers (80%) such as rickshaw driver or farm labourers. Few patients were engaged in small-scale farming ($n=5$; 8%) or were salaried workers ($n=4$; 6%). The median monthly income of an economically active patient ($n=60$) was Rs. 6,000 or 81 US\$ (range 27-162 US\$). The median monthly income of the household ($n=168$) was Rs. 10,243 or 138 US\$ (range 87-163 US\$) giving a median per capita monthly income of Rs. 1,882 or 25 US\$ (range 20-30 US\$).

The vast majority of households lived in non-permanent housing structures either consisting entirely of natural materials (49%) or a combination, usually mud walls and metal sheets as roof (45%). Eighteen per cent of households owned land ($n=30$); 27 of these households cultivated some crops on their land, mainly paddy rice and wheat. Most households also owned some livestock (83%), usually goats (56% of households; median 3 heads), cows (44%, median 2.5) or chickens (36%, median 5). Fifty-eight per cent of households owned a bicycle, 56% a mobile phone and 36% a radio.

Health-seeking behaviour

Patients visited a median of 2 health providers including the one who eventually treated them (IQR 1-2) (table 2). For 91 households (55%), a public provider was the patient's first point of contact; other households first visited a private (qualified) provider ($n=34$; 20%), a traditional healer ($n=26$; 15%) or a chemist or pharmacy ($n=16$; 10%). The main reasons behind the choice of the first provider were proximity (49%) and the perceived (good) reputation of the health provider (38%). Traditional healers were chosen for their proximity, public providers (health centre or hospital) most often for their reputation while for private providers it was a mix of both. Ninety patients (54%) were submitted to a VL diagnostic test on their first visit but this varied considerably by type of provider: all patients presenting at public hospitals were

tested for VL, compared to 56% at public health centres and 39% at private providers. Of the patients that did not receive a VL diagnostic test at a public health centre or a private provider on their first visit, respectively 60% and 55% were subsequently referred by the provider to a public hospital for testing and treatment. Households that used the services of an unqualified provider first, were more likely to visit either another unqualified provider or a private provider afterwards. Approximately 21% (n=35) of households visited three different providers; 4% (n=7) of households visited 4 different types of health providers.

The median delay between the onset of symptoms and presentation to a qualified health provider (i.e. patient delay) was 25 days (IQR 20-30). Once the patient had presented to a qualified provider, the median delay to correct diagnosis of VL was 3 days (IQR 2-7). The total median delay from onset of symptoms to start of treatment was 31 days (IQR 23-35). While none of the delays varied with age or gender, there was a significant and positive relationship between the total delay and the number of providers visited ($p < 0.01$).

Treatment regimens

The vast majority of patients in our study were treated with either miltefosine (83%) or conventional amphotericin B (15%) in case of relapse as recommended by the 2005 guidelines of the VL elimination initiative (WHO 2005). Four patients were treated with SSG (2%), two of them by a public provider. The other two patients treated with SSG by a private provider were unexpected in our survey because these patients are usually not included in the DPHO records. In addition, both patients reported not to have paid for the SSG drugs. Upon closer inspection, these patients were from the same village in Morang district on the border with Bihar state (India). After discussion with the local vector control officer, they had probably obtained the SSG free of charge from a private charitable hospital in Bihar and subsequently received the injections at a health facility in Nepal by trained health workers.

Direct costs

The average and median direct household costs incurred by type of provider are given in table 3. All but three patients received VL treatment at a public hospital, the remaining 3 patients were treated at a public health centre (n=1) or a private qualified provider (n=2).

The median direct cost of an episode of VL across all providers was Rs. 4,905 (IQR 3,025-7,125) or US\$ 66 (IQR 41-96). Direct medical costs were Rs. 2,390 (IQR 1,100-4,290) and non-medical costs Rs. 2,300 (IQR 1,550-3,350) or 51% and 49% respectively of total median direct costs. Direct medical costs arose from expenditures on consultation fees, miscellaneous drugs and laboratory investigations (including diagnostic tests). The survey confirmed that none of the

Table 2: Health seeking behaviour of households (n=168)

| Variable | N° (%) of patients |
|-----------------------------------------------------------|--------------------|
| Type of health provider first visited | |
| Traditional | 26 (15.5) |
| Chemist or pharmacy | 16 (9.5) |
| Village health worker | 6 (3.6) |
| Public, primary | 23 (13.7) |
| Public, hospital | 63 (37.5) |
| Private doctor/clinic | 34 (20.2) |
| Delay to presentation (in days) (median; IQR) | 25 (20-30) |
| Delay to diagnosis (in days) (median; IQR) | 3 (2-7) |
| Number of health providers visited | |
| 1 | 63 (37.5) |
| 2 | 70 (41.6) |
| 3 | 28 (16.7) |
| 4 | 7 (4.2) |
| Diagnosed with VL at first visit? | |
| Yes | 90 (53.6) |
| No | 78 (46.4) |
| Mode of transportation to first facility | |
| Foot | 38 (22.6) |
| Bicycle | 28 (16.7) |
| Bus | 96 (57.1) |
| Other | 6 (3.6) |
| Mode of transportation to treatment facility ¹ | |
| Foot | 4 (2.4) |
| Bicycle | 10 (6.0) |
| Motorbike | 2 (1.2) |
| Bus | 148 (88.6) |
| Other | 3 (1.8) |
| Distance between home and treatment facility (kilometers) | |
| < 20 | 45 (26.8) |
| 20-60 | 86 (51.2) |
| > 60 | 37 (22.0) |

¹ The treatment facility is the health provider where the patient received VL treatment. In 98% of cases this was a public hospital.

households had to pay for VL drugs. Median direct medical costs were highest for households visiting private providers (median Rs. 2000; IQR 1,475-3,575), in particular payments for ancillary drugs (e.g. antibiotics, antipyretics or vitamin injections). The direct non-medical costs, consisting of transportation, food and other expenses (i.e. small daily expenses) were highest at the public hospital. The high food costs at public hospitals (median: Rs. 1,400; IQR 700-2,000) arose from the hospitalization of the patient and accompanying family member(s) for VL treatment. The median duration of hospitalisation was 10 days (IQR 7-16) and was the same for patients receiving SSG or miltefosine but higher for patients treated with conventional amphotericin B (median: 14 days; IQR 8-20). Direct (medical and non-medical) costs did not vary by gender or income quintiles, but direct medical costs increased with the patient's age.

Indirect costs

VL is a syndrome characterized by prolonged fever, weight loss, anaemia, fatigue and enlargement of the liver and spleen. As a result patients are either severely limited or not able at all to carry out their daily activities and need much support from family members. Among the 168 patients, 95% (n=160) reported that VL illness had a severe impact on their normal functioning and resulted in a loss of income to the household, either wage losses to the patient or caretakers, losses in agricultural output or other earnings. Patients reported not being able to carry out their normal daily activities for a median number of 57 days (IQR 51-65) (table 4). As a result the median loss of income was Rs. 12,400 for economically active patients. Since only 36% of patients were economically active, the value of time lost across all patients, both the economically active and non-active, was on average Rs. 4,731.

Patients were attended by on average 1.1 household members (range: 1-2). These caretakers reported a median loss of 15 workdays (IQR 10-30) mainly due to accompanying the patient to the various health providers and staying with him/her for the full duration of hospitalization. The median loss of income to caretakers was Rs. 2,583 (on average Rs. 2,279 across all caretakers). The median total value of time lost to the household per episode of VL was Rs. 4,500 (IQR 1,500-12,167).

Table 3: Direct medical and non-medical costs of treatment per patient by type of provider (Rs. 2010)

| | Traditional (n=28) | | Chemist/pharmacy (n=25) | | Village health worker (n=6) | |
|--------------------------------|--------------------|--------------------|-------------------------|--------------------|-----------------------------|--------------------|
| | Mean (sd) | Median (IQR 25-75) | Mean (sd) | Median (IQR 25-75) | Mean (sd) | Median (IQR 25-75) |
| Direct medical costs | | | | | | |
| Consultation | 210 (363) | 50 (0-200) | 74 (76) | 50 (10-100) | 72 (114) | 30 (10-60) |
| Ancillary drugs | 425 (847) | 0 (0-750) | 1,242 (1,526) | 700 (500-1,400) | 657 (242) | 670 (500-800) |
| Laboratory investigations | 45 (137) | 0 (0-0) | 225 (218) | 100 (0-400) | 67 (103) | 0 (0-200) |
| Total direct medical costs | 679 (973) | 300 (0-1,000) | 2,034 (2,950) | 1,150 (500-2,000) | 795 (325) | 785 (650-1,000) |
| Direct non-medical costs | | | | | | |
| Transportation | 7 (30) | 0 (0-0) | 113 (194) | 0 (0-100) | 0 (0) | 0 (0-0) |
| Food | 314 (567) | 0 (0-350) | 107 (149) | 0 (0-200) | 17 (26) | 0 (0-50) |
| Other | 0 (0) | 0 (0-0) | 2 (10) | 0 (0-0) | 8 (20) | 0 (0-0) |
| Total direct non-medical costs | 321 (564) | 0 (0-350) | 222 (319) | 100 (0-200) | 25 (42) | 0 (0-50) |
| Total direct costs | 1,001 (1,150) | 775 (75-1,700) | 2,256 (3,158) | 1,150 (500-2,200) | 820 (350) | 785 (700-1,000) |

Table 3: Continued

| | Public, primary (n=32) | | Public, hospital (n=165) | | Private doctor/clinic (n=52) | |
|---------------------------------------|------------------------|--------------------|--------------------------|---------------------|------------------------------|---------------------|
| | Mean (sd) | Median (IQR 25-75) | Mean (sd) | Median (IQR 25-75) | Mean (sd) | Median (IQR 25-75) |
| Direct medical costs | | | | | | |
| Consultation | 56 (85) | 25 (10-50) | 79 (191) | 50 (25-70) | 222 (155) | 200 (200-250) |
| Ancillary drugs | 1,002 (845) | 600 (500-1,345) | 1,120 (1,580) | 650 (400-1,080) | 1,714 (1,567) | 1,200 (825-2,000) |
| Laboratory investigations | 317 (349) | 200 (95-425) | 442 (603) | 300 (200-500) | 748 (767) | 500 (300-875) |
| Total direct <i>medical</i> costs | 1,375 (1,107) | 975 (608-1,900) | 1,550 (1,703) | 1,000 (720-1,688) | 2,684 (2,038) | 2,000 (1,475-3,575) |
| Direct non-medical costs | | | | | | |
| Transportation | 75 (114) | 23 (0-100) | 431 (343) | 450 (200-500) | 390 (392) | 300 (150-500) |
| Food | 123 (144) | 100 (0-200) | 1,424 (968) | 1,400 (700-2,000) | 414 (571) | 200 (100-500) |
| Other | 29 (97) | 0 (0-0) | 239 (216) | 200 (100-400) | 68 (183) | 0 (0-0) |
| Total direct <i>non-medical</i> costs | 226 (290) | 125 (23-275) | 2,090 (1,207) | 2,000 (1,300-2,700) | 882 (1,007) | 600 (300-1,100) |
| Total direct costs | 1,601 (1,266) | 1,075 (693-2,260) | 3,640 (2,531) | 3,130 (2,135-4,223) | 3,582 (2,545) | 2,500 (2,000-5,000) |

| | Total costs all providers | | |
|---------------------------------------|---------------------------|------|---------------------|
| | Mean | (sd) | Median (IQR 25-75) |
| Direct medical costs | | | |
| Consultation | 203 (297) | | 88 (40-270) |
| Ancillary drugs | 2,134 (2,137) | | 1,500 (700-2,850) |
| Laboratory investigations | 786 (872) | | 600 (300-1,000) |
| Total direct <i>medical</i> costs | 3,123 (2,745) | | 2,390 (1,100-4,290) |
| Direct non-medical costs | | | |
| Transportation | 574 (481) | | 500 (300-800) |
| Food | 1,619 (1,018) | | 1,500 (1,000-2,250) |
| Other | 260 (252) | | 200 (100-480) |
| Total direct <i>non-medical</i> costs | 2,453 (1,400) | | 2,300 (1,550-3,350) |
| Total direct costs | 5,576 (3,552) | | 4,905 (3,025-7,125) |

Table 4: Indirect costs (Rs. 2010)

| | Mean | (sd) | Median | (IQR 25-75) |
|----------------------------------------------------------|--------------|----------------|--------------|-----------------------|
| Patients' duration of illness (days) [§] | 60 | (18) | 57 | (51-65) |
| Number of attendants per patient | 1.1 | (0.3) | 1.0 | (1.0-1.0) |
| Workdays lost by attendants | 21 | (16) | 15 | (10-30) |
| Loss of income; working patients only (n=61) | 13,030 | (5,638) | 12,400 | (9,800-15,400) |
| Loss of income; all patients (n=168) | 4,731 | (7,136) | 0 | (0-10,700) |
| Loss of income working attendants only (n=134) | 3,112 | (2,300) | 2,583 | (1,500-4,000) |
| Loss of income; all attendants (n=183) | 2,279 | (2,404) | 1,500 | (0-3,100) |
| <i>Total loss of income to the household[£]</i> | <i>7,213</i> | <i>(7,217)</i> | <i>4,500</i> | <i>(1,500-12,167)</i> |
| Total payment on loan* | 2,611 | (2,176) | 2,080 | (1000-3,300) |
| Total indirect cost | 8,084 | (7,391) | 5,167 | (3,000-13,290) |

[§] Consists of the various types of delays plus the treatment duration
[£] Across all patients & attendants
* For those with interest payments

Coping strategies

Households used a number of strategies to cope with the costs of VL illness. Many of these strategies resulted in additional costs to the household; e.g. in terms of interest payments on loans or hiring labour to replace the sick household member. The survey identified three strategies to cope with the financial costs of VL illness: mobilizing cash/savings, taking a loan and sales of livestock. Mobilizing cash or savings was the most frequent coping strategy. Seventy-one per cent (n=120) of households used their savings to pay for health care expenditures, although for 54% of these households it was not enough to cover all medical costs. Out of those 120 households, 75% (n=90) of households reported that the savings were supposed to buy food, in other cases assets (n=30). Fifty-six per cent (n=94) of households took a loan to finance the costs of care, most often from a member of the same village (71%), followed by friends or peers (17%) or an informal money lender (6%). When borrowing from friends, the loan was interest free. In other cases (n=57), the amount to be repaid was on average 140% the original amount borrowed, usually through monthly instalments. A collateral was not often provided for the loan: four households provided assets as collateral, one household that took a loan from a bank provided their house. Seventeen per cent of households sold livestock to cover the costs of care (n=29). Forty-two per cent of households chose more than one strategy (n=71), mainly using cash/savings and a loan (n=47), while 2% (n=4) had to revert to all three strategies to cope with the costs of VL illness.

Households also used various strategies to compensate the labour lost due to VL illness. Nine households hired external labour to replace either the patient or caretakers in the field at a rate of Rs. 200 per day for a median of 60 days (IQR 60-60). Twenty-three per cent of patients were replaced by a family member that was a school-going child for the duration of their illness (n=14).

Table 5: Summary of direct and indirect costs per VL episode (Rs. and in US\$ 2010)

| Item | Mean | | Median | |
|------------------------------|---------------|--------------|---------------|--------------|
| | Rs. | US\$ | Rs. | US\$ |
| Direct and indirect costs | | | | |
| Direct medical cost | 3,116 | 42.2 | 2,385 | 32.3 |
| Direct non-medical costs | 2,444 | 33.1 | 2,297 | 31.1 |
| Indirect cost | 8,084 | 110.7 | 5,167 | 70.8 |
| <i>Total household costs</i> | <i>13,659</i> | <i>187.1</i> | <i>12,050</i> | <i>165.1</i> |
| Annual income | | | | |
| Household | 127,074 | 1,720.7 | 122,665 | 1,661.0 |
| Per capita | 23,366 | 316.4 | 22,539 | 305.2 |
| Median costs as a % of | | | | |
| Annual household income | 11% | | | |
| Annual per capita income | 57% | | | |

Exchange rate 1 US\$=74 Rs. (Sept. 2010)

DISCUSSION

During the past five years, considerable efforts were made by public health authorities in Nepal towards the elimination of VL such as the decentralisation and provision of free diagnosis and treatment and the introduction of the oral drug miltefosine in the public health system. In this study we studied the health seeking behaviour and documented household costs and coping strategies for one episode of VL from the perspective of the patient in a miltefosine-based treatment program. From our findings, the following observations can be made.

First, our results showed that the cost of a VL episode to patients and their family was high notwithstanding the free provision of drugs and diagnostics by the government. With a median total cost of US\$ 165 per episode, the economic burden of VL across all households was 11% of annual household income or 57% of median annual per capita income. This cost included

both direct costs (medical and non-medical out-of-pocket expenditures) and indirect costs (productive time losses due to illness). While about half (51%) of the households exceeded the catastrophic threshold of 10% of annual household income, it would be wrong to conclude that the economic consequences of VL illness were not significant for the other households. Because VL is a disease of poverty primarily affecting the poorest income groups (Boelaert et al., the ability to cope with the costs of VL illness are limited. This was evident from the coping strategies households used whereby a majority of households were forced to take a loan to pay for the costs of care and/or use all their savings. However, without the free provision of VL drugs the median cost of an episode of VL would be US\$ 226 and the proportion of households exceeding the catastrophic threshold would increase from 51% to 74% (assuming a drug cost of US\$63 for a 28-days course of miltefosine at WHO preferential prices (Olliaro & Sundar 2009)).

Secondly, direct costs accounted for 47% of total costs and were largely caused by out-of-pocket expenditures households made on ancillary drugs and food. Households that visited private-for-profit health providers incurred substantial expenditures on ancillary drugs. These ancillary drugs, most frequently antibiotics, antipyretics or vitamin injections, were given to patients prior to their referral to a public hospital for VL treatment. From our study we cannot say whether the prescription of these ancillary drugs were justified on medical grounds. Besides the ancillary drugs, another important direct cost component were the high food costs for the patient and accompanying relatives and was caused by the extensive stay at the hospital for treatment. Although miltefosine is an oral drug, patients stayed at the hospital for a median of 10 days.

A number of studies had been carried out prior to the VL elimination initiative (Rijal et al. 2006; Adhikari & Maskay 2005; Adhikari et al. 2009). Our findings seem to suggest that, compared to these studies, the economic burden of VL (as a % of household income) has decreased. In particular the magnitude of indirect costs (as a % of total costs) was less in our study likely due to a shorter patient delay. We cannot say whether the shorter patient delay observed in this study was the result of increased patient knowledge of VL, but the previous studies reported that traditional providers were most commonly the patient's first choice of provider, which may have lengthened the delay until VL diagnosis. However, these conclusions need to be considered as tentative. For instance the total cost of VL in our study was higher compared to Rijal et al. (2006) but much lower compared to Adhikari & Maskay (2005) and Adhikari et al. (2009) while the household income of VL affected households in our study was higher compared to all three studies (to enable comparisons the cost data in these articles were adjusted to the year 2010 using the consumer price index (World Bank 2012)). While

daily wages in Nepal have increased over the years, partly to compensate for the high inflation rate, methodological differences between the studies and in particular the small sample sizes of the first two studies (respectively 18 and 7 households) limit their generalizability beyond the communities or villages where these studies were carried out.

Despite the provision of free VL drugs, we have shown that households still incurred substantial medical out-of-pocket expenditures, especially at private providers. It remains to be seen, however, if these medical costs can be prevented or at least diminished. While prepayment schemes, such as community-based health insurance, may be a solution, coverage in Nepal is very low and expansion of coverage to VL affected households is unlikely in the near future. More realistic and feasible approaches consist of reducing indirect costs and vector control. Treatment duration can be reduced substantially by considering alternative VL drugs to miltefosine as single-dose liposomal amphotericin B or a short course combination therapy (Meheus et al. 2010). Vector control, such as indoor residual spraying, has been shown to be effective in reducing the number of sandflies inside the house (Chowdhury et al. 2011a; Chowdhury et al. 2011b) and may therefore reduce disease transmission. Since 2011, the Government of Nepal has also introduced a conditional cash transfer programme whereby households receive Rs. 1,000 (US\$13.5) upon completion of treatment at a public hospital (Adhikari et al. 2011). This payment can be used to cover transportation costs but was not yet in place when we carried out the household survey. However from our study, the total transportation costs of households were smaller than Rs. 1,000 while food costs were higher. The conditional cash programme therefore ought to be expanded to include food costs as well.

This study had a number of limitations. Because patients were selected from the medical records of the District Public Health Offices and the BPKIHS, only patients treated at public health facilities were included in the study. The records kept by the District Public Health Offices are obtained through passive surveillance from cases detected and treated by public health facilities. Private for-profit providers are not required to report patients treated at their facilities. Due to the low incidence of VL and the high number of private practitioners in Nepal, it would have been difficult and costly to find and interview these patients. Because patients exclusively treated at private-for-profit providers were excluded, we probably underestimated the true burden of VL. Despite this limitation, our findings were still representative for a large proportion of the VL population because in Nepal, contrary to India, a relatively small proportion of patients seek VL treatment from private-for-profit providers (about 11% of patients according to (Mondal et al. 2009). Nonetheless, efforts should be made in the future to include the private sector in the control of VL.

A second limitation is related to the recall bias. With decreasing VL incidence rates in Nepal, we have chosen a recall of 12 months to allow the identification of a sufficient number of households. To minimize the recall bias we only analysed the most recent case of VL in the household. Because of the clustering of VL in communities and villages, several cases of VL often occur in the same household. For instance, in 44% of households in our study one or more members had been treated for VL before, often within a few years. The occurrence of more than one case of VL in the same household would significantly increase the economic impact of VL to households. And even if these cases do not occur in the same year, the risk of impoverishment and indebtedness would still be much higher.


CONCLUSION

With a shorter delay to diagnosis once the patient has presented to a qualified health provider and the free provision of the correct first-line drug, our study indicates that the case management component of the Nepalese VL programme performs rather well. In particular free VL diagnosis and drugs at public health facilities have been an important policy measure in Nepal to lower financial barriers and improve access to VL diagnosis and care. Without this policy the economic burden of VL would have been much higher. However, the economic impact of VL is still considerable and intensified efforts are needed to further reduce the burden of VL to affected households or prevent the transmission of VL. These include shortening the duration of stay at the hospital and expanding demand side financing mechanisms to cover a wider range of costs incurred by households.

REFERENCES

- Adhikari SR, Maskay NM (2005) Economic cost and consequences of kala-azar in Dhanusha and Mahottari districts of Nepal. *Indian J Community Med* 30: 121–25.
- Adhikari SR, Maskay NM, Sharma BP (2009) Paying for hospital-based care of Kala-azar in Nepal: assessing catastrophic, impoverishment and economic consequences. *Health Policy Plan* 24: 129–39.
- Adhikari SR, Prasai DP, Sharma SK (2011) A review of demand side financing schemes in the health sector in Nepal. *Nepal Health Sector Support Programme*. pp.1–44.
- Alvar J, Yactayo S, Bern C (2006) Leishmaniasis and poverty. *Trends Parasitol* 22: 552–57.
- Boelaert M, Meheus F, Sanchez A, et al. (2009) The poorest of the poor: a poverty appraisal of households affected by visceral leishmaniasis in Bihar, India. *Trop Med Int Health* 14: 639–44.
- Chowdhury R, Dotson E, Blackstock AJ, et al. (2011a) Comparison of insecticide-treated nets and indoor residual spraying to control the vector of visceral leishmaniasis in Mymensingh District, Bangladesh. *Am J Trop Med Hyg* 84: 662–67.
- Chowdhury R, Huda MM, Kumar V, et al. (2011b) The Indian and Nepalese programmes of indoor residual spraying for the elimination of visceral leishmaniasis: performance and effectiveness. *Ann Trop Med Parasitol* 105: 31–35.
- Epidemiology and Diseases Control Division (2010) Kala-azar annual report of Nepal. Kathmandu: Ministry of Health and Population, Department of Health Services, Govt. of Nepal. pp. 97–101.
- Joshi DD, Sharma M, Bhandari S (2006) Visceral leishmaniasis in Nepal during 1980–2006. *J Commun Dis* 38: 139–48.
- Meheus F, Balasegaram M, Olliaro P, et al. (2010) Cost-effectiveness analysis of combination therapies for visceral leishmaniasis in the Indian subcontinent. *PLoS Negl Trop Dis* 4: e818.
- Mills A (1993) The household costs of malaria in Nepal. *Trop Med Parasitol* 44: 9–13.
- Mondal D, Singh SP, Kumar N, et al. (2009) Visceral leishmaniasis elimination programme in India, Bangladesh, and Nepal: reshaping the case finding/case management strategy. *PLoS Negl Trop Dis* 3: e355.
- O'Donnell O, van Doorslaer E, Wagstaff A, Lindelow M (2008) Analyzing health equity using household survey data: a guide to techniques and their implementation. Washington D.C.: The World Bank.
- Olliaro P, Sundar S (2009) Anthropometrically derived dosing and drug costing calculations for treating visceral leishmaniasis in Bihar, India. *Trop Med Int Health* 14: 88–92.
- Ranson MK (2002) Reduction of catastrophic health care expenditures by a community-based health insurance scheme in Gujarat, India: current experiences and challenges. *Bull World Health Organ* 80: 613–21.
- Rijal S, Chappuis F, Singh R, et al. (2003) Treatment of visceral leishmaniasis in south-eastern Nepal: decreasing efficacy of sodium stibogluconate and need for a policy to limit further decline. *Trans R Soc Trop Med Hyg* 97: 350–54.

- Rijal S, Koirala S, Van der Stuyft P, Boelaert M (2006) The economic burden of visceral leishmaniasis for households in Nepal. *Trans R Soc Trop Med Hyg* 100: 838–41.
- Singh SP, Hasker E, Picado A, et al. (2010) Risk factors for visceral leishmaniasis in India: further evidence on the role of domestic animals. *Trop Med Int Health* 15 Suppl 2: 29–35.
- Singh SP, Reddy DC, Rai M, Sundar S (2006) Serious underreporting of visceral leishmaniasis through passive case reporting in Bihar, India. *Trop Med Int Health* 11: 899–905.
- Singh VP, Ranjan A, Topno RK, et al. (2010) Estimation of under-reporting of visceral leishmaniasis cases in Bihar, India. *Am J Trop Med Hyg* 82: 9–11.
- World Bank (2012). World Development Indicators. Available: data.worldbank.org/data-catalog/world-development-indicators. Accessed 15 March 2012.
- World Health Organization (2005) Regional strategic framework for elimination of kala-azar from the South-East Asia region (2005–2015). New Delhi: WHO Regional Office for Southeast Asia.
- World Health Organization (2007) Health system in Nepal: challenges and strategic options. New Delhi: WHO Country Office for Nepal.



CHAPTER 6

The economic burden of visceral leishmaniasis in Sudan: an assessment of provider and household costs

Meheus F
Abuzaid AA
Baltussen R
Brima YM
Balasegaram M
Khalil AG
Boelaert M
Musa M

Submitted for publication; 2013.

ABSTRACT

Visceral leishmaniasis (VL) is a neglected parasitic disease that is fatal if left untreated and is endemic in eastern Sudan. We estimated the direct and indirect costs of treatment for VL from the perspective of the provider and the household at three public hospitals in Gedaref State. The median total cost for one VL episode was estimated to be US\$ 450. Despite the free provision of VL drugs at public hospitals, households bore 53% of the total cost of VL with one episode of VL representing 40% of the annual household income. More than 75% of households incurred catastrophic out-of-pocket expenditures. The length of treatment of 30 days led to important costs for both health providers and households. Alternative treatment regimens that reduce the duration of treatment are urgently needed.

INTRODUCTION

Visceral leishmaniasis (VL) is a neglected parasitic disease endemic in many parts of East Africa in particular in eastern Sudan with important foci also found in South Sudan, Ethiopia, Kenya and Uganda. VL is caused by *Leishmania donovani* and transmitted to humans through the bite of a female phlebotomine sandfly. The disease, also known as kala-azar, is systemic and results in the death of the infected individual if treatment is not provided in time. Signs and symptoms of VL include prolonged fever, fatigue and weakness, anaemia and enlarged lymph nodes, spleen and liver (Murray *et al.* 2005)

The public health importance of VL in East Africa is undervalued not least because of the limited knowledge on the disease burden, including the socio-economic aspects of the disease. In the Indian subcontinent (India, Nepal & Bangladesh) studies examining the financial and economic burden of VL have led to a better understanding of the impact and the financial consequences of VL illness at the individual and household level (Meheus *et al.* 2006; Sharma *et al.* 2006; Uranw *et al.* 2013). Other tropical neglected diseases, in particular the helminthic infections, have been successful in garnering international attention and funding by demonstrating the low cost of control interventions and their cost-effectiveness. However, to date, no studies on the costs of VL illness, either from the household or the health-care provider perspective, have been carried out in East Africa. Better cost data would allow the use of economic evaluation as a tool to inform policy decisions and could help to make the case for increased investment and resource allocation in VL control policies and programmes.

Therefore we carried out a costing study on VL in Gedaref State in eastern Sudan. The aim of this study was to estimate both the cost of providing VL diagnosis and treatment services (i.e. health-care provider perspective) as well as the cost to patients and their family to access these services (i.e. household perspective).

MATERIALS AND METHODS

This study is part of a research project on the cost-effectiveness of treatment alternatives for VL in Sudan initiated by the Leishmaniasis East Africa Platform (LEAP) and the Drugs for Neglected Diseases Initiative (DNDi).

Study sites

The study was conducted in Gedaref State, Sudan¹ between October 2009 and May 2011. Gedaref State is located in eastern Sudan bordering Ethiopia and had an estimated population of 1.35 million in 2008, representing 3.4% of the Sudanese population (Central Bureau of Statistics Sudan). Agriculture is the main economic activity in the State with main products including sorghum, sesame, millet and peanuts. Gedaref State is the main VL endemic area in the country and recently experienced an important increase in the number of reported VL cases (deaths) from 2,792 (109) in 2006 to 5,050 (142) in 2010 (Gedaref State Ministry of Health, unpublished data). However, because many people do not have access to health services, the true incidence and mortality of VL are probably much higher Collin *et al.* 2006). It is estimated that about 20,000 VL cases occur annually in Sudan. The number of cases is expected to rise because of migration into endemic areas, climate change, HIV/VL co-infection and the overall lack of efforts to control the disease (Desjeux 1996).

The public health care system in Sudan is organized across three levels: the Federal Ministry of Health (FMOH), the State Ministries of Health (SMOH) and the district level. The delivery of health care also follows a three-tiered structure. Primary health care units are the lowest tier followed by health centres and then rural/community hospitals (WHO Regional Office for the Eastern Mediterranean 2006). The National Malaria, Schistosomiasis and Leishmaniasis Administration (NMSLA) of the FMOH is responsible for VL control activities in Sudan. Currently, VL control activities consist only of passive case detection and treatment at rural/community hospitals. Due to the low level of human and financial resources, VL treatment services are often provided in collaboration with private-not-for-profit partners, either through existing health facilities (e.g. Kassab hospital supported by the Institute of Endemic Diseases (IEND) and DNDi) or by establishing independent health facilities (e.g. Um-el-Kher hospital used to be operated by Médecins sans Frontières (MSF) and now by the Ministry of Health). Although there are no routine VL prevention activities, MSF has been involved in the distribution of bednets in endemic areas since 1995 (Malaria Consortium 2010; Ritmeijer *et al.* 2007).

We purposefully selected 3 health facilities in Gedaref State as study sites: a public rural hospital located in Doka, the State's second largest city; a rural hospital in Kassab and a hospital in Bazora town. The hospital in Doka, funded by the Ministry of Health (MoH), is the largest of the three facilities and has a capacity of 82 beds. It is a referral hospital providing a wide range of medical services including surgery and obstetrical services. Kassab hospital has a capacity of 68 beds and receives funding from both the MoH and DNDi. This hospital, located

between Doka and the State capital Gedaref, mainly provides VL treatment services. Both Doka and Kassab hospital cover the area between Rahad and Atbara rivers together with other rural hospitals in the same area. Bazora hospital is the smallest of the three facilities with a capacity of 51 beds and covers the Rahad river basin area. It is operated by the MoH and also receives funds from the World Health Organization.

Current VL case management in Sudan

Sodium stibogluconate (SSG), a pentavalent antimony compound, is the first-line treatment for VL in Sudan (Federal Ministry of Health Sudan 2004) and is administered intramuscularly on a daily basis for 30 days at a dosage of 20 mg per kg per day. Although widespread resistance to SSG has been reported in the Indian subcontinent (Rijal *et al.* 2003; Sundar *et al.* 2000), the drug is still effective in Sudan with a cure rate of 92% (Hailu *et al.* 2010; Melaku *et al.* 2007; Musa *et al.* 2012) but has numerous side-effects including nausea, diarrhoea, muscle pains, arrhythmia and pancreatitis. Other treatment options are currently being evaluated (Omollo *et al.* 2011). Diagnosis of VL is made with either the rK39 rapid diagnostic dipstick test, the Direct Agglutination Test (DAT) and/or microscopic confirmation of the parasite in lymph node or bone marrow aspirate (Zijlstra *et al.* 1992; Zijlstra *et al.* 2001). Both diagnosis and treatment for VL are provided free of charge at public facilities. Other medical and non-medical costs (registration fee, laboratory investigations, food) need to be paid out-of-pocket by the patient. All patients diagnosed with VL are admitted for the full duration of treatment to ensure 100% adherence and to monitor for side-effects because patients are usually high-risk presenting with severe anaemia and malnutrition (Collin *et al.* 2004).

Costing methodology

The costs of VL were assessed from the perspective of the health-care provider as well as the household. Each perspective involved a different methodology and data collection process described in more detail below (table 1).

¹ Sudan here refers to North Sudan and excludes the Republic of South Sudan that became an independent State in 2011.

buildings was based on the rental price per square meter of office space in the area. Capital costs were annualized using a discount rate of 10% which is the rate mostly used for project appraisals in Africa. A discount rate of 0% (i.e. straight-line depreciation) and standard rates of 3% and 5% (Drummond *et al.* 2005; Gold *et al.* 1996) were used in the sensitivity analysis. We assumed a lifespan of 30 years for buildings, 10 years for equipment and furniture and 5 years for vehicles.

The *step-down costing* was applied as described by Conteh & Walker (2004) using an adapted version of the WHO-CHOICE2 CostIt software package for estimating hospital unit costs. Briefly described, the various departments/units at each health facility were divided into direct and indirect cost centres. Direct costs centres provided services directly to the patient (e.g. outpatient unit or wards), while indirect cost centres provided general services that are necessary to run a hospital, but were not directly related to patient care such as the administration or maintenance department. The final (direct) cost centre of interest to this study was the inpatient ward (all wards were grouped together)³. In a first step we allocated costs directly to the inpatient ward when resource usage could be identified. This was for instance the case with personnel costs whereby medical staff were assigned a proportion of their time (and thus cost) to the inpatient ward based on interviews with the staff. Other costs that could not be assigned directly to the inpatient ward were allocated on the basis of criteria reflecting service use including floor space (for costs of maintenance, cleaning and utilities) and number of staff (for the administration costs).

Once all costs had been assigned to the inpatient ward, a unit cost per inpatient day was calculated by dividing the total cost of the inpatient ward with the total number of inpatient bed-days. Information about the number of inpatient bed-days at each facility was not readily available and was estimated by multiplying the number of admissions with the average length of stay (ALOS). The number of admissions of VL and non-VL patients was retrieved from the statistical records. The ALOS of VL patients was 30 days and obtained from the same sample of 250 medical records described above. The ALOS of non-VL patients was estimated by taking a random sample of 120 medical records at each facility and recording the date of admission and date of discharge⁴. The ALOS for non-VL patients ranged from 2 days at Kassab to 3.5 days at Doka hospital.

² World Health Organization, Choosing Interventions that are Cost Effective (WHO-CHOICE): http://www.who.int/choice/toolkit/cost_it/en/index.html

³ None of the health facilities had disease-specific wards but were differentiated by gender and age (male, female and paediatric ward).

⁴ At Kassab, all medical records of non-VL patients were consulted.

The median total cost of VL care from the provider perspective was obtained by multiplying the unit cost per inpatient day with the ALOS for a VL patient and adding to this figure the medical costs per VL patient obtained with the ingredients approach.

Costs from the household perspective

Costs from the household perspective included direct and indirect costs. These were collected with a hospital exit survey using a pre-tested structured questionnaire. A total of 75 patients were interviewed at Kassab hospital (n=45) and Bazora HC (n=30) between December 2010 and May 2011. At the time of the survey there were no patients attending Doka hospital due to a shortage of SSG at the hospital. The interviews were conducted in Arabic by a medical doctor with extensive experience in VL diagnosis and treatment (one of the authors, AA).

The monetary expenditures by patients and their family to access and receive VL services were recorded separately for all health care providers that were visited. Respondents were asked about all direct medical costs they incurred such as expenditure on registration fees, drugs, laboratory investigations and medical supplies at each provider. The survey also included questions on non-medical costs including the cost of transportation to and from the various health providers as well as food costs incurred while travelling to a health provider and during hospitalization at the treatment facility.

Indirect costs represented the loss of productive time of patients and family members taking care of the patient due to VL illness. Patients and their caretakers were asked about the number of days they were unable to engage in productive activities, and this was multiplied by a median daily income to obtain the indirect cost of a VL episode. Most patients and caretakers that contributed to the household income combined subsistence farming with casual labour during the off-season when there was no field work. To estimate the median daily income (as well as household income in general), the survey collected data on the economic activities of all household members, the number of months these activities were done and the daily or monthly income in the case of casual labour. For subsistence farming, information was collected on the annual production of each produce which was valued with local market prices and divided by the number of agriculturally active household members. Data was also gathered on the income from sales of animals and animal products (e.g. milk) and remittances from family members. Farming activities were done on average 6 months per year, while the rest of the year consisted of casual labour usually in construction (e.g. brick making) or the collection and sale of wood.

The direct and indirect costs were added together to obtain the median total cost of an episode of VL from the perspective of the household. The costs of VL were considered catastrophic if they exceeded 10% of the annual household income (O'Donnell *et al.* 2008; Ranson 2002).

Data analysis

The data was entered in Excel (Microsoft) and analysed with Excel and STATA v10.1 (Stata Corp., College Station Tx, USA). Cost data collected prior to 2010 were inflated using the consumer price index (World Bank 2012) to the year 2010. All cost data are presented in US\$ whereby US\$ 1 = SDP 2.64 (2010).

RESULTS

Costs from the provider perspective

Table 2 shows the median cost of VL case management per patient from the provider perspective as well as activity data for each of the three health facilities. The total cost includes both the “hotel” and the medical component.

Table 2: Activity statistics and unit costs by health facility

| | Kassab hosp. | Doka hosp. | Bazora hosp. |
|-------------------------------------------------|---------------------------------|---------------------------------|-------------------|
| Area covered | Between Rahad and Atbara rivers | Between Rahad and Atbara rivers | Rahad river basin |
| Beds | 68 | 82 | 51 |
| Admissions VL (non-VL) | 805 (95) | 102 (3,049) | 580 (1,198) |
| In-patient days VL (non-VL) | 24,150 (187) | 3,060 (10,641) | 17,400 (3,810) |
| Total in-patient days | 24,337 | 13,701 | 21,210 |
| Average length-of-stay VL (non-VL) ¹ | 30 (2) | 30 (3.5) | 30 (3) |
| Bed occupancy rate | 111% | 46% | 114% |
| Median cost per patient (IQR) (US\$) | 154 (137-186) | 366 (349-399) | 117 (100-147) |

¹ The ALOS was estimated from a random sample of medical records of VL and non-VL patients.

There were marked differences in the median total cost per patient between facilities ranging from US\$117 at Bazora, US\$155 at Kassab and US\$366 at Doka hospital. The medical cost of VL represented 13%, 30% and 38% of total costs at Doka, Kassab and Bazora respectively. The median medical cost was US\$45 (IQR \$28-\$75) per patient with the cost of SSG representing 91% of medical costs. The large variation between facilities in the total cost per patient was caused by the difference in the (hotel) unit cost per inpatient day. While the unit cost per

inpatient day was similar at Bazora and Kassab hospitals (US\$2.5 and US\$3.8 respectively), it was nearly 3 times higher at Doka hospital compared to Kassab (US\$11 per inpatient day).

Varying the discount rate between 0% and 10% did not have much impact on the cost per inpatient day. At Bazora hospital, where capital costs as a proportion of total hospital costs were the largest (23% vs. 12% at Kassab and 11% at Doka), the unit cost per inpatient day changed from US\$ 2.1 (0% discount rate) to US\$ 2.5 (10% discount rate).

Costs from the household perspective

Patient and household characteristics

A total of 75 patients attending Kassab (n=45) and Bazora hospital (n=30) were interviewed; we did not interview any patients at Doka hospital because of a shortage of VL drugs at the time of the study. The cost data from the household perspective collected from Kassab and Bazora hospital were combined in the analysis (tables 3 and 4). Site-specific household cost data is provided in appendix 1.

The majority of patients were males (68.0%) and young with 60% of patients below the age of 15 corresponding to the age profile of other studies showing the epidemic in East Africa is concentrated amount the young. The median age of the sample was 13 (range 3 to 40); female patients were younger than males (median 7.5 (range 2-34) vs. 13 (range 2-40); p<0.05). Overall the level of education was low. Amongst patients aged 6 and above (n=61), 29 (47.5%) had not received any formal education (including Koranic schools). Only 3 patients had completed primary school. There were major differences in the level of education between males and females whereby male patients were more likely to have received some primary education (p<0.01). The median household size was 6 members.

Twenty-nine per cent of patients were economically active (n=22), mainly engaged in subsistence farming (n=13; 59.1%) and daily labour (n=7; 31.9%). Two patients were formally employed. Of those engaged in farming, 7 patients were working as casual labourers during the off-season (defined as secondary occupation). The median annual income of working patients was US\$ 471 (IQR US\$ 244-1,049).

Subsistence farming was the principal economic activity for 80% of households (n=60). A median of two different crops were cultivated by the same household (IQR 2-3; range 1-4); sorghum was the most common crop, followed by sesame, millet and peanuts. Although 40% of households reported owning livestock, either sheep, goats and/or cattle, the headcount was

low and only 47% of these households sold livestock over the past year (n=14) contributing to 6% of their household income. About 19% of households reported receiving remittances, nearly all from outside Sudan (92.9%). The median amount of remittances received was US\$ 378 per year (IQR US\$ 114-454). Including all sources of income, the median annual household income was US\$1,116 (IQR US\$ 744-1,818) and the median annual per capita income was US\$ 208 (IQR US\$ 140-341).

Direct costs at health providers prior to admission

Prior to admission for treatment at Kassab or Bazora hospital, patients had visited on average 3 other health providers (IQR 2-4). A public provider - either a village health worker (43%), a health centre (20%) or a hospital (25%) - was most often the first choice of provider mainly due to their proximity to the patients’ home while the remainder of patients visited either a chemist (3%) or a private general practitioner (9%). Because there are few private (formal) health providers in rural areas, less than 10% of households had initially visited a private provider. In subsequent visits, households consulted a private general practitioner more frequently (24% of households had consulted a private general practitioner after the first provider) or a public hospital until eventually all patients consulted or were referred to either Kassab or Bazora hospital and admitted for treatment.

Households incurred a median total cost of US\$ 33 during the health seeking phase (IQR US\$ 9-73) (table 3) and this included expenses on consultation (US\$ 1), drugs (US\$ 14), laboratory investigations (US\$ 3) and transportation (US\$ 1) (table 4). Households paid the most at private formal providers (US\$ 51), twice as much as at public hospitals (US\$ 24) (p<0.01), of which 94% were direct medical costs for consultation, drugs and diagnosis/laboratory investigations. Households that visited private laboratories were usually referred by a previous provider to be tested for VL.

Direct costs at the treatment facility

The combined direct medical and non-medical costs by households for Kassab and Bazora hospital are presented in table 4. All interviewed patients were treated with sodium stibogluconate; our study confirmed that sodium stibogluconate was provided free of charge to patients in these two facilities. Very few laboratory investigations were done once the patient was admitted. The median direct medical cost was US\$ 14 (IQR US\$ 10-22). Over 85% of costs were non-medical, mainly food costs (median US\$ 112) caused by the long stay at the hospital (median 30 days). The food costs were for the patient, caretaker(s) and other accompanying relatives. All patients were accompanied by at least one adult caretaker, usually the mother,

Table 3: Mean/median cost per provider visited during the health seeking phase prior to admission (US\$ 2010)

| Health provider | n | Mean | (sd) | Median | (IQR) |
|-----------------------|----|------|---------|--------|--------------|
| Traditional healer | 3 | 10.2 | (9.2) | 10.2 | (3.7-16.7) |
| Chemist | 2 | 8.7 | (4.5) | 8.7 | (5.6-11.9) |
| VHW | 37 | 14.6 | (19.8) | 9.3 | (4.5-16.0) |
| Public health centre | 22 | 33.3 | (44.7) | 15.3 | (10.0-42.8) |
| Public hospital | 30 | 56.8 | (73.8) | 24.2 | (12.1-55.8) |
| Private doctor | 26 | 90.7 | (125.4) | 51.2 | (32.7-101.2) |
| Private laboratory | 13 | 27.9 | (16.8) | 26.0 | (17.9-34.2) |
| Total cost per person | 75 | 60.8 | (103.5) | 33.1 | (9.3-73.3) |

Note: All data presented in this table was calculated using data collected at 2 hospitals (Kassab and Bazora). Household costs by facility are provided in appendix table A.1

and one or two young children that stayed with the patient at the hospital for the full duration of treatment. The median direct cost of VL care at the treatment facility was US\$ 148 (IQR US\$ 128-184).

Indirect costs of VL

Sixteen out of 22 patients (73%) reported they were unable to carry out either their primary or secondary occupation because of VL illness which resulted in an income loss. The median number of workdays lost was 51 days (IQR 44-63). The median loss of income for working patients was US\$ 101 (IQR US\$ 62-233) while the average loss of income across all patients was US\$ 41 (sd US\$ 135). Among adult caretakers, 20 reported an income loss (out of 99 caretakers), with a median loss of 39 working days (IQR 31-58) because they had to take care of the patient. The median loss of income to caretakers was US\$ 95 (IQR US\$ 45-179). The average income loss across all attendants, both working and non-working was US\$ 44 (sd US\$ 85).

Total costs of a VL episode

The median total cost for one VL episode was US\$ 450 (IQR US\$ 387-544) (table 5). The median cost of VL case management across the three health facilities was US\$ 211 (IQR US\$ 197-244). The median direct expenditure by households, including the health seeking phase and the costs incurred at the treatment facility, was US\$ 185 (IQR US\$ 158-240) while the median income loss (i.e. indirect costs) was US\$ 22 (IQR US\$ 0-113.9). Overall, households bore 53% of the total cost of VL and public health facilities 47%. Direct costs, in particular non-medical costs, were the main cost driver from the perspective of the household representing nearly 86% of the

Table 4: Mean/median direct costs of VL care during the health seeking and treatment phase (US\$ 2010)

| | Health seeking phase | | | Admission & treatment | | |
|---------------------------------------|----------------------|--------|------------|-----------------------|--------|---------------|
| | Mean (sd) | Median | (IQR) | Mean (sd) | Median | (IQR) |
| Direct medical costs | | | | | | |
| Consultation | 6.6 (13.0) | 1.1 | (0.0-6.0) | 3.3 (1.7) | 3.7 | (1.1-4.8) |
| Ancillary drugs | 26.0 (47.7) | 13.8 | (6.7-37.2) | 11.1 (10.0) | 7.4 | (4.3-15.6) |
| Laboratory investigations | 13.2 (24.9) | 3.3 | (0.0-16.7) | 4.1 (7.0) | 1.9 | (0.2-4.1) |
| <i>Total direct medical costs</i> | 45.9 (78.4) | 24.2 | (9.3-54.0) | 18.5 (14.9) | 14.0 | (9.9-22.3) |
| Direct non-medical costs | | | | | | |
| Transportation | 7.5 (14.7) | 1.1 | (0.0-7.4) | 6.5 (10.1) | 2.6 | (0.7-8.9) |
| Food | 6.5 (19.9) | 0.0 | (0.0-4.8) | 121.5 (44.5) | 111.6 | (111.6-141.4) |
| Other | 1.0 (4.4) | 0.0 | (0.0-0.2) | 5.0 (4.1) | 4.5 | (2.0-6.9) |
| <i>Total direct non-medical costs</i> | 15.0 (31.1) | 2.6 | (0.0-18.6) | 133 (46.0) | 126.5 | (113.9-157.8) |
| Total direct costs | 60.8 (103.5) | 33.1 | (9.3-73.3) | 151.5 (49.1) | 148.5 | (128.2-184.2) |

Note: All data presented in this table was calculated using data collected at 2 hospitals (Kassab and Bazora). Household costs by facility are provided in appendix tables A.2 and A.3

median household cost while indirect costs represented 14% of this cost. The median annual household income was estimated to be US\$ 1,116. The economic burden of VL to households, including direct and indirect costs, was equal to 23% of the median annual household income.

Table 5: Summary of direct and indirect costs for one VL episode (US\$ 2010)

| | Mean | (sd) | Median | (IQR) |
|--------------------------|-------|---------|--------|---------------|
| Direct costs | | | | |
| Household ¹ | 212.4 | (122.8) | 185.1 | (158.5-240.2) |
| Provider ² | 220.3 | (31.4) | 211.1 | (197.3-243.9) |
| Indirect costs | 118.8 | (181.9) | 22.3 | (0-113.9) |
| Total cost | | | | |
| Household | 297.1 | (250.4) | 238.4 | (171.8-333.2) |
| Household + provider | 508.2 | (250.3) | 450.0 | (386.9-544.3) |
| Median costs as a % of | | | | |
| Annual household income | 23% | | | |
| Annual per capita income | 122% | | | |

¹ Health seeking and treatment facility costs; Calculated using the mean/median data from 2 hospitals (Kassab and Bazora) since no patients attended Doka hospital at the time of the study
² Calculated using the average hotel unit cost of US\$ 5.74 across the three hospitals

DISCUSSION

This is the first study to provide a comprehensive set of estimates on the cost and economic burden of VL in Sudan and East Africa in general. We collected data from both the provider and the household perspective. Data on provider costs were collected from three health facilities in Gedaref State; direct costs to access treatment and indirect costs of VL illness from the perspective of the household were collected using a structured exit questionnaire with 75 patients in two of these facilities.

The median total cost per episode of VL was estimated to be US\$ 450. Over 75% of households incurred catastrophic out-of-pocket expenditures (defined as expenditures exceeding 10% of annual median income) when considering only direct costs while 83% of households exceeded this threshold when also including indirect costs. These findings concur with studies that were carried out in the Indian subcontinent (Nepal, India and Bangladesh) which also found a huge economic burden of VL illness for households (Meheus *et al.* 2006; Rijal *et al.* 2006; Sarnoff *et al.* 2010; Sundar *et al.* 2010; Uranw *et al.* 2013). There were however some differences with regard to the distribution of costs. Indirect cost as a proportion of the total household cost

was smaller in Sudan, mainly linked with a different age profile. In our study we found that 60% of patients were below the age of 15 while for instance in Bangladesh it was 47%³. These differences are consistent with the demographic profile of VL across countries reported by Harhay *et al.* (2011). Because patients were younger in Sudan, fewer reported earning an income. For instance we found that 29% of patients were economically active and generating an income to the household compared to 36% in Nepal (Uranw *et al.* 2013) and 42% in India (Meheus *et al.* 2006).

In terms of the provider costs, the cost of hospitalization (which did not include the medical costs of VL) was the main cost driver and varied between 62% and 87% of the total provider cost at Bazora and Doka hospital respectively. We observed large differences in the unit cost per inpatient day across the three facilities. There are several factors that explain the higher unit cost per inpatient day at Doka hospital. From the cost side, because Doka hospital is a secondary level referral facility, it has more skilled health staff resulting in higher personnel costs. For instance the ratio of medical doctors working at Doka compared to Bazora hospital was 5:1. The capital costs were also higher compared to the other two health facilities. But the factor that influenced the unit cost most was the workload at each hospital, and in particular the number of VL patients. Because most costs are fixed (including personnel costs on the short and medium term) and do not vary with the number of inpatients at the hospital, the fewer the inpatients, the higher the unit cost per inpatient day. A total of 1,487 patients were admitted for VL treatment in the three study facilities in 2008, with important differences in case load across facilities. Overall the bed occupancy⁵ rate at Doka hospital was low (46%) while in Kassab and Bazora it exceeded 100% (on average, there were not sufficient beds to accommodate all patients). Doka hospital admitted the least VL patients in 2008 due to frequent shortages of SSG. These patients were then referred to Kassab hospital, located approximately 50kms away. The number of VL patients at Bazora hospital was high because the hospital covers a very large area (the Rahad River Basin) with important transmission of VL.

Our estimates were also lower than the WHO-CHOICE hotel unit cost estimates for Sudan (i.e. US\$ 13.5 inflation-adjusted for a secondary-level hospital at 80% occupancy rate) (WHO-CHOICE 2012). The variation in unit costs between facilities in this study was caused by differences in the cost structure of the health facilities, but especially by the patient load since we observed a low occupancy rate at Doka hospital (46%) and rates exceeding full occupancy at Kassab and Bazora. During the time of the study, most VL patients who attended Doka hospital were referred to Kassab hospital because of a shortage in SSG drugs. Given these

logistical problems the unit costs of hospitalization should be interpreted with care and not considered as an indicator of efficient resource usage. As Hansen *et al.* (200) noted in a study on the hospital costs of HIV/AIDS care, low unit costs may also indicate insufficient resources at these health facilities. Although an evaluation of optimal resource usage for VL care and other services was not within the scope of this study, there were clear indications that for example the hospital in Bazora was under-staffed compared to its patient load.

A limitation to our study was that we did not capture the seasonal pattern of costs. While this is less relevant for provider costs that were based on data for a complete financial year, the household survey was carried out half-way the dry season, which takes place approximately from November to April. This may have had an impact on the health seeking behaviour of households and transportation costs. Since health facilities are more accessible during the dry season, patients are likely to present faster to a health facility and transportation costs will be lower. For example, Gerstl *et al.* (2006) in a study examining the accessibility of VL treatment centres in Gedaref State found that transportation costs were 2 to 3 times higher during the rainy season. In addition the time of the survey may also have had an impact on the opportunity cost of time lost due to VL illness and treatment. Patients involved in subsistence farming reported they were not working at the time of the survey because it was the off-season. As such we may have underestimated the indirect cost of VL illness for those patients that did not have a secondary occupation.

Secondly, data on costs were based on actual consumption of resources with the major cost elements (medical costs and personnel) obtained through micro-costing. However the hospital records on activity statistics were poor and the unit costs per inpatient day are particularly sensitive to inaccurate activity data. Information on the number of admissions for both VL and non-VL patients were available at each facility, but the average length of stay needed to be extrapolated from a random sample of medical records. While the average length of stay for VL patients is no doubt close to 30 days because of the duration of the drug regimen, there was more uncertainty regarding the ALOS of non-VL patients.

Finally, we were not able to obtain data for all cost categories at each hospital included in the study. From the provider perspective, detailed information from medical records to estimate the direct medical costs was only available at Kassab hospital. Data from the household perspective was collected from Kassab and Bazora hospital and not Doka hospital since there were no patients attending the latter facility at the time of the study due to a shortage of SSG. Given the limitations in data availability and since the aim of the study was to estimate the cost of

⁵ The bed occupancy rate was calculated as the number of inpatient days in a year divided by the number of beds x 365.

providing and accessing VL diagnosis and treatment services, rather than making comparisons across facilities (in terms of e.g. efficiency or access), results were presented in terms of the mean/median across the three facilities. This is the first study of its kind for VL carried out in Sudan (and eastern Africa in general). We hope that in the future, cost data on VL diagnosis and treatment will be more readily available to support on-going control efforts in the country.

In the absence of a vaccine and preventive interventions, the diagnosis and treatment of VL is currently the main control strategy available in Sudan. We showed VL diagnosis and treatment to be expensive not only to the public health sector but also to the households compared to their income. Policies are therefore needed that reduce this cost to both providers and households. A common determinant of costs for both perspectives was the length of treatment. The current first line treatment is SSG and patients are admitted for the full duration of treatment of 30 days to ensure adherence to treatment and monitor for possible side-effects. In recent years there have been a number of therapeutic innovations that open the door to treatment alternatives with shorter duration. These treatment alternatives would not only reduce the cost of hospitalization from the provider perspective but also the substantial food costs patients and caretakers incur. One of the alternatives is a combination of SSG and paromomycin for 17 days (Melaku *et al.* 2007) that showed an acceptable safety and efficacy profile in a recent randomized-controlled trial (Musa *et al.* 2012). This treatment is in fact now being rolled out in Gedaref state. Other alternatives include combinations of an oral drug miltefosine with either SSG or liposomal amphotericin B (Omollo *et al.* 2011) that may reduce treatment to 10 days (for the latter combination) or a multi-dose regimen of liposomal Amphotericin B for 10 days (Edwards *et al.* 2011). Liposomal Amphotericin B (AmBisome®) is already registered in Sudan and used as second line treatment but is not considered as first-line because of its very high cost despite preferential pricing agreements between the WHO and its manufacturer Gilead Sciences (US\$ 270 for total dose of 21 mg/kg for a 35kg patient). However a donation by Gilead to treat 50,000 patients opens new perspectives for using AmBisome® as first line treatment in East Africa (Burki 2012).

The cost data that we presented here will facilitate the use of cost-effectiveness analysis to compare the various treatment alternatives presently available or in the near future for Sudan. This study highlighted the importance of including the household perspective in such analysis since VL caused catastrophic health expenditure in almost all households despite drugs and diagnostics being free. In the absence of effective prevention, the best way to alleviate this burden is through a treatment regimen with shorter duration. The article is hopefully the first in a series of studies investigating the behavioural and socio-economic aspects of VL and its control. Until we are able to close this knowledge gap, there will be little interest from the

government and the international community to actively engage in the control of VL in Sudan and East Africa in general.

ACKNOWLEDGMENTS

The authors are grateful to the administrative and health staff at the hospitals of Doka, Kassab and Bazora for their much appreciated cooperation. This study received financial support from the Drugs for Neglected Diseases Initiative and the European Commission under the Health Cooperation Work Programme of the 7th Framework Programme (FP7) to the NIDIAG consortium (www.nidiag.org). Filip Meheus was supported by the ITM VL Fund and the SOFI-A programme funded by the Flemish Government.

REFERENCES

- Bureau of Statistics Sudan. *Fifth population and housing census 2008*. Available at: <http://www.cbs.gov.sd/en/node/6>. Accessed on: July 15, 2012
- Burki T (2012) Drug donated for treatment of visceral leishmaniasis. *Lancet Infect Dis* 12: 106–107.
- Collin S, Davidson R, Ritmeijer K, et al. (2004). Conflict and kala-azar: determinants of adverse outcomes of kala-azar among patients in southern Sudan. *Clin Infect Dis* 38: 612–19.
- Collin SM, Coleman PG, Ritmeijer K, Davidson RN (2006) Unseen Kala-azar deaths in south Sudan (1999–2002). *Trop Med Int Health* 11: 509–12.
- Conteh L, Walker D (2004) Cost and unit cost calculations using step-down accounting. *Health Policy Plan* 19: 127–35.
- Creese, A, Parker, D (1994) *Cost analysis in primary health care: a training manual for programme managers*. Geneva: World Health Organization.
- Desjeux P (1996) Leishmaniasis. Public health aspects and control. *Clin Dermatol* 14: 417–23.
- Drummond M, Sculpher MJ, Torrance GW, O'Brien BJ, Stoddart GL (2005) *Methods for the economic evaluation of health care programmes*. 3rd ed. Oxford: Oxford University Press.
- Edwards T, Omollo R, Khalil EA, et al. (2011) Single-dose liposomal amphotericin B (AmBisome(R)) for the treatment of visceral leishmaniasis in East Africa: study protocol for a randomized controlled trial. *Trials* 12: 66.
- Federal Ministry of Health Sudan (2004) *Manual for diagnosis and treatment of leishmaniasis*. Federal Ministry of Health: Leishmaniasis Administration.
- Gerstl S, Amsalu R, Ritmeijer K (2006). Accessibility of diagnostic and treatment centres for visceral leishmaniasis in Gedaref State, northern Sudan. *Trop Med Int Health* 11: 167–75.
- Gold MR, Russell LB, Siegel JE, Weinstein MC (1996) *Cost-effectiveness in health and medicine*. New York: Oxford University Press.
- Hailu A, Musa A, Wasunna M, et al. (2010) Geographical variation in the response of visceral leishmaniasis to paromomycin in East Africa: a multicentre, open-label, randomized trial. *PLoS Negl Trop Dis* 4: e709.
- Hansen K, Chapman G, Chitsike I, Kasilo O, Mwaluko G (2000) The costs of HIV/AIDS care at government hospitals in Zimbabwe. *Health Policy Plan* 15: 432–40.
- Harhay MO, Olliaro PL, Vaillant M, et al. (2011) Who is a typical patient with visceral leishmaniasis? Characterizing the demographic and nutritional profile of patients in Brazil, East Africa, and South Asia. *Am J Trop Med Hyg* 84: 543–50.
- Malaria Consortium (2010) *Leishmaniasis control in eastern Africa: Past and present efforts and future needs*.
- McPake B, Hongoro C, Russo G (2011) Two-tier charging in Maputo Central Hospital: costs, revenues and effects on equity of access to hospital services. *BMC Health Serv Res* 11: 143.
- Meheus F, Boelaert M, Baltussen R, Sundar S (2006) Costs of patient management of visceral leishmaniasis in Muzaffarpur, Bihar, India. *Trop Med Int Health* 11: 1715–24.
- Melaku Y, Collin SM, Keus K, Gatluak F, Ritmeijer K, Davidson RN (2007) Treatment of kala-azar in southern Sudan using a 17-day regimen of sodium stibogluconate combined with paromomycin: a retrospective comparison with 30-day sodium stibogluconate monotherapy. *Am J Trop Med Hyg* 77: 89–94.
- Murray HW, Berman JD, Davies CR, Saravia NG (2005) Advances in leishmaniasis. *Lancet* 366: 1561–77.
- Musa A, Khalil E, Hailu A, et al. (2012) Sodium Stibogluconate (SSG) & Paromomycin combination compared to SSG for visceral leishmaniasis in East Africa: a randomised controlled trial. *PLoS Negl Trop Dis* 6: e1674.
- O'Donnell O, van Doorslaer E, Wagstaf A, Lindelow M (2008) *Analyzing health equity using household survey data: a guide to techniques and their implementation*. Washington D.C.: The World Bank.
- Omollo R, Alexander N, Edwards T, et al. (2011) Safety and efficacy of miltefosine alone and in combination with sodium stibogluconate and liposomal amphotericin B for the treatment of primary visceral leishmaniasis in East Africa: study protocol for a randomized controlled trial. *Trials* 12: 166.
- Ranson MK (2002) Reduction of catastrophic health care expenditures by a community-based health insurance scheme in Gujarat, India: current experiences and challenges. *Bull World Health Organ* 80: 613–21.
- Rijal S, Chappuis F, Singh R, et al. (2003) Treatment of visceral leishmaniasis in south-eastern Nepal: decreasing efficacy of sodium stibogluconate and need for a policy to limit further decline. *Trans R Soc Trop Med Hyg* 97: 350–54.
- Rijal S, Koirala S, Van der Stuyft P, Boelaert M (2006) The economic burden of visceral leishmaniasis for households in Nepal. *Trans R Soc Trop Med Hyg* 100: 838–41.
- Ritmeijer K, Davies C, van Zorge R, et al. (2007) Evaluation of a mass distribution programme for fine-mesh impregnated bednets against visceral leishmaniasis in eastern Sudan. *Trop Med Int Health* 12: 404–14.
- Sarnoff R, Desai J, Desjeux P, et al. (2010) The economic impact of visceral leishmaniasis on rural households in one endemic district of Bihar, India. *Trop Med Int Health* 15(S2): 42–49.
- Sharma AD, Bern C, Varghese B, et al. (2006) The economic impact of visceral leishmaniasis on households in Bangladesh. *Trop Med Int Health* 11: 757–64.
- Sundar S, Arora R, Singh SP, Boelaert M, Varghese B (2010) Household cost-of-illness of visceral leishmaniasis in Bihar, India. *Trop Med Int Health* 15(S2): 50–54.
- Sundar S, More DK, Singh MK, et al. (2000) Failure of pentavalent antimony in visceral leishmaniasis in India: report from the center of the Indian epidemic. *Clin Infect Dis* 31: 1104–07.
- Uranw S, Meheus F, Baltussen R, Rijal S, Boelaert M (2013) The economic burden of visceral leishmaniasis on households in eastern Nepal. *PLoS Neglected Tropical Diseases* 7: e2062
- Van der Plaetse B, Hlatiwayo G, Van EL, Meessen B, Criel B (2005) Costs and revenue of health care in a rural Zimbabwean district. *Health Policy Plan* 20: 243–51.
- WHO Regional Office for the Eastern Mediterranean (2006) *Health system profile Sudan*. Egypt: Cairo.
- WHO-CHOICE. *Country specific unit costs*. Available at: http://www.who.int/choice/country/country_specific/en/index.html. Accessed on: July 15, 2012

WHO (2010) *Control of the leishmaniasis: report of a meeting of the WHO expert committee on the control of leishmaniasis, Geneva, 22-26 March 2010*. Geneva: World Health Organization.

World Bank. *World databank: World development indicators & Global development finance*. Available at: <http://data.worldbank.org/indicator/FP.CPI.TOTL>. Accessed on: July 15, 2012

Zijlstra EE, Ali MS, el-Hassan AM, et al. (1992) Kala-azar: a comparative study of parasitological methods and the direct agglutination test in diagnosis. *Trans R Soc Trop Med Hyg* 86: 505–07.

Zijlstra EE, el-Hassan AM (2001) Leishmaniasis in Sudan. Visceral leishmaniasis. *Trans R Soc Trop Med Hyg* 95(S1): S27–S58.

Table A.1: Mean/median cost per provider visited during the health-seeking phase prior to admission (US\$ 2010)

| | Bazora | | | | | | Kassab | | | | | | Total | | | | | |
|-----------------------|--------|------|------|--------|------|-------|--------|------|-------|--------|------|------|-------|------|-------|--------|------|-------|
| | n | mean | sd | median | p25 | p75 | n | mean | sd | median | p25 | p75 | n | mean | sd | median | p25 | p75 |
| Traditional healer | 0 | | | | | | 3 | 10.2 | 9.2 | 10.2 | 3.7 | 16.7 | 3 | 10.2 | 9.2 | 10.2 | 3.7 | 16.7 |
| Chemist | 1 | 5.6 | | 5.6 | 5.6 | 5.6 | 1 | 11.9 | | 11.9 | 11.9 | 11.9 | 2 | 8.7 | 4.5 | 8.7 | 5.6 | 11.9 |
| VHW | 16 | 20.2 | 28.9 | 11.2 | 7.8 | 20.8 | 21 | 10.4 | 6.7 | 9.1 | 4.3 | 14.0 | 37 | 14.6 | 19.8 | 9.3 | 4.5 | 16.0 |
| Public health centre | 6 | 54.2 | 73.2 | 27.3 | 12.3 | 47.3 | 16 | 24.4 | 23.9 | 12.8 | 9.3 | 38.3 | 22 | 33.3 | 44.7 | 15.3 | 10.0 | 42.8 |
| Public hospital | 8 | 36.0 | 50.5 | 13.9 | 10.1 | 40.9 | 22 | 65.5 | 81.3 | 29.0 | 13.8 | 99.5 | 30 | 56.8 | 73.8 | 24.2 | 12.1 | 55.8 |
| Private doctor | 4 | 73.7 | 59.3 | 93.0 | 7.1 | 120.9 | 22 | 94.6 | 137.7 | 47.3 | 35.7 | 89.7 | 26 | 90.7 | 125.4 | 51.2 | 32.7 | 101.2 |
| Private laboratory | 0 | | | | | | 13 | 27.9 | 16.8 | 26.0 | 17.9 | 34.2 | 13 | 27.9 | 16.8 | 26.0 | 17.9 | 34.2 |
| Total cost per person | 30 | 38.1 | 54.7 | 15.9 | 4.5 | 40.9 | 45 | 76.0 | 124.3 | 43.2 | 21.2 | 80.7 | 75 | 60.8 | 103.5 | 33.1 | 9.3 | 73.3 |

Table A.2: Mean/median direct costs of VL care during the health seeking phase (US\$ 2010)

| | Bazora | | | | Kassab | | | | Total | | | |
|--------------------------------|--------|------|--------|-----|--------|------|-------|--------|-------|------|------|-------|
| | mean | sd | median | p25 | p75 | mean | sd | median | p25 | p75 | mean | sd |
| Direct medical costs | | | | | | | | | | | | |
| Consultation | 4.2 | 11.3 | 0.0 | 0.0 | 2.2 | 8.2 | 13.9 | 2.2 | 0.0 | 7.4 | 6.6 | 13.0 |
| Ancillary drugs | 16.9 | 17.5 | 11.2 | 3.7 | 26.0 | 32.1 | 59.4 | 18.2 | 8.2 | 38.0 | 26.0 | 47.7 |
| Laboratory investigations | 5.2 | 9.2 | 0.6 | 0.0 | 8.6 | 18.6 | 30.2 | 7.4 | 1.1 | 24.2 | 13.2 | 24.9 |
| Total direct medical costs | 26.3 | 31.2 | 13.4 | 4.5 | 40.9 | 58.9 | 96.2 | 35.7 | 21.2 | 65.7 | 45.9 | 78.4 |
| Direct non-medical costs | | | | | | | | | | | | |
| Transportation | 7.6 | 19.3 | 0.0 | 0.0 | 4.5 | 7.3 | 10.9 | 2.2 | 0.0 | 10.4 | 7.5 | 14.7 |
| Food | 3.1 | 7.6 | 0.0 | 0.0 | 0.0 | 8.7 | 24.8 | 0.9 | 0.0 | 5.2 | 6.5 | 19.9 |
| Other | 1.1 | 5.2 | 0.0 | 0.0 | 0.2 | 1.0 | 3.8 | 0.0 | 0.0 | 0.2 | 1.0 | 4.4 |
| Total direct non-medical costs | 11.9 | 26.9 | 0.0 | 0.0 | 5.6 | 17.1 | 33.7 | 4.7 | 0.0 | 21.6 | 15.0 | 31.1 |
| Total direct costs | 38.1 | 54.7 | 15.9 | 4.5 | 40.9 | 76.0 | 124.3 | 43.2 | 21.2 | 80.7 | 60.8 | 103.5 |

Table A.3: Mean/median direct costs of VL care at the treatment facility (US\$ 2010)

| | Bazora | | | | Kassab | | | | Total | | | |
|--------------------------------|--------|------|--------|-------|--------|-------|------|--------|-------|-------|-------|------|
| | mean | sd | median | p25 | p75 | mean | sd | median | p25 | p75 | mean | sd |
| Direct medical costs | | | | | | | | | | | | |
| Consultation | 1.7 | 1.3 | 1.1 | 1.1 | 2.2 | 4.5 | 0.9 | 4.8 | 4.8 | 4.8 | 3.3 | 1.7 |
| Ancillary drugs | 18.0 | 11.5 | 16.4 | 11.2 | 22.3 | 6.5 | 5.1 | 5.6 | 2.6 | 7.4 | 11.1 | 10.0 |
| Laboratory investigations | 8.3 | 9.6 | 4.8 | 3.3 | 9.7 | 1.3 | 1.5 | 0.7 | 0.0 | 1.7 | 4.1 | 7.0 |
| Total direct medical costs | 28.0 | 18.9 | 23.3 | 18.6 | 29.0 | 12.3 | 6.0 | 11.5 | 7.8 | 14.0 | 18.5 | 14.9 |
| Direct non-medical costs | | | | | | | | | | | | |
| Transportation | 7.4 | 11.3 | 3.7 | 0.7 | 7.4 | 5.9 | 9.3 | 1.9 | 0.7 | 8.9 | 6.5 | 10.1 |
| Food | 116.0 | 50.3 | 111.6 | 111.6 | 141.4 | 125.2 | 40.3 | 111.6 | 111.6 | 137.7 | 121.5 | 44.5 |
| Other | 5.3 | 4.2 | 4.9 | 3.5 | 6.1 | 4.8 | 4.1 | 3.7 | 1.9 | 6.9 | 5.0 | 4.1 |
| Total direct non-medical costs | 128.7 | 50.5 | 121.1 | 112.4 | 155.5 | 135.8 | 43.0 | 128.0 | 114.4 | 158.9 | 133.0 | 46.0 |
| Total direct costs | 156.6 | 56.6 | 149.4 | 136.2 | 184.2 | 148.1 | 43.8 | 148.5 | 124.8 | 172.5 | 151.5 | 49.1 |



PART III

**The cost-effectiveness of
treatment strategies for visceral
leishmaniasis**



CHAPTER 7

Combination therapy for visceral leishmaniasis

Van Griensven J
Balasegaram M
Meheus F
Alvar J
Lynen L
Boelaert M

SUMMARY

Combination therapy for the treatment of visceral leishmaniasis has increasingly been advocated as a way to increase treatment efficacy and tolerance, reduce treatment duration and cost, and limit the emergence of drug resistance. We reviewed the evidence and potential for combination therapy, and the criteria for the choice of drugs in such regimens. The first phase 2 results of combination regimens are promising, and have identified effective and safe regimens as short as 8 days. Several phase 3 trials are underway or planned in the Indian subcontinent and east Africa. The limited data available suggest that combination therapy is more cost-effective and reduces indirect costs for patients. Additional advantages are reduced treatment duration (8–17 days), with potentially better patient compliance and lesser burden on the health system. Only limited data are available on how best to prevent acquired resistance. Patients who are co-infected with visceral leishmaniasis and HIV could be a reservoir for development and spread of drug-resistant strains, calling for special precautions. The identification of a short, cheap, well-tolerated combination regimen that can be given in ambulatory care and needs minimal clinical monitoring will most likely have important public health implications. Effective monitoring systems and close regulations and policy will be needed to ensure effective implementation. Whether combination therapy could indeed help delay resistance, and how this is best achieved, will only be known in the long term.

INTRODUCTION

Visceral leishmaniasis, also known as kala-azar, is a disseminated protozoan infection caused by the *Leishmania donovani* complex and transmitted via phlebotomine sandflies (Chappuis *et al.* 2007). The zoonotic form, for which dogs are the main reservoir, is present in the Mediterranean basin, China, the Middle East, and South America, and is caused by *Leishmania infantum* or *Leishmania chagasi*. The anthroponotic form (human reservoir) is caused by *L. donovani* and is prevalent in east Africa and the Indian subcontinent (Chappuis *et al.* 2007; Herwaldt 1999). Although the disease is endemic in more than 60 countries, with 200 million people at risk, 90% of the 500 000 cases every year happen in five countries: India, Bangladesh, Nepal, Sudan, and Brazil (figure 1) (Chappuis *et al.* 2007; Desjeux 2004a; Guerin *et al.* 2002; Desjeux 2004b).




Figure 1: Geographical distribution of visceral leishmaniasis

Anthroponotic visceral leishmaniasis happens mainly in east Africa and the Indian subcontinent. The zoonotic form is prevalent in South America, the Mediterranean basin, China, and the Middle East. More than 90% of cases happen in five countries: India, Nepal, Bangladesh, Sudan, and Brazil. Reprinted from Desjeux (2004b) with permission from Macmillan Publishers Ltd.

For most of the past 70 years, the therapeutic armoury for treatment of visceral leishmaniasis has been extremely limited (Alvar *et al.* 2006). Pentavalent antimonials were introduced in the 1940s, and include sodium stibogluconate and meglumine antimoniate. Use of amphotericin

B followed after a few decades, and was later joined by paromomycin, a cheap and effective parenteral drug with an acceptable toxicity profile that can easily be given by intramuscular injection (Sundar *et al.* 2008). The development of miltefosine, the only drug at present that can be given orally for visceral-leishmaniasis treatment, has been a major breakthrough (Berman 2006; Croft *et al.* 2006). This drug is the mainstay of the recently launched visceral-leishmaniasis elimination plan in the Indian subcontinent, (WHO 2004) and benefits from a preferential pricing scheme that puts it at the same price as generic pentavalent antimonials if large quantities are purchased. Finally, different lipid formulations of amphotericin B (ie, liposomal amphotericin B) have been developed, which are similar to amphotericin B in efficacy but with fewer toxic effects (table 1) (Bern *et al.* 2006). Although these formulations were initially prohibitively expensive, the preferential price now offered to governments of endemic countries, WHO, and non-governmental organisations make them an option for low-income and middle-income countries. Although other compounds are being developed, these drugs are likely to constitute the main therapeutic options for visceral leishmaniasis in the years to come (Alvar *et al.* 2006; Maltezou 2008). These drugs belong to chemically unrelated classes and are thought to have distinct targets. All of them have several important disadvantages (table 1).

There are several reasons why consensus has grown over the past few years towards the use of combination regimens in visceral leishmaniasis (Alvar *et al.* 2006; Bryceson 2001; Croft *et al.* 2006; Den Boer *et al.* 2006; Sundar *et al.* 2007; Singh *et al.* 2006). First, combining drugs from different chemical classes could reduce treatment duration or total drug doses, resulting in fewer toxic effects, higher compliance, and less burden on the health system. This could also reduce the overall costs (direct and indirect) and provide a more cost-effective option. Increasing reports of treatment failure with pentavalent antimonials from the Indian subcontinent have raised the issue of acquired drug resistance (Croft *et al.* 2006; Lira *et al.* 1999; Sundar 2001). This concern now extends to miltefosine, because of its long half-life and susceptibility to develop resistance with a single point mutation (Sundar *et al.* 2005; Perez-Victoria *et al.* 2006; Seifert *et al.* 2007. Combination therapy might help to delay the emergence of resistance and increase the therapeutic lifespan of the respective drugs, as has been seen for diseases like malaria, tuberculosis, and HIV (Bryceson 2001). Finally, combination therapy could improve treatment efficacy for complicated cases, such as patients coinfectd with HIV, for whom treatment outcomes with monotherapy have been consistently poor (Alvar *et al.* 2008). We review the evidence and explore the potential of combination therapy for visceral leishmaniasis in areas of anthroponotic transmission—in particular, the Indian subcontinent and east Africa. Given the anthroponotic pattern, these areas have the highest threat of drug resistance and bear the highest burden of visceral leishmaniasis. We will discuss the evidence and the criteria for a rational design of such combination regimens that focus on the parallel or

Table 1: Drugs currently used for treatment of visceral leishmaniasis

| | Marketing* (trade name of drug) | Regimen | Clinical efficacy | Resistance | Toxicity | Cost of drug course (US\$) | Disadvantages |
|------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------|-----------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------|--------------------------------------------------------------------------------------------|
| Pentavalent Antimonials (sodium stibogluconate, meglumine antimoniate) | Sodium stibogluconate: Albert David (SSG), GlaxoSmithKline (Pentostam). Meglumine antimoniate: Sanofi Aventis (Glucontime) | 20 mg antimony per kg bodyweight for 20-30 days (depending on geographic area), intravenous or intramuscular | 35-95% (depending on geographic area) | Treatment failure up to 60% (Bihar, India) | Moderately toxic: cardiac effects, pancreatitis, nephrotoxicity hepatotoxicity | 53 (generic) to 198 (branded SSG) | Quality control; length of treatment; painful injection; resistance in India |
| Amphotericin B | Bristol Meyers Squibb (Fungizone); generic companies | 1 mg/kg every other day for up to 30 days (15 mg/kg total dose), intravenous | > 97% all regions | Not documented | Moderately toxic: nephrotoxicity (inpatient care needed) | About 21 (generic) | Need for slow intravenous infusion; dose-limiting nephrotoxicity; heat stability |
| Liposomal amphotericin B | Gilead (AmBisome) | 5-20 mg/kg total dose in 4-10 doses over 10-20 days, intravenous | Asia: > 97%; India, single dose: 91%; Africa: not fully established | Not documented | Nephrotoxicity (limited) | 280 (preferential) for 20mg/kg dose; about 3000 (non-preferential) | Price; need for slow intravenous infusion; heat stability (needs to be stored below 25°C) |
| Miltefosine | Paladin, Montreal, Canada (Impavido) | 2-2.5 mg/kg daily over 28 days (India only), oral | Asia: 94% (India); Africa: 60% (single field study), 93% in patients not infected with HIV and excluding those lost to follow up | Only in laboratory isolates | Gastrointestinal effects (20-55% of patients, usually mild), nephrotoxicity hepatotoxicity possibly teratogenic | About 74 (preferential), about 150 (non-preferential) | Price; possibly teratogenic; potential for resistance (half-life); poor patient compliance |
| Paromomycin sulphate | Institute for One World Health; *Gland Pharma, Hyderabad, India | 15 mg/kg daily for 21 days (India only), intramuscular | Asia: 94% (India); Africa: under evaluation | Only in laboratory isolates | Nephrotoxicity ototoxicity hepatotoxicity (all relatively rare) | About 15 | Efficacy varies between and within regions; potential for resistance? |

* Marketing authorization holder. † Paromomycin was granted orphan drug status by the US Food and Drug Administration and the European Medicines Agency in 2005. Adapted with permission from the Drugs for Neglected Diseases initiative from data presented during Fourth World Congress on Leishmaniasis (Feb 3-7, 2009).

sequential administration of separate drugs (co-administration), and are not co-formulations as used for tuberculosis or malaria.

EFFICACY AND SAFETY

Preclinical data

Few preclinical data on the efficacy and safety of combination therapies for visceral leishmaniasis are available. An early study looked at interactions between sodium stibogluconate and paromomycin (Neal *et al.* 1995). Whereas a marked potentiation was reported against *L. donovani* in vitro, a less-pronounced, additive effect of the antimonial drug was noted in mice (Neal *et al.* 1995). Another study specifically focused on interactions in efficacy between miltefosine and sodium stibogluconate, amphotericin B, paromomycin, and sitamaquine (an oral aminoquinoline) (Seifert *et al.* 2006). In vivo, the highest enhancement of miltefosine activity was seen with amphotericin B, which preceded paromomycin. No activity enhancement was seen with miltefosine combined with sodium stibogluconate. Whereas the combination of miltefosine and amphotericin B could theoretically have some advantages over the other combinations, its clinical relevance remains unknown. More recent findings have also shown a synergistic interaction between amphotericin B and paromomycin (Seifert *et al.* 2009). [https://vpnua2.ua.ac.be/+CSCO+dh756767633A2F2F6A6A2E677572796E617072672E70627A++/journals/laninf/article/PIIS1473-3099\(10\)70011-6/fulltext](https://vpnua2.ua.ac.be/+CSCO+dh756767633A2F2F6A6A2E677572796E617072672E70627A++/journals/laninf/article/PIIS1473-3099(10)70011-6/fulltext) - bib26 Preclinical studies on several drug combinations have been done, with no major safety concerns identified (R Don, Drugs for Neglected Diseases initiative, personal communication, Jan 15, 2010).

Clinical data

The combination of pentavalent antimonials and paromomycin was the first regimen to be studied in India, at a time when clinical failure with pentavalent antimonials was increasingly being reported (Oliaro *et al.* 2005). Overall, these studies showed that 21-day regimen of paromomycin as monotherapy or combined with pentavalent antimonials were efficacious for visceral leishmaniasis. Subsequently, promising data became available on (shortened) monotherapy regimens. A phase 2 study showed that even with liposomal amphotericin B given as a single dose (5 mg/kg), a high proportion of patients could be cured (about 90%) (Sundar *et al.* 2011b; Sundar *et al.* 2003). Equally high proportions could be achieved with 14 days of miltefosine (Sundar *et al.* 2000). These observations provided the rationale for a phase 2, non-comparative randomised trial in India, which assessed different combinations of a single dose of liposomal amphotericin B followed by miltefosine for 7—14 days (table 2)

(Sundar *et al.* 2008b). All combinations were highly efficacious (more than 95% of patients cured) and well tolerated, irrespective of the duration of miltefosine treatment. A phase 2 trial studying the combination of liposomal amphotericin B (5 mg/kg) with miltefosine for 14 days is underway in India and planned in Bangladesh (B Arana, WHO Special Programme for Research and Training in Tropical Diseases, personal communication, Nov 3, 2009) (ClinicalTrials.gov.a). Several short combinations are being studied in a large non-inferiority phase 3 trial in India (table 2), (ClinicalTrials.gov.b) which has now moved into its second stage adding children into the study. In 2009, similar studies will be started in Bangladesh and Nepal; the first results from India are expected by 2010.

In Africa, combination therapy of sodium stibogluconate and paromomycin was studied in the late 1980s, (Chunge *et al.* 1990) and was subsequently used by Médecins Sans Frontières, who needed a shorter treatment regimen when faced with large numbers of patients during an epidemic in Sudan (Davidson *et al.* 2009; Melaku *et al.* 2007; Seaman *et al.* 1996; Seaman *et al.* 1993). Retrospective cohort data from more than 4000 patients showed that, relative to monotherapy with pentavalent antimonials (sodium stibogluconate), combination therapy was associated with clearly reduced mortality and fewer complications during treatment (Melaku *et al.* 2007). This experience formed the basis for the leishmaniasis in east Africa platform (LEAP) 0104 trial, which was started in 2004 (final results are expected in early 2010) (ClinicalTrials.gov. c; Wakabi 2007). This phase 3 trial, which was done in Sudan, Ethiopia, Kenya, and Uganda, initially compared two monotherapy regimens—sodium stibogluconate (20 mg/kg for 30 days) and paromomycin sulphate (15 mg/kg for 21 days)—with the combination of both drugs at the same dose for 17 days (table 2) (ClinicalTrials.gov.c). In 2006, because of unexpectedly low efficacy with paromomycin monotherapy, the protocol was amended, and the dose of paromomycin was increased to 20 mg/kg in the second monotherapy group (Mudawi 2009). Whether the low efficacy related to drug resistance, differences in susceptibility, or pharmacokinetics is currently being investigated.

No large trials on other combination regimens have been done in Africa. Studies on miltefosine and amphotericin B as monotherapy are limited (Berman *et al.* 1998; Mueller *et al.* 2006; Mueller *et al.* 2008; Ritmeijer *et al.* 2006). A phase 2 study, which is due to start recruiting patients early in 2010, will assess the use of miltefosine (including pharmacokinetics) and combinations of liposomal amphotericin B plus either miltefosine or pentavalent antimonial (sodium stibogluconate) in an HIV-negative population (our unpublished data, Drugs for Neglected Diseases initiative). This study will provide important data on the efficacy of miltefosine within a standard clinical trial in east Africa, intends to facilitate drug registration

Table 2: Studies on combination therapy for visceral leishmaniasis in the Indian subcontinent and East Africa

| | Study design | ClinicalTrials.gov registration | Country | Study period | Patients enrolled | Drug combinations studied | Definitive cure (95% CI) at 9 months in intention-to-treat analysis |
|----------------------------------------|---------------------------------------------------------------------------------|---------------------------------|------------------------------------|----------------------------------------------|------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------|
| <i>Completed trials</i> | | | | | | | |
| Sundar <i>et al.</i> ³¹ | Phase 2, non-randomised, non-comparative, group-sequential trial | NCT00370825 | India | 2008-08 | 181 adults: 45 each in groups A, C, D and E; 46 in group B | Group A: SD L-AmB 5 mg/kg alone; Group B: SD L-AmB 5 mg/kg followed by miltefosine 100 mg for 14 days; Group C: SD L-AmB 5 mg/kg followed by miltefosine 100 mg for 10 days; Group D: SD L-AmB 3.75 mg/kg followed by miltefosine 100 mg for 14 days; Group E*: SD L-AmB 5 mg/kg followed by miltefosine 100 mg for 7 days; | Group A: 91% (78-97%); Group B: 98% (87-100%); Group C: 96% (84-99%); Group D: 96% (84-99%); Group E: 98% (87-100%) |
| <i>Planned or ongoing trials</i> | | | | | | | |
| DNDi (VLCOMBO-07 trials) ³³ | Phase 3, randomised, open-label, non-inferiority trial | NCT00696969 | India, Bangladesh, and Nepal | 2008-09 (India), 2009-10 (Bangladesh, Nepal) | 624 adults and children† | Group 1: AmB 1 mg/kg every other day for 30 days; Group 2: SD L-AmB 5 mg/kg followed by miltefosine 2.5 mg/kg for 7 days; Group 3: SD L-AmB 5 mg/kg followed by paromomycin 15 mg/kg for 10 days; Group 4: miltefosine 2.5 mg/kg plus paromomycin 15 mg/kg for 10 days | .. |
| Banaras Hindu University ³² | Phase 2, non-randomised, open-label, historical control, safety/ efficacy trial | NCT00371995 | India | Ongoing | 150 adults and children† | SD L-AmB 5 mg/kg followed by miltefosine 2.5 mg/kg for 14 days | .. |
| TDR‡ | Phase 2 | .. | Bangladesh | Planned | 150† | SD L-AmB 5 mg/kg followed by miltefosine 2.5 mg/kg for 14 days | .. |
| DNDi (LEAP0104A/B trial) ³⁴ | Phase 3, randomised, open-label, active control, safety/ efficacy trial | NCT00255567 | Kenya, Ethiopia, Sudan, and Uganda | 2004-09 (LEAP0104A), 2007-10 (LEAP0104B) | 1100 adults and children† | Active comparator group 1: SSG 20 mg/kg for 30 days; Group 2: paromomycin 15mg/kg (LEAP0104A) or 20 mg/kg (LEAP 0104B) for 21 days; § Group 3: SSG 20 mg/kg plus paromomycin 15mg/kg for 17 days | .. |
| DNDi (VLCOMBO trial) [†] | Phase 2, exploratory, randomised, open-label trial | .. | Sudan and Kenya | Planned for 2009-10 | 189 adults and children; 63 in each group | L-AmB followed by miltefosine; L-AmB followed by SSG; Miltefosine 2.5 mg/kg for 30 days. | .. |

*Non-randomised group assigned when it had become apparent that all other regimens were effective. †Estimated. ‡B Arana, WHO Special Programme for Research and Training in Tropical Diseases, personal communication, Nov 3, 2009. §Due to poor outcomes on paromomycin 15 mg/kg in monotherapy in Sudan in LEAP0104A, the dose for paromomycin monotherapy was increased to 20 mg/kg in LEAP0104B. ¶Our unpublished data. AmB=amphotericin B. DNDi=Drugs for Neglected Diseases initiative. L-AmB=Liposomal amphotericin B. LEAP=leishmaniasis in east Africa platform. SD=single-dose. SSG=sodium stibogluconate. TDR=WHO Special Programme for Research and Training in Tropical Diseases. VLCOMBO=visceral leishmaniasis combination treatment.

in the region, and will also provide the first data on combination therapy based on liposomal amphotericin B. Because pentavalent antimonials are still highly effective in east Africa and have the additional advantage of being cheap, they indeed deserve further exploration in combination regimens when given for shorter duration and at lower total dose.

HEALTH-SYSTEM ISSUES

Access to and capacity of health systems

At present, major barriers exist in terms of access to diagnosis and care for visceral leishmaniasis, monitoring of treatment, and quality assurance of care. A recent study assessed the use of care by patients and delays in diagnosis and treatment in four endemic districts in India, Nepal, and Bangladesh (Mondal *et al.* 2009). In India, even poor people prefer to see a private medical practitioner for treatment of visceral leishmaniasis. In Bangladesh and Nepal, most patients rely on the public-health-care sector, although use of local pharmacists was high in Nepal (Mondal *et al.* 2009). Long delays in diagnosis and treatment were also reported. Overall, 23% of patients interrupted their treatment, mainly because of a lack of resources (67%) or side-effects (16%) (Mondal *et al.* 2009). In Africa, provision of visceral-leishmaniasis care has, until recently, mainly relied on non-governmental organisations. Diverse populations, ranging from semi-nomadic people to migrant workers, need to be reached in different contexts and environments (figure 2) (Desjeux 2004a; Den Boer *et al.* 2006; Desjeux 2001).

Therefore, the identification of an effective, well-tolerated, short regimen that can be given at the primary health-care level, needing little monitoring, and that is affordable both from the patient's and society's perspective would have a major public health impact on these populations. However, combination therapy might bring additional complexity in terms of logistics, service delivery, and programme monitoring. If not carefully considered and planned, combination therapy might actually increase the burden on the health-care system.

The combination of miltefosine and paromomycin might be an interesting option, because it could be given entirely within ambulatory care, and might even be suitable for delivery at the primary-care level. The weak health-care infrastructure would seem to argue against more complex intravenous administrations (e.g., amphotericin B or liposomal amphotericin B) and drugs that need more complicated laboratory and clinical monitoring (e.g., amphotericin B). The need to transport liposomal amphotericin B below 25°C could be another barrier for national programmes, although technological solutions exist that can guarantee transport of drugs below 25°C for 4–5 days. Steps could be taken to strengthen the supply chain of liposomal amphotericin B, and a single dose given at presentation could substantially ease the treatment protocol if the remaining drug could be given on an outpatient basis. Of note,



Figure 2: Visceral leishmaniasis happens in diverse contexts (Images from François Chappuis)
Typical environment in Bihar, India (A). Seminomadic lifestyle in east Africa (B).

ambulatory treatment also needs sufficient capacity to assure good monitoring and follow-up. In cases in which miltefosine is used, an effective system to provide contraception also has to be considered because of the risk of teratogenicity. Despite extensive counselling on contraception, two cases of pregnancy were reported among the 227 women (≥ 12 years) treated in a phase 4 trial of miltefosine in India, with the conception date close to the exposure date (Bhattacharya *et al.* 2007). Fortunately, no birth anomalies were reported.

Cost-effectiveness

Thought must also be given to aspects of cost and cost-effectiveness of the various treatments available. In addition to the public health perspective, it is also important to consider the viewpoint of the household affected by visceral leishmaniasis. Studies found that the median total expenditure by the patient on visceral-leishmaniasis treatment was 1.2 times the annual per head income in Bangladesh (Sharma *et al.* 2006), 1.3 times in India (Meheus *et al.* 2006) and 1.1 times in Nepal (Rijal *et al.* 2006). Only limited data are available on the cost and cost-effectiveness of visceral-leishmaniasis treatment, and studies have mainly focused on monotherapy (Meheus *et al.* 2006; Rijal *et al.* 2006; Vanlerberghe *et al.* 2007). Olliaro and Sunder (Olliaro *et al.* 2009) recently calculated the cost of drugs on the basis of international drug prices and anthropometric data from a single health-care facility in Bihar, India. They found that paromomycin was the cheapest option (US\$7.4 per patient) and that liposomal amphotericin B was the most expensive (\$162–229 per patient) (Olliaro *et al.* 2009). Private

treatment with miltefosine cost \$119 per patient, and \$64—75 at the WHO-negotiated preferential price. However, these calculations did not include other direct or indirect costs, and might differ widely from country to country.

Combination therapies have the potential to reduce the cost to the public health system and patients by reducing the duration of treatment. This not only lowers the burden to the health system but also reduces the economic inactivity of patients. Preliminary findings on the cost-effectiveness of combination therapies in India, Nepal, and Bangladesh, showed combination therapies to be a viable alternative to monotherapies, with liposomal amphotericin B and paromomycin the best combination economically (Meheus *et al.* 2009). However, when we use the preferential drug price of miltefosine in the analysis, it seemed that miltefosine and paromomycin would be the preferred option from an economical perspective (our unpublished observations). No data exist on the cost-effectiveness of combination regimens in east Africa, but a study is due to start in Kenya, Ethiopia, and Sudan, with results expected in 2010 (Meheus F, unpublished).

Regulations and policy

Irrational drug use is a potential threat to the lifespan of any drug, and has probably contributed to the high level of treatment failure with pentavalent antimonials in Bihar, India (Sundar *et al.* 2001a). The unrestricted availability of antimonials in India resulted in widespread misuse by unqualified practitioners, leading to incomplete treatment courses. According to a survey of drug resistance in India, only 26% of patients were treated according to WHO guidelines, and patients often stopped treatment on their own initiative (Sundar *et al.* 1994). The high reliance on the private sector and local pharmacists in the Indian subcontinent even today highlight the need for tightened regulations on the modalities of visceral-leishmaniasis treatment, and for treatment to be made available at no cost (Den Boer *et al.* 2006; Sundar *et al.* 2005; Mondal *et al.* 2009). When policy makers opt for combination therapy, they should take measures to limit the use of monotherapy, particularly in incomplete courses. The fact that miltefosine is available in India without prescription or regulation on dispensed quantities is worrying, since this could facilitate the development of drug resistance (Sundar *et al.* 2005).

PREVENTION OF DRUG RESISTANCE

The problem of drug resistance in visceral leishmaniasis has been extensively reviewed elsewhere (Croft *et al.* 2006). Treatment failure can manifest as initial treatment failure (failure to clear parasites at the end of the treatment course) or relapse (reappearance of parasites after initial cure, usually within 6 months of follow-up). Although pentavalent antimonials have been

successfully used throughout the world for decades, poor treatment response (mainly due to initial treatment failure) has increasingly been reported since the 1980s from Bihar, India, with geographical and temporal clustering in several hyperendemic districts (Sundar 2001a; Peters 1981). Although treatment outcomes could initially be improved with higher total doses, the improvement was only temporary (Thakur *et al.* 1984; Thakur *et al.* 1988; Thakur *et al.* 1991). In subsequent reports, therapy failed in up to 60% of patients that were newly diagnosed (Sundar *et al.* 1995; Sundar *et al.* 1997; Sundar *et al.* 2000b). At the same time, misuse of the drugs was reported (Sundar *et al.* 1994). Increased treatment failure has also been reported in Nepal, in districts that neighbour Bihar (Rijal *et al.* 2003; Rijal *et al.* 2009). Although treatment failure can have several causes, including factors related to drug, host, and parasite, substantial evidence exists that acquired drug resistance is a key issue. Reduced drug sensitivity has been reported with *L. donovani* strains from non-responsive cases in vitro (Lira *et al.* 1999; Laurent *et al.* 2007; Dube *et al.* 2005). Reduced susceptibility to pentavalent antimonials has also been reported with *L. infantum* in both human beings and animals (Carrio *et al.* 2002; Carrio *et al.* 2001; Faraut-Gambarelli *et al.* 1997). In these studies, post-treatment isolates had reduced sensitivity compared with pretreatment isolates, supporting the notion of acquired drug resistance. However, more recent studies have reported less clear associations of in-vitro susceptibility and clinical outcomes, underscoring the need of improved and standardised methods (Rijal *et al.* 2009). The limited understanding of the mechanism of resistance towards pentavalent antimonials, and the shortcomings of drug sensitivity assays, make it difficult to predict the risk of acquired resistance in other regions or drugs and to assess the need for combination therapy to help prevent resistance. However, on the basis of the evidence, acquired drug resistance should be thought to be a potentially serious threat to visceral-leishmaniasis control, and comprehensive strategies should be developed, including the use of combination therapy (Bryceson 2001; Croft *et al.* 2006; Sundar 2001a; den Boer *et al.* 2009).

Rationale

For individual drugs, the ease with which resistance develops will mainly depend on the parasite burden, the probability of spontaneous development of resistance mutations, and the fitness cost associated with those mutations (Hastings 2001). The level and pattern of drug use in a population constitutes the selection force for the development of resistance, and intact host immunity is generally thought protective. The potency of the drug, therapeutic index, and pharmacokinetic properties of the drug also play a part (Hastings 2001). Combination therapy could delay resistance if two drugs with different modes of action and mechanisms of resistance are used. The combination of synergistic drugs is preferred, because if more effective replication can be inhibited, resistance is less likely during treatment.

For resistance prevention, both drugs should ideally have similar pharmacokinetics. If parasites always confront both drugs, the probability of the emergence of double-resistant parasites would be expected to be extremely rare (ie, the product of their individual per-parasite probabilities). A rapid elimination phase minimises the duration of subtherapeutic drug concentrations, which could provide an opportunity for amplification or selection of resistant parasites (Bryceson 2001; Hastings 2001; Stepniewska *et al.* 2008). In studies of malaria, the combination of one very active drug with a short half-life with a slower acting drug with a longer half-life to clear the remaining parasites has been explored as a way to shorten treatment duration and improve treatment compliance (Nosten *et al.* 2007). However, recent studies have focused on the terminal elimination phase of the second drug, which can act as a selective filter for resistant malaria parasites (Pongtavornpinyo *et al.* 2008; White 2004). Artemisinin resistance has recently been reported in Cambodia, but underlying reasons remain to be established (Noedl *et al.* 2008; Noedl *et al.* 2009; Dondorp *et al.* 2009). Finally, drugs can be combined to target different biological stages of the infectious agent. This has been done for tuberculosis and malaria, although the drugs are essentially targeted at preventing relapse and would only indirectly prevent or delay resistance.

Pharmacological considerations for combination therapy

Although the mechanisms of action and resistance remain poorly understood for all anti-leishmanial drugs in use (except amphotericin B), they are all thought to act on different targets (Alvar *et al.* 2006). Recent findings from India suggest that field isolates from areas with high-level resistance to pentavalent antimonials show reduced sensitivity towards other anti-leishmanial drugs such as amphotericin B and miltefosine (Prajapati 2009; Kumar *et al.* 2009). However, true cross-resistance between the various drugs has not been reported so far. Several combinations have shown activity enhancement in animal experiments (Seifert *et al.* 2006; Seifert *et al.* 2009).

Clear differences in pharmacokinetics exist (Bryceson 2001). Miltefosine might be particularly vulnerable to the emergence of resistance, because of its narrow therapeutic index and long half-life, which has been estimated at around 7 days (Perez-Victoria *et al.* 2006; Berman *et al.* 2006). Recent data from patients with cutaneous leishmaniasis suggested a terminal half-life of 31 days, with miltefosine still detectable 5–6 months after the end of treatment (Dorlo *et al.* 2008). Resistant strains could be selected and amplified during this period because of subtherapeutic drug concentrations, either from relapsing patients, or from newly acquired infections (Perez-Victoria *et al.* 2006; Berman *et al.* 2006). If confirmed, this might have important repercussions on the risk of emerging resistance and on the duration of contraceptive measures.

Paromomycin has a short half-life (2–3 h in patients without visceral leishmaniasis), but has a low therapeutic index. Resistance can easily be induced in vitro (Maarouf *et al.* 1998), clinical resistance has been noted with its antibacterial use (Bryceson 2001; Davidson *et al.* 2009; Teklemariam *et al.* 1994), and some have argued against its use in monotherapy (Den Boer *et al.* 2006). Most of a pentavalent antimonial (about 99%) is eliminated within a few hours, followed by a slower elimination phase with a half-life of 76 h (Chulay *et al.* 1988). At least in east Africa, these drugs remain highly effective.

Amphotericin B could be thought less likely to induce resistance given its high efficacy and a relatively short half-life of 24 h (Bryceson 2001; Croft *et al.* 2006; Di Giorgio *et al.* 1999; Durand *et al.* 1998; Lachaud *et al.* 2009). Although resistance can be induced in vitro, clinical cases of amphotericin-B resistance have not been reported (Singh *et al.* 2001; Papadopoulou *et al.* 1998; Ouellette *et al.* 2004; Mbongo *et al.* 1998). Liposomal amphotericin B has a bioavailability in tissues for several weeks despite a relatively short plasma half-life (Gagneux *et al.* 1996; Adler-Moore *et al.* 2002). Given this long tissue half-life, a single dose of liposomal amphotericin B followed by daily administration of a second drug (eg, sodium stibogluconate, paromomycin, or miltefosine) would result in simultaneous exposure of the parasite to both drugs. The use of a single dose of liposomal amphotericin B (10 mg/kg) in monotherapy is being explored in India (ClinicalTrials.gov. d). Although this might be a simple and effective approach, concerns of resistance and cost should also be taken into account.

Compliance with treatment

Besides the intrinsic characteristics of the combination regimen, the use of and compliance to the regimen also affects the risk of drug resistance. All conventional monotherapies (apart from liposomal amphotericin B) need a long treatment duration (21–30 days), making compliance more challenging. This is of particular concern for treatment with miltefosine, the only oral drug, for which the risk of premature treatment interruption is high. Even in a phase 4 trial, 4.5% of patients were lost to follow-up before the end of treatment, and 14.5% were not available for assessment by 6 months after treatment (Bhattacharya *et al.* 2007). High default rates (up to 30%) were noted in a pilot study of miltefosine in India in ten districts (our unpublished data). Shorter treatment duration, particularly if the drug is also more tolerable, might help to increase compliance, as has been the case for patients on combination regimens for malaria (Bryceson 2001).

The lower costs to patients associated with shortened combination therapy could also improve access to and acceptability of visceral-leishmaniasis treatment. Some have suggested that the directly observed treatment strategy, which has been successfully used for tuberculosis,

might have a role in ensuring good compliance to miltefosine, although this will increase the indirect and direct costs (Croft *et al.* 2006; Sundar *et al.* 2007; Perez-Victoria *et al.* 2006). The elimination programme for visceral leishmaniasis in south Asia has opted for miltefosine as first-line drug, but will need to engage in the monitoring of clinical treatment outcomes and pharmacovigilance to ensure effective management (Sundar *et al.* 2007).

SPECIAL POPULATIONS: HIV COINFECTION

In east Africa, coinfection with HIV is a major challenge in the treatment of visceral leishmaniasis, with up to 30% of cases infected with HIV in some regions (Den Boer *et al.* 2006; Alvar *et al.* 2008; Lyons *et al.* 2003). This problem also seems to be on the increase in the Indian subcontinent (Mathur *et al.* 2006; Redhu *et al.* 2006). Because asymptomatic leishmania infections are thought to outnumber symptomatic infections, (Chappuis *et al.* 2007; Guerin *et al.* 2002) the dramatically increased risk of progression to visceral leishmaniasis after infection with HIV could lead to increased disease burden (Alvar *et al.* 2008), as has also been seen with tuberculosis (Harries 1990). Additionally, coinfection with leishmania and HIV is associated with a high mortality and a high rate of treatment failure and relapse with all visceral-leishmaniasis drugs (Alvar *et al.* 2008; Ritmeijer *et al.* 2006; Ter Horst *et al.* 2008; Ritmeijer *et al.* 2009). The efficacy of combination therapy for coinfection with leishmania and HIV has not yet been studied in a controlled trial.

Although widespread use of antiretroviral therapy has resulted in large reductions in the incidence of visceral-leishmaniasis—HIV coinfection in southern Europe, it seems to be only partly protective against relapses (Alvar *et al.* 2008; Ter Horst *et al.* 2008). Patients with incurable disease (who present with relapse or post kala-azar dermal leishmaniasis) could serve as a reservoir for the development and spread of drug-resistant strains, particularly because such patients seem to be more infectious (Molina *et al.* 2003). To increase efficacy, combination therapy might be particularly relevant for coinfecting patients to prevent resistance, because repeated exposure to single anti-leishmanial drugs will facilitate the emergence of resistant parasites. Even if relapses cannot be prevented, combination therapy might preserve drug sensitivity. Although one study reported decreased sensitivity towards amphotericin B after several treatment courses in individuals infected with HIV (Di Giorgio *et al.* 1999), this has not been confirmed (Durand *et al.* 1998; Lachaud *et al.* 2009).

Given the overall poor treatment outcomes, patients infected with HIV are most likely to need a different first-line therapy from patients not infected with HIV, at least in terms of treatment duration. However, none of the planned or continuing phase 3 trials include patients infected with HIV. Whether secondary prophylaxis could prevent relapses remains unclear. WHO does not recommend secondary prophylaxis in foci of anthroponotic visceral leishmaniasis (WHO 2007).

FUTURE PERSPECTIVES

Although the principle of combination therapy is generally accepted, the rationale for the choice of the drugs in such combination regimens is still under debate (Bryceson 2001; Croft *et al.* 2006; Den Boer *et al.* 2006). Several combinations have been taken forward in clinical studies, which will most likely provide us with several efficacious treatment options in the near future. However, it remains to be defined what exactly we should expect from combination therapy, and which factors will be important in selecting a specific therapy within a given context.

The current or planned clinical trials on combination therapy for visceral leishmaniasis in Asia and Africa will essentially provide data on safety and efficacy of the different regimens. Whereas drug—drug interactions, both in terms of pharmacokinetics and toxicity, have been prime considerations in the development of combination therapy for other infectious diseases, only limited data on anti-leishmanial drugs are available. Since the leishmania parasite targets reticulo-endothelial cells within specific organs, factors such as tissue distribution and uptake into macrophages of the individual components within combination therapy might also be relevant (Seifert *et al.* 2006). Whereas reduced toxic effects with combination therapy could possibly improve drug tolerance, because of reduced total doses of the individual compounds, increased toxicity could also be possible. Although the available data from animal and human studies seem reassuring, more studies to address these issues should be done. At least some of the combination studies that have been planned by the Drugs for Neglected Disease initiative will include pharmacokinetic substudies (our unpublished data).

Forthcoming results of clinical trials will not address the future public health benefits of combination therapy nor how it should be applied in field settings. Particularly, data from east Africa on liposomal amphotericin B suggest that many patients present with advanced and complicated visceral leishmaniasis and that perhaps relatively intensive treatment might be needed for cure (Berman *et al.* 1998; Collin *et al.* 2004; Mueller *et al.* 2007; Seaman *et al.* 1995). Because these patients are generally not included in phase 3 trials, these issues should be further addressed in field studies. Even in the recently reported phase 4 trial of miltefosine

in India, 10% of patients were excluded (Bhattacharya *et al.* 2007). More research on safety issues also needs to be done.

It will take several years before sufficient clinical data will be available and combination therapy will be effectively implemented in the field. In the meantime, care should be taken to minimise the development of resistance to the individual drugs. Compared with miltefosine, liposomal amphotericin B might be a more cautious option, because substantial resistance towards liposomal amphotericin B is unlikely to happen when used as large-scale monotherapy for a few years. Recent data from India on the use of standard dose of liposomal amphotericin B in operational settings have shown excellent treatment outcomes (Lima 2009).

Most likely, regional factors will play an important part in deciding on the best combination therapy for a particular area, region, or country. Regional differences in natural or acquired drug resistance or pharmacokinetics should be considered. Differences in nutritional status and prevalence of coinfections such as HIV and tuberculosis might also determine the choice of combination therapies. Finally, regional factors will also determine which combination could be made available with the largest coverage and compliance in a sustainable and stable manner, and which delivery model would be most cost-effective.

KEY QUESTIONS AND CHALLENGES

Trial data of several combination therapies will become available over the next few years, and should address some of the issues that have been discussed. However, many questions and challenges remain (panel). Whereas combination therapy aims to increase the lifespan of available drugs, this could also lead to rapid loss of two therapeutic options, if not applied in a controlled and regulated way. Care should be taken to ensure that the increased complexity of the logistics of combination therapy does not hamper effective implementation. The evidence base for combination therapy should be regularly re-assessed within each specific context.

How an effective treatment strategy should be integrated within the private and public-health sectors and how this determines the choice of (combination) regimen are unclear. Without increased access to and capacity of health care, both in terms of diagnosis and treatment, the overall effect of any (combination) therapy will be small. Efforts to reduce transmission also need to be addressed. The factors that affect the acceptability of visceral-leishmaniasis care and how to improve it will also need to be taken into account.

Close monitoring of combination therapy will be important at several levels. Programme monitoring will allow assessment of the overall effect, to identify barriers, and to supervise

the correct implementation and use of combination therapy. Systems should also be put in place to detect counterfeit drugs or illegal drug supply. In areas with high prevalence of HIV or tuberculosis, close links and integration with the national programmes will be pivotal.

Panel: Key questions and challenges

- Key questions and challenges
- To what extent are the available anti-leishmanial drugs threatened by the development of drug resistance when used in monotherapy?
- Will combination therapy effectively delay acquired drug resistance and how is this best achieved?
- How do we set up an effective surveillance system to detect the development of acquired drug resistance?
- What will be the field efficacy of combination therapy?
- How do we set up a pharmacovigilance system?
- How do we assure good compliance with ambulatory treatment?
- Is there a need for a (daily) directly observed therapy strategy for miltefosine?
- How do we effectively implement a combination therapy strategy and, at the same time, increase access to care and diagnostic and therapeutic capacity?
- What is most cost-effective and feasible delivery model for visceral-leishmaniasis care in a specific setting?
- How do we regulate and control prescription of combination therapy, and how do we detect or prevent extra-legal drug use?
- What is the place of the private health care in visceral-leishmaniasis treatment?
- Do public—private partnerships have a place in the provision of care and treatment for visceral leishmaniasis?

Although reliable programme data can show the effectiveness of treatment regimens in use, they do not allow emerging resistance to be traced at an early stage. Parasite drug susceptibility should be monitored within surveillance systems, as being deployed for tuberculosis, malaria, and HIV. Because in-vitro susceptibility assays used at present have substantial limitations, further optimisation and standardisation remain necessary (Croft 2001; Rijal *et al.* 2007). The development of molecular markers of resistance, as pursued by several research groups, could be a way forward in the long run if also made available in endemic countries (Perez-Victoria *et al.* 2006). Moreover, visceral-leishmaniasis therapies carry a substantial risk of toxic effects, and there are is obviously no long-term experience with combination therapy. Particularly if miltefosine is to be used on a large scale, problems related to teratogenicity or other unknown side-effects should be traced at an early stage.

Whether combination therapy also has an important place in the prevention of drug resistance, and how this is best achieved, will only be known in the long run. Aiming for a combination therapy that is highly effective, widely accessible, affordable, and with high rates of compliance should probably be the priority.

Search strategy and selection criteria

Articles cited in this Review were obtained through searches of PubMed or Medline for papers published up to June 1, 2009, with terms including, but not restricted to, the following combinations: “visceral leishmaniasis”, “treatment”, “combination therapy”, “safety”, “efficacy”, “cost”, “cost-effectiveness”, “resistance”, “compliance”, “health system”, and “human immunodeficiency virus”. The search was limited to English. Reference lists of these articles were then searched to identify other relevant publications. Clinical trial websites were verified, and experts in clinical research in visceral leishmaniasis were contacted for information on planned or ongoing trials. Abstracts of recent international conferences on infectious diseases were also reviewed. We structured our Review around the following criteria: efficacy and safety, health-system aspects (feasibility, cost-effectiveness and regulations), potential for drug resistance, and use in special populations.

REFERENCES

- Adler-Moore J, Proffitt RT (2002) AmBisome: liposomal formulation, structure, mechanism of action and pre-clinical experience. *J Antimicrob Chemother* 49 (S1): 21–30.
- Alvar J, Aparicio P, Aseffa A, et al. (2008) The relationship between leishmaniasis and AIDS: the second 10 years. *Clin Microbiol Rev* 21: 334–59.
- Alvar J, Croft S, Olliaro P (2006) Chemotherapy in the treatment and control of leishmaniasis. *Adv Parasitol* 61: 223–74.
- Berman J, Bryceson AD, Croft S, et al. (2006) Miltefosine: issues to be addressed in the future. *Trans R Soc Trop Med Hyg* 100(S1): S41–S44.
- Berman JD (2006) Development of miltefosine for the leishmaniasis. *Mini Rev Med Chem* 6: 145–51.
- Berman JD, Badaro R, Thakur CP, et al. (1998) Efficacy and safety of liposomal amphotericin B (AmBisome) for visceral leishmaniasis in endemic developing countries. *Bull World Health Organ* 76: 25–32.
- Bern C, Adler-Moore J, Berenguer J, et al. (2006) Liposomal amphotericin B for the treatment of visceral leishmaniasis. *Clin Infect Dis* 43: 917–24.
- Bhattacharya SK, Sinha PK, Sundar S, et al. (2007) Phase 4 trial of miltefosine for the treatment of Indian visceral leishmaniasis. *J Infect Dis* 196: 591–98.
- Bryceson A (2001) A policy for leishmaniasis with respect to the prevention and control of drug resistance. *Trop Med Int Health* 6: 928–34.
- Carrio J, Portus M (2002) In vitro susceptibility to pentavalent antimony in *Leishmania infantum* strains is not modified during in vitro or in vivo passages but is modified after host treatment with meglumine antimoniate. *BMC Pharmacol* 2: 11.
- Carrio J, Riera C, Gallego M, Ribera E, Portus M (2001) In vitro susceptibility of *Leishmania infantum* to meglumine antimoniate in isolates from repeated leishmaniasis episodes in HIV-coinfected patients. *J Antimicrob Chemother* 47: 120–21.
- Chappuis F, Sundar S, Hailu A, et al. (2007) Visceral leishmaniasis: what are the needs for diagnosis, treatment and control? *Nat Rev Microbiol* 5: 873–82.
- Chulay JD, Fleckenstein L, Smith DH (1988) Pharmacokinetics of antimony during treatment of visceral leishmaniasis with sodium stibogluconate or meglumine antimoniate. *Trans R Soc Trop Med Hyg* 82: 69–72.
- Chunge CN, Owate J, Pamba HO, Donno L (1990) Treatment of visceral leishmaniasis in Kenya by aminosidine alone or combined with sodium stibogluconate. *Trans R Soc Trop Med Hyg* 84: 221–25.
- ClinicalTrials.gov. Efficacy/safety of sodium stibogluconate (SSG) versus paromomycin (PM) and SSG/PM combination to treat V leishmaniasis. NCT00255567. <http://clinicaltrials.gov/ct2/show/NCT00255567>. Accessed on: Jan 23, 2010.
- ClinicalTrials.gov. Safety and efficacy study to evaluate different combination treatment regimens for visceral leishmaniasis. NCT00696969. <http://clinicaltrials.gov/ct2/show/NCT00696969>. Accessed on: Jan 23, 2010.

- ClinicalTrials.gov. Short course of miltefosine and liposomal amphotericin B for kala-azar. NCT00371995. <http://clinicaltrials.gov/ct2/show/NCT00371995>. Accessed on: Jan 23, 2010.
- ClinicalTrials.gov. Single infusion of liposomal amphotericin B in Indian visceral leishmaniasis. NCT00628719. <http://clinicaltrials.gov/ct2/show/NCT00628719> Accessed on: Jan 25, 2010.
- Collin S, Davidson R, Ritmeijer K, et al. (2004) Conflict and kala-azar: determinants of adverse outcomes of kala-azar among patients in southern Sudan. *Clin Infect Dis* 38: 612–19.
- Croft SL (2001) Monitoring drug resistance in leishmaniasis. *Trop Med Int Health* 6: 899–905.
- Croft SL, Engel J (2006) Miltefosine - discovery of the antileishmanial activity of phospholipid derivatives. *Trans R Soc Trop Med Hyg* 100(S1): S4–S8.
- Croft SL, Sundar S, Fairlamb AH (2006) Drug resistance in leishmaniasis. *Clin Microbiol Rev* 19: 111–26.
- Davidson RN, Den Boer M, Ritmeijer K (2009) Paromomycin. *Trans R Soc Trop Med Hyg* 103: 653–60.
- Den Boer M, Davidson RN (2006) Treatment options for visceral leishmaniasis. *Expert Rev Anti Infect Ther* 4: 187–97.
- den Boer ML, Alvar J, Davidson RN, Ritmeijer K, Balasegaram M (2009) Developments in the treatment of visceral leishmaniasis. *Expert Opin Emerg Drugs* 14: 395–410.
- Desjeux P (2001) The increase in risk factors for leishmaniasis worldwide. *Trans R Soc Trop Med Hyg* 95: 239–43.
- Desjeux P (2004) Leishmaniasis. *Nat Rev Microbiol* 2: 692–93.
- Desjeux P (2004) Leishmaniasis: current situation and new perspectives. *Comp Immunol Microbiol Infect Dis* 27: 305–18.
- Di Giorgio C, Faraut-Gambarelli F, Imbert A, Minodier P, Gasquet M, Dumon H (1999) Flow cytometric assessment of amphotericin B susceptibility in *Leishmania infantum* isolates from patients with visceral leishmaniasis. *J Antimicrob Chemother* 44: 71–76.
- Dondorp AM, Nosten F, Yi P, et al. (2009) Artemisinin resistance in *Plasmodium falciparum* malaria. *N Engl J Med* 361: 455–67.
- Dorlo TP, van Thiel PP, Huitema AD, et al. (2008) Pharmacokinetics of miltefosine in Old World cutaneous leishmaniasis patients. *Antimicrob Agents Chemother* 52: 2855–60.
- Dube A, Singh N, Sundar S, Singh N (2005) Refractoriness to the treatment of sodium stibogluconate in Indian kala-azar field isolates persist in in vitro and in vivo experimental models. *Parasitol Res* 96: 216–23.
- Durand R, Paul M, Pratlong F, et al. (1998) *Leishmania infantum*: lack of parasite resistance to amphotericin B in a clinically resistant visceral leishmaniasis. *Antimicrob Agents Chemother* 42: 2141–43.
- Faraut-Gambarelli F, Piarroux R, Deniau M, et al. (1997) In vitro and in vivo resistance of *Leishmania infantum* to meglumine antimoniate: a study of 37 strains collected from patients with visceral leishmaniasis. *Antimicrob Agents Chemother* 41:827–30.


- Gangneux JP, Sulahian A, Garin YJ, Farinotti R, Derouin F (1996) Therapy of visceral leishmaniasis due to *Leishmania infantum*: experimental assessment of efficacy of AmBisome. *Antimicrob Agents Chemother* 40: 1214–18.
- Guerin PJ, Olliaro P, Sundar S, et al. (2002) Visceral leishmaniasis: current status of control, diagnosis, and treatment, and a proposed research and development agenda. *Lancet Infect Dis* 2: 494–501.
- Harries AD (1990) Tuberculosis and human immunodeficiency virus infection in developing countries. *Lancet* 335: 387–90.
- Hastings IM (2001) Modelling parasite drug resistance: lessons for management and control strategies. *Trop Med Int Health* 6: 883–90.
- Herwaldt BL (1999) Leishmaniasis. *Lancet* 354: 1191–99.
- Kumar D, Kulshrestha A, Singh R, Salotra P (2009) In vitro susceptibility of field isolates of *Leishmania donovani* to miltefosine and amphotericin B: correlation with sodium antimony gluconate susceptibility and implications for treatment in areas of endemicity. *Antimicrob Agents Chemother* 53: 835–38.
- Lachaud L, Bourgeois N, Plourde M, Leprohon P, Bastien P, Ouellette M (2009) Parasite susceptibility to amphotericin B in failures of treatment for visceral leishmaniasis in patients coinfectd with HIV type 1 and *Leishmania infantum*. *Clin Infect Dis* 48: e16–e22.
- Laurent T, Rijal S, Yardley V, et al. (2007) Epidemiological dynamics of antimonial resistance in *Leishmania donovani*: genotyping reveals a polyclonal population structure among naturally-resistant clinical isolates from Nepal. *Infect Genet Evol* 7: 206–12.
- Lima N (2009) Experience of Médecins Sans Frontières after 1 year of using liposomal amphotericin B (AmBisome) treatment for Indian visceral leishmaniasis under routine programme conditions in Bihar, India. Proceedings of the 4th World Congress on Leishmaniasis 2009 (WorldLeish4), Lucknow, India Feb 3–7.
- Lira R, Sundar S, Makharia A, et al. (1999) Evidence that the high incidence of treatment failures in Indian kala-azar is due to the emergence of antimony-resistant strains of *Leishmania donovani*. *J Infect Dis* 180: 564–67.
- Lyons S, Veeken H, Long J (2003) Visceral leishmaniasis and HIV in Tigray, Ethiopia. *Trop Med Int Health* 8: 733–39.
- Maarouf M, Adeline MT, Solignac M, Vautrin D, Robert-Gero M (1998) Development and characterization of paromomycin-resistant *Leishmania donovani* promastigotes. *Parasite* 5: 167–73.
- Maltezou HC (2008) Visceral leishmaniasis: advances in treatment. *Recent Pat Antiinfect Drug Discov* 3: 192–98.
- Mathur P, Samantaray JC, Vajpayee M, Samanta P (2006) Visceral leishmaniasis/human immunodeficiency virus co-infection in India: the focus of two epidemics. *J Med Microbiol* 55: 919–22.
- Mbongo N, Loiseau PM, Billion MA, Robert-Gero M (1998) Mechanism of amphotericin B resistance in *Leishmania donovani* promastigotes. *Antimicrob Agents Chemother* 42: 352–57.

- Meheus F, Olliaro P, Rijal S, Sundar S, Faiz M, Boelaert M (2009) Preliminary cost-effectiveness analysis of combination therapies to combat parasite resistance to antileishmanial drugs in the Indian subcontinent. Proceedings of the 4th World Congress on Leishmaniasis 2009 (WorldLeish4), Lucknow, India Feb 3–7, 2009. Abstract 691.
- Meheus F, Boelaert M, Baltussen R, Sundar S (2006) Costs of patient management of visceral leishmaniasis in Muzaffarpur, Bihar, India. *Trop Med Int Health* 11: 1715–24.
- Melaku Y, Collin SM, Keus K, Gatluak F, Ritmeijer K, Davidson RN (2007) Treatment of kala-azar in southern Sudan using a 17-day regimen of sodium stibogluconate combined with paromomycin: a retrospective comparison with 30-day sodium stibogluconate monotherapy. *Am J Trop Med Hyg* 77: 89–94.
- Molina R, Gradoni L, Alvar J (2003) HIV and the transmission of *Leishmania*. *Ann Trop Med Parasitol* 97(S1): 29–45.
- Mondal D, Singh SP, Kumar N, et al. (2009) Visceral leishmaniasis elimination programme in India, bangladesh, and Nepal: reshaping the case finding/case management strategy. *PLoS Negl Trop Dis* 3: e355.
- Mudawi AM (2009) New treatments for visceral leishmaniasis in east Africa: the story so far. Proceedings of the 4th World Congress on Leishmaniasis 2009 (WorldLeish4), Lucknow, India Feb 3-7. http://www.worldleish4.org/program_details.php?controlId=symposia&&symp=dndi. Accessed on: Jan 18, 2010.
- Mueller M, Balasegaram M, Koummuki Y, Ritmeijer K, Santana MR, Davidson R (2006) A comparison of liposomal amphotericin B with sodium stibogluconate for the treatment of visceral leishmaniasis in pregnancy in Sudan. *J Antimicrob Chemother* 58: 811–15.
- Mueller M, Ritmeijer K, Balasegaram M, Koummuki Y, Santana MR, Davidson R (2007) Unresponsiveness to AmBisome in some Sudanese patients with kala-azar. *Trans R Soc Trop Med Hyg* 101: 19–24.
- Mueller Y, Nguimfack A, Cavailler P, et al. (2008) Safety and effectiveness of amphotericin B deoxycholate for the treatment of visceral leishmaniasis in Uganda. *Ann Trop Med Parasitol* 102: 11–19.
- Neal RA, Allen S, McCoy N, Olliaro P, Croft SL (1995) The sensitivity of *Leishmania* species to aminosidine. *J Antimicrob Chemother* 35: 577–84.
- Noedl H, Se Y, Schaecher K, Smith BL, Socheat D, Fukuda MM (2008) Evidence of artemisinin-resistant malaria in western Cambodia. *N Engl J Med* 359: 2619–20.
- Noedl H, Socheat D, Satimai W (2009) Artemisinin-resistant malaria in Asia. *N Engl J Med* 361: 540–41.
- Nosten F, White NJ (2007) Artemisinin-based combination treatment of falciparum malaria. *Am J Trop Med Hyg* 77: 181–92.
- Olliaro P, Sundar S (2009) Anthropometrically derived dosing and drug costing calculations for treating visceral leishmaniasis in Bihar, India. *Trop Med Int Health* 14: 88–92.
- Olliaro PL, Guerin PJ, Gerstl S, Haaskjold AA, Rottingen JA, Sundar S (2005) Treatment options for visceral leishmaniasis: a systematic review of clinical studies done in India, 1980—2004. *Lancet Infect Dis* 5: 763–74.

- Ouellette M, Drummelsmith J, Papadopoulou B (2004) Leishmaniasis: drugs in the clinic, resistance and new developments. *Drug Resist Updat* 7: 257–66.
- Papadopoulou B, Kundig C, Singh A, Ouellette M (1998) Drug resistance in *Leishmania*: similarities and differences to other organisms. *Drug Resist Updat* 1: 266–78.
- Perez-Victoria FJ, Sánchez-Cañete MP, Seifert K, et al. (2006) Mechanisms of experimental resistance of *Leishmania* to miltefosine: implications for clinical use. *Drug Resist Updat* 9: 26–39.
- Peters W (1981) The treatment of kala-azar – new approaches to an old problem. *Indian J Med Res* 73(S): 1–18.
- Pongtavornpinyo W, Yeung S, Hastings IM, Dondorp AM, Day NP, White NJ (2008) Spread of anti-malarial drug resistance: mathematical model with implications for ACT drug policies. *Malar J* 7: 229.
- Prajapati V (2009) Intrinsic macrophage sensitivity of different field isolates in respect to antileishmanial drugs. Proceedings of the 4th World Congress on Leishmaniasis 2009 (WorldLeish4), Lucknow, India Feb 3–7, 2009. Abstract 134.
- Redhu NS, Dey A, Balooni V, Singh S (2006) *Leishmania*—HIV co-infection: an emerging problem in India. *AIDS* 20: 1213–15.
- Rijal S, Bhandari S, Koirala S, et al. (2009) Clinical risk factors for therapeutic failure in kala-azar patients treated with pentavalent antimonials in Nepal. *Trans R Soc Trop Med Hyg* 104: 225–29
- Rijal S, Chappuis F, Singh R, et al. (2003) Treatment of visceral leishmaniasis in south-eastern Nepal: decreasing efficacy of sodium stibogluconate and need for a policy to limit further decline. *Trans R Soc Trop Med Hyg* 97: 350–54.
- Rijal S, Koirala S, Van der Stuyft P, Boelaert M (2006) The economic burden of visceral leishmaniasis for households in Nepal. *Trans R Soc Trop Med Hyg* 100: 838–41.
- Rijal S, Yardley V, Chappuis F, et al. (2007) Antimonial treatment of visceral leishmaniasis: are current in vitro susceptibility assays adequate for prognosis of in vivo therapy outcome? *Microbes Infect* 9: 529–35.
- Ritmeijer K, Ter Horst R, Chane S, Davidson RN (2009) Poor effectiveness of liposomal amphotericin-B (Ambisome) in HIV co-infected visceral leishmaniasis patients in Ethiopia. 2009. Proceedings of the 4th World Congress on Leishmaniasis 2009 (WorldLeish4), Lucknow, India Feb 3–7, 2009. Abstract 159.
- Ritmeijer K, Dejenie A, Assefa Y, et al. (2006) A comparison of miltefosine and sodium stibogluconate for treatment of visceral leishmaniasis in an Ethiopian population with high prevalence of HIV infection. *Clin Infect Dis* 43: 357–64.
- Seaman J, Boer C, Wilkinson R, et al. (1995) Liposomal amphotericin B (AmBisome) in the treatment of complicated kala-azar under field conditions. *Clin Infect Dis* 21: 188–93.
- Seaman J, Mercer AJ, Sondorp HE, Herwaldt BL (1996) Epidemic visceral leishmaniasis in southern Sudan: treatment of severely debilitated patients under wartime conditions and with limited resources. *Ann Intern Med* 124: 664–72.

- Seaman J, Pryce D, Sondorp HE, Moody A, Bryceson AD, Davidson RN (1993) Epidemic visceral leishmaniasis in Sudan: a randomized trial of aminosidine plus sodium stibogluconate versus sodium stibogluconate alone. *J Infect Dis* 168: 715–20.
- Seifert K, Syeda T, Munday J, Croft S (2009) Drug combination studies in experimental leishmaniasis models—lessons learned and future directions. Proceeding of the 4th World Congress on Leishmaniasis (WorldLeish4), Lucknow, India Feb 3–7, 2009. Abstract 423.
- Seifert K, Croft SL (2006) In vitro and in vivo interactions between miltefosine and other antileishmanial drugs. *Antimicrob Agents Chemother* 50: 73–79.
- Seifert K, Perez-Victoria FJ, Stettler M, et al. (2007) Inactivation of the miltefosine transporter, LdMT, causes miltefosine resistance that is conferred to the amastigote stage of *Leishmania donovani* and persists in vivo. *Int J Antimicrob Agents* 30: 229–35.
- Sharma A, Bern C, Varghese B, et al. (2006) The economic impact of visceral leishmaniasis on households in Bangladesh. *Trop Med Int Health* 11: 757–64.
- Singh AK, Papadopoulou B, Ouellette M (2001) Gene amplification in amphotericin B-resistant *Leishmania tarentolae*. *Exp Parasitol* 99: 141–47.
- Singh RK, Pandey HP, Sundar S (2006) Visceral leishmaniasis (kala-azar): challenges ahead. *Indian J Med Res* 123: 331–44.
- Stepniewska K, White NJ (2008) Pharmacokinetic determinants of the window of selection for antimalarial drug resistance. *Antimicrob Agents Chemother* 52: 1589–96.
- Sundar S (2001) Drug resistance in Indian visceral leishmaniasis. *Trop Med Int Health* 6: 849–54.
- Sundar S, Agrawal G, Rai M, Makharia MK, Murray HW (2001) Treatment of Indian visceral leishmaniasis with single or daily infusions of low dose liposomal amphotericin B: randomised trial. *BMJ* 323: 419–22.
- Sundar S, Chakravarty J (2008) Paromomycin in the treatment of leishmaniasis. *Expert Opin Investig Drugs* 17: 787–94.
- Sundar S, Jha TK, Thakur CP, Mishra M, Singh VP, Buffels R (2003) Single-dose liposomal amphotericin B in the treatment of visceral leishmaniasis in India: a multicenter study. *Clin Infect Dis* 37: 800–04.
- Sundar S, Makharia A, More DK, et al. (2000) Short-course of oral miltefosine for treatment of visceral leishmaniasis. *Clin Infect Dis* 31: 1110–13.
- Sundar S, More DK, Singh MK, et al. (2000) Failure of pentavalent antimony in visceral leishmaniasis in India: report from the center of the Indian epidemic. *Clin Infect Dis* 31: 1104–07.
- Sundar S, Murray HW (2005) Availability of miltefosine for the treatment of kala-azar in India. *Bull World Health Organ* 83: 394–95.
- Sundar S, Oliaro PL (2007) Miltefosine in the treatment of leishmaniasis: clinical evidence for informed clinical risk management. *Ther Clin Risk Manag* 3: 733–40.

- Sundar S, Rai M, Chakravarty J, et al. (2008) New treatment approach in Indian visceral leishmaniasis: single-dose liposomal amphotericin B followed by short-course oral miltefosine. *Clin Infect Dis* 47: 1000–06.
- Sundar S, Rosenkaimer F, Lesser ML, Murray HW (1995) Immunochemotherapy for a systemic intracellular infection: accelerated response using interferon-gamma in visceral leishmaniasis. *J Infect Dis* 171: 992–96.
- Sundar S, Singh VP, Sharma S, Makharia MK, Murray HW (1997) Response to interferon-gamma plus pentavalent antimony in Indian visceral leishmaniasis. *J Infect Dis* 176: 1117–19.
- Sundar S, Thakur BB, Tandon AK, et al. (1994) Clinicoepidemiological study of drug resistance in Indian kala-azar. *BMJ* 308: 307.
- Teklemariam S, Hiwot AG, Frommel D, Miko TL, Ganlov G, Bryceson A (1994) Aminosidine and its combination with sodium stibogluconate in the treatment of diffuse cutaneous leishmaniasis caused by *Leishmania aethiopica*. *Trans R Soc Trop Med Hyg* 88: 334–39.
- Ter Horst R, Collin SM, Ritmeijer K, Bogale A, Davidson RN (2008) Concordant HIV infection and visceral leishmaniasis in Ethiopia: the influence of antiretroviral treatment and other factors on outcome. *Clin Infect Dis* 46: 1702–09.
- Thakur CP, Kumar M, Kumar P, Mishra BN, Pandey AK (1988) Rationalisation of regimens of treatment of kala-azar with sodium stibogluconate in India: a randomised study. *BMJ* 296: 1557–61.
- Thakur CP, Kumar M, Pandey AK (1991) Evaluation of efficacy of longer durations of therapy of fresh cases of kala-azar with sodium stibogluconate. *Indian J Med Res* 93: 103–10.
- Thakur CP, Kumar M, Singh SK, et al. (1984) Comparison of regimens of treatment with sodium stibogluconate in kala-azar. *BMJ* 288: 895–97.
- Vanlerberghe V, Diap G, Guerin PJ, et al. (2007) Drug policy for visceral leishmaniasis: a cost-effectiveness analysis. *Trop Med Int Health* 12: 274–83.
- Wakabi W (2007) Hope for improved leishmaniasis treatment in Africa. *Lancet Infect Dis* 7: 638.
- White NJ (2004) Antimalarial drug resistance. *J Clin Invest* 113: 1084–92.
- WHO Special Programme for Research and Training in Tropical Diseases (2004) *Countries of the south-east Asia region plan to eliminate kala azar*. TDR News, page <http://www.who.int/tdr/publications/tdrnews/pdf/TDRnews-issue-73.pdf>. Accessed on: Jan 18, 2010.
- WHO. *Report of the fifth consultative meeting on Leishmania/HIV coinfection*. Addis Ababa, Ethiopia, 20–22 March 2007. http://www.who.int/leishmaniasis/resources/Leishmaniasis_hiv_coinfection5.pdf. Accessed on: Jan 20, 2010.



CHAPTER 8

Cost-effectiveness analysis of combination therapies for visceral leishmaniasis in the Indian subcontinent

Meheus F
Balasegaram F
Olliaro F
Sundar S
Rijal R
Abul Faiz M
Boelaert M

ABSTRACT

Background

Visceral leishmaniasis is a systemic parasitic disease that is fatal unless treated. We assessed the cost and cost-effectiveness of alternative strategies for the treatment of visceral leishmaniasis in the Indian subcontinent. In particular we examined whether combination therapies are a cost-effective alternative compared to monotherapies.

Methods and findings

We assessed the cost-effectiveness of all possible mono- and combination therapies for the treatment of visceral leishmaniasis in the Indian subcontinent (India, Nepal and Bangladesh) from a societal perspective using a decision analytical model based on a decision tree. Primary data collected in each country was combined with data from the literature and an expert poll (Delphi method). The cost per patient treated and average and incremental cost-effectiveness ratios expressed as cost per death averted were calculated. Extensive sensitivity analysis was done to evaluate the robustness of our estimations and conclusions. With a cost of US\$92 per death averted, the combination miltefosine-paromomycin was the most cost-effective treatment strategy. The next best alternative was a combination of liposomal amphotericin B with paromomycin with an incremental cost-effectiveness of \$652 per death averted. All other strategies were dominated with the exception of a single dose of 10mg per kg of liposomal amphotericin B. While strategies based on liposomal amphotericin B (AmBisome) were found to be the most effective, its current drug cost of US\$20 per vial resulted in a higher average cost-effectiveness. Sensitivity analysis showed the conclusion to be robust to variations in the input parameters over their plausible range.

Conclusions

Combination treatments are a cost-effective alternative to current monotherapy for VL. Given their expected impact on the emergence of drug resistance, a switch to combination therapy should be considered once final results from clinical trials are available.

INTRODUCTION

Despite their toxicity, pentavalent antimonials are still widely used as first line treatment for visceral leishmaniasis (VL) except in the Indian subcontinent where emerging drug resistance in Bihar State in India (Sundar *et al.* 2000b) and Nepal (Rijal *et al.* 2003) required a change in drug policy. Current therapeutic options include amphotericin B deoxycholate (AmB), liposomal amphotericin B (L-AmB), miltefosine (MF) and paromomycin (PM). The VL elimination initiative launched in 2005 by the governments of India, Nepal and Bangladesh adopted miltefosine as the first line treatment (Sundar *et al.* 2008; WHO 2005). More recently paromomycin was registered in India as a first line regimen for VL (Sundar *et al.* 2007). However, parasite resistance to MF and PM can be induced experimentally (Seifert *et al.* 2003) and is expected to emerge naturally if optimal adherence cannot be ensured (Bryceson 2001). The World Health Organization has recommended the use of liposomal amphotericin B (L-AmB) based on high efficacy and safety (Bern *et al.* 2006). While the development of resistance has not yet been demonstrated for AmB and L-AmB, practicalities (requirements for cold chain and intravenous perfusion) and the high drug cost have so far delayed its adoption as first line treatment. As there are no new compounds for VL expected to come to the market in the near future, policies that delay or prevent the emergence of resistance to the currently available drugs are therefore required. A possible strategy that has been successfully used for malaria and tuberculosis is the use of combination therapies (van Griensven *et al.* 2010). Combination therapies may also increase tolerability, reduce treatment duration and possibly (direct and indirect) costs.

Phase III clinical trials of combination therapies for VL are currently underway testing the efficacy and safety of several combinations and results are expected in 2010 (Clinicaltrial.gov, Identifier: NCT00696969; for more information see <http://clinicaltrials.gov/>). Choices in VL drug policy should be based on efficacy, safety as well as the cost of treatment, the process of patient management and the factors influencing treatment effectiveness, such as adherence. The latter factor is particularly important as some regimens (e.g. injectables) are likely to lead to higher compliance than others.

The objective of the present study was to assess the cost-effectiveness of various treatment options for VL, and in particular to evaluate whether combination therapies are a cost-effective alternative to monotherapy.

METHODS

Description of alternatives

We considered 10 alternative treatment strategies: (1) all monotherapies that are either already implemented or under consideration and (2) combination therapies currently included in a phase III clinical trial (See table 1). AmB has infusion-related and delayed toxicities (e.g.nephrotoxicity) (Sundar *et al.* 2004) and requires prolonged parenteral administration and hospitalisation. MF has the advantages of an oral drug but causes serious adverse events in 2–3% of patients (Bhattacharya *et al.* 2007) , has a long half-life and is possibly teratogenic. It can thus not be used in pregnant women and women in child-bearing age should accept contraception over the treatment period and up to two months after (Sundar *et al.* 2004a). PM seems a safer option - though phase IV results are still pending - and relatively cheap, but requires intramuscular injections. L-AmB is highly efficacious (>90%) even in a single dose of 5–10 mg/kg in India (Sunder *et al.* 2002b; Sunder *et al.* 2003b; Sundar *et al.* 2010) and is safe, but it is expensive despite a preferential price offered by the manufacturer to the public sector and requires an efficient cold chain. All of the other more “affordable” monotherapies listed above (MF, PM, AmB) require prolonged treatment which is problematic in very poor population groups that are dependent on daily labour and pay much out-of-pocket.

Table 1: Overview of treatment strategies included in decision analysis model.

| Strategy | Drug | Length of treatment (days) |
|----------|---------------------------------------------------------|----------------------------|
| A | L-AmB (5MK) + Miltefosine (50/100 MD) | 8 |
| B | L-AmB (5MK) + Paromomycin sulphate (15 MKD) | 11 |
| C | Miltefosine (50/100 MD) + Paromomycin sulphate (15 MKD) | 10 |
| D | SSG (20 MKD) + Paromomycin sulphate (15 MKD) | 17 |
| E | Miltefosine (50/100 MD) | 28 |
| F | Paromomycin sulphate (15 MKD) | 21 |
| G | Amphotericine B deoxycholate (1 MK eod) | 30 |
| H | L-AmB10 (10 MK) | 1 |
| I | L-AmB20 (5 MKD) | 4 |
| J | Sodium Stibogluconate (20 MKD) | 30 |

L-AmB : Liposomal Amphotericine B

MK = mg/kg single dose; MD = mg per day; MKD = mg/kg body weight per day

Miltefosine is given at 50 mg/ day if body weight is <25 or 100 mg if body weight ≥25 kg per day

Decision model

A decision tree model, depicted in figure 1, was developed using TreeAge Pro Suite v2009 (TreeAge Software Inc., Williamstown, MA, USA) to determine the outcome of a single confirmed VL patient receiving first-line treatment at a primary health care facility. The outcome was expressed in terms of number of deaths averted and we assumed a case-fatality rate of 100% in the absence of treatment. For each treatment strategy, the patient either adheres or does not. Those adhering are either cured or experience treatment failure. Patients not adhering to treatment were considered lost to follow-up and we assigned a value of 0 deaths averted. For strategies combining L-AmB with MF or PM, we assigned a value of 0.91 in case of non-adherence since patients will have received a single dose of 5mg/kg of L-AmB (with 91% cure rate) (Sundar *et al.* 2008b; Sundar *et al.* 2001) on the first day before they are lost to follow-up. Since MF is contraindicated in pregnant and breastfeeding women or women in child bearing age because of its potential teratogenic effect, the path for strategies including MF is different from those without MF. MF can only be given if the patient accepts the use of contraceptive measures during treatment and up to two months after completion of treatment. This is captured in the model by including an additional probability representing the contraceptive prevalence in the community. We calculated the cost per case treated and the average and incremental cost-effectiveness ratios (ICERs) expressed as the cost per death averted. The ICER represents the additional cost to gain an additional unit of effectiveness (i.e. one additional death averted) and is calculated by dividing the incremental cost of a given strategy by its incremental effectiveness compared to the previous not dominated strategy.

Furthermore we assumed in the baseline analysis the patient to be hospitalized for at least one day for all strategies; for treatment with AmB the patient is hospitalized for the entire duration of treatment (30 days) as this drug needs to be given under close supervision. Strategies with PM and SSG are provided on an outpatient basis whereby the patient visits the health facility daily to receive the intramuscular injection. In the case of treatment with MF, the patient visits the health facility weekly to receive a 1-week supply of the drug. We also assumed patients to undergo weekly routine laboratory investigations (blood count, liver and renal function tests), a pregnancy test for women in child-bearing age and an HIV test.




Figure 1: Root decision tree with different pathways depending on whether miltefosine is included in the strategy.

Probabilities

The probabilities used in the model are shown in table 2. These consist of a most plausible value used in the baseline analysis and the range used in the sensitivity analysis. We used anthropometric data from a sample of 1496 patients attending a dedicated VL treatment centre in Muzaffarpur, Bihar (India) (Olliaro & Sundar 2009) to derive probabilities on patient characteristics (age, weight and sex). Other values were obtained from expert opinion and published literature. To derive the probabilities related to efficacy and compliance for therapies that are still in clinical trials, we consulted a group of VL experts in an adapted Delphi process to reach consensus after two consultation rounds. In the first round seven clinical experts were presented with a survey asking for efficacy and compliance values for all treatments. Subsequently results from this round were summarized and presented to the experts to revise their earlier answers (round 2). For estimates derived from the literature, we used data from clinical trials using the pooled estimate in the baseline analysis and used the minimum and maximum for sensitivity analysis. All estimates represent definite cure rates defined as the absence of VL at 6 month follow-up; failure, relapse and fatal toxicity are included in the estimates. While minor side effects, such as diarrhoea and vomiting may occur, we did not consider these in the model since they do not hamper the completion of treatment.

We assumed treatments of short duration (strategies A, B, C, H and I) to result in high compliance. Similarly, treatment with AmB was assumed to lead to high compliance since treatment is provided on an inpatient basis. On the other hand, patients receiving MF for 28 days receive a 1-week supply of drug at a time for self-administration and compliance is anticipated to be lower than for the other strategies, consistent with findings from a miltefosine phase IV study by Bhattacharya et al (2007) (Bhattacharya *et al.* 2007) where the final cure rate was 82% on intent-to-treat analysis due to the high losses to follow-up.

Costs

Table 3 summarizes cost estimates presented in 2008 US dollars (US\$). Costs were obtained from primary data collected in 2008 using an ingredients based approach (i.e. collecting information on quantities and prices) and supplemented by data from the literature. We adopted a societal perspective including both provider and patient costs. These costs consist of direct medical costs (e.g. antileishmanial drugs, administration kits (intravenous sets, syringes and needles), laboratory investigations and the cost of hospitalization and outpatient care); direct non-medical costs (transportation to/from the health facility) and indirect costs representing the loss of income to the patient. The cost of drugs was obtained from WHO, Médecins sans Frontières (MSF) and the Institute of One World Health (iOWH). The cost of L-AmB was US\$ 20 per 50 mg vial (AmBisome, Gilead, USA), MF (Impavido previously Zentaris, Germany, now

Table 2: Model parameters (a)

| Variable | Likeliest (base) | Minimum | Maximum | Source |
|---------------------------------------------------|------------------|---------|---------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Demographic parameters of sample (%) ^b | | | | |
| Women in the sample | 39 | 30 | 50 | Olliaro & Sundar 2009; Thakur <i>et al.</i> 2000b |
| Women of childbearing age (15-49 yrs) of all VL | 17 | 10 | 35 | Olliaro & Sundar 2009; Thakur <i>et al.</i> 2000b |
| Patients weighing less than 25 kg | 41 | 20 | 60 | Olliaro & Sundar 2009; Thakur <i>et al.</i> 2000b |
| Children (0-14 years) | 51 | 25 | 75 | Olliaro & Sundar 2009; Thakur <i>et al.</i> 2000b |
| Adults (15-80 years) | 49 | 25 | 75 | Olliaro & Sundar 2009; Thakur <i>et al.</i> 2000b |
| Drug efficacy (%) | | | | |
| L-AmB + MF | 95 | 90 | 99 | a; Sundar <i>et al.</i> 2008b |
| L-AmB + PM | 95 | 90 | 99 | a |
| MF + PM | 95 | 91 | 99 | a |
| SSG + PM | 90 | 85 | 98 | a; Thakur <i>et al.</i> 2000b; Thakur <i>et al.</i> 1992 |
| MF | 94 | 82 | 94 | Sundar <i>et al.</i> 2002a; Thakur <i>et al.</i> 2000b; Bhattacharya <i>et al.</i> 2004; Sundar <i>et al.</i> 2003a; Sundar <i>et al.</i> 2000a |
| PM | 94 | 89 | 95 | a; Sundar <i>et al.</i> 2007; Jha <i>et al.</i> 1998; Sundar <i>et al.</i> 2009; Thakur <i>et al.</i> 2000a |
| AmB | 97 | 96 | 99 | Sundar <i>et al.</i> 2007; Sundar <i>et al.</i> 2004; Sundar <i>et al.</i> 2002a; Thakur <i>et al.</i> 1999 |
| L-AmB10 | 95 | 93 | 98 | a; Sundar <i>et al.</i> 2010 |
| L-AmB20 | 95 | 93 | 99 | a; Thakur 2001 |
| SSG | 70 | 35 | 93 | Sundar <i>et al.</i> 2000b; Rijal <i>et al.</i> 2003; Thakur <i>et al.</i> 2000b; Jha <i>et al.</i> 1998; Thakur <i>et al.</i> 2000a; Thakur & Narayan 2004; Thakur <i>et al.</i> 2008; Thakur <i>et al.</i> 1998; Sundar <i>et al.</i> 1997 |

Table 2: Model parameters (b)

| Variable | Likeliest (base) | Minimum | Maximum | Source |
|------------------------------|------------------|---------|---------|------------------------------------|
| Compliance to treatment (%) | | | | |
| L-AmB + MF | 95 | 80 | 97 | a |
| L-AmB + PM | 95 | 85 | 97 | a |
| MF + PM | 95 | 80 | 97 | a |
| SSG + PM | 83 | 75 | 90 | a |
| MF | 80 | 60 | 90 | a; Bhattacharya <i>et al.</i> 2007 |
| PM | 85 | 75 | 90 | a |
| AmB | 90 | 80 | 90 | a |
| L-AmB10 | 100 | - | - | a |
| L-AmB20 | 98 | 95 | 100 | a |
| SSG | 75 | 60 | 90 | a |
| Contraceptive prevalence (%) | 55 | 30 | 70 | a; WHO-WHOSISc |

^a Estimates obtained from expert opinion (Delphi method).
^b Baseline values were varied ±50% in sensitivity analysis
^c Average of figures reported for Bangladesh 58.1% (2004); India 56.3% (2006); Nepal 48.0% (2006)

Paladin, Canada) US\$ 1.41 per 100 mg capsule (or US\$ 79 per blister of 56 capsules; the market price of US\$ 2.68 per capsule was used as the maximum value in the range), PM US\$ 0.71 per 1000 mg ampoule (Gland Pharma Ltd, India), AmB US\$ 1.90 per mg vial (Combinopharm, Spain) and SSG US\$ 8.25 per vial (Albert David, India). The average drug cost per patient for each strategy was estimated using the anthropometric database. The baseline cost of laboratory investigations includes the cost of equipment, supplies, reagents, the technician’s time and indirect laboratory costs (i.e. overhead costs obtained through step-down costing) and is an average cost calculated at a VL treatment centre in India (Muzaffarpur) and a health facility in Nepal (Dharan). The range consists of prices charged to patients at public health facilities and private laboratories. The unit cost per inpatient bed-day and outpatient visit was estimated at a charitable clinic in India (Meheus *et al.* 2006). The maximum value used in the range was derived from WHO-CHOICE estimates for the South Asian region (Mulligan *et al.* 2003). Average income was estimated with the human capital approach and was estimated at US\$ 1.48 per day (Meheus *et al.* 2006). We assumed that the patient was not able to work for the full duration of treatment. Indirect costs were varied from 0 (i.e. excluding indirect costs) to twice the baseline value in sensitivity analysis. Costs related to diagnosis of VL were not included in the analysis since these are the same for all strategies.

Table 3: Cost estimates of each treatment strategy per patient treated (US\$ 2008).

| Strategy | Drug cost | Other direct medical ¹ | Non-medical & indirect | Total cost ² |
|----------|-----------|-----------------------------------|------------------------|-------------------------|
| L-AmB+MF | 95.7 | 14.8 | 12.8 | 123.4 |
| L-AmB+PM | 87.1 | 20.5 | 25.3 | 132.9 |
| MF+PM | 29.5 | 19.5 | 23.8 | 72.9 |
| SSG+PM | 45.1 | 29.9 | 43.6 | 118.6 |
| MF | 62.8 | 22.0 | 45.4 | 130.2 |
| PM | 14.9 | 30.6 | 51.1 | 96.6 |
| AmB | 20.9 | 131.6 | 45.4 | 197.9 |
| L-AmB10 | 140.0 | 11.0 | 2.5 | 153.4 |
| L-AmB20 | 280.0 | 24.7 | 6.9 | 311.6 |
| SSG | 57.8 | 40.7 | 73.4 | 171.8 |

¹ Includes costs of contraceptives, administration (intravenous kits, solutions, syringes), laboratory investigations. It also includes the cost per inpatient bed-day and outpatient visit obtained.

² Total costs of strategies with MF in this table do not include cost of AmB given to women in childbearing age that refuse to take contraceptives and are therefore different from total costs mentioned in table 4.

All costs were adjusted to the 2008 national currency of each country using the consumer price index and converted to US dollars using the exchange-rate prevailing at that time.

Sensitivity analysis

To examine the uncertainty around variables and how these affect the outcome and conclusions of our study, we conducted a series of one-way and two-way sensitivity analyses. Since values for drug efficacy and compliance were largely based on expert opinion we varied the values of these variables over the plausible range specified in table 2. On the cost side, we examined the impact of changing drug prices. While the price of most drugs, such as PM or AmB is unlikely to change much in the future given their low cost, there is uncertainty with regard to the pricing of MF and L-AmB. MF was recently acquired by Paladin Labs Inc., Canada from Zentaris, and it is at the time of writing unclear if the current negotiated differential prices will be maintained. For L-AmB, despite substantial price reductions, the cost per vial remains high and there may be room for further price reductions. To test the robustness of our results we (i) varied each drug cost separately; (ii) conducted a threshold analysis to determine at what price level strategies with L-AmB become the most cost-effective; and (iii) conducted a two-way sensitivity analysis of the price of MF and L-AmB. We also varied the unit cost per inpatient bed-day and outpatient visit. Finally, we examined the impact of indirect costs. In

the baseline analysis we assumed the patient would not be able to work for the full duration of treatment. But in practice, with effective treatment, patients may already feel better after a week of treatment and resume their activities. The indirect cost of strategies with longer treatment duration could therefore be overestimated. In addition the inclusion of indirect costs is a controversial issue, mainly due to the valuation method (Tan Torres *et al.* 2003). We therefore looked at the impact of indirect costs by (i) limiting productivity losses to a week, and (ii) excluding indirect costs from estimations.

RESULTS

Baseline analysis

Table 4 shows the expected cost and effectiveness for each treatment strategy using baseline values. Strategies were ranked in ascending order of costs. Strategies based on treatment with L-AmB, either used as a single agent or in combination, were found to be more effective compared to other strategies with a single dose of 10mg/kg of L-AmB (strategy H) being the most effective and averting 95% of deaths. This high effectiveness of strategies with L-AmB is explained by the combination of high drug efficacy and a short treatment duration resulting in high expected compliance to treatment. After strategies with L-AmB, the next best alternative is the co-administration of MF and PM (strategy C) averting 90% of deaths. Monotherapies with either SSG (strategy J) or MF (strategy E) had the lowest effectiveness. For SSG this is due to the low efficacy of the drug and for MF due to the low expected treatment compliance.

Table 4: Results in the baseline analysis.

| Strategy | Cost (C) | Incremental Cost* | Effectiveness (E) | Incremental Effectiveness* | C/E | Incremental C/E (ICER)** |
|------------|----------|-------------------|-------------------|----------------------------|-----|--------------------------|
| MF + PM | 82.5 | | 0.900 | | 92 | |
| PM | 96.6 | 14.1 | 0.799 | -0.101 | 121 | (Dominated) |
| SSG + PM | 118.6 | 36.1 | 0.747 | -0.153 | 159 | (Dominated) |
| L-AmB + MF | 129.1 | 46.6 | 0.942 | 0.042 | 137 | 1123** |
| L-AmB + PM | 132.9 | 3.8 | 0.948 | 0.006 | 140 | 652 |
| MF | 135.4 | 2.5 | 0.761 | -0.186 | 178 | (Dominated) |
| L-AmB 10 | 153.4 | 20.6 | 0.950 | 0.002 | 162 | 8224 |
| SSG | 171.8 | 18.4 | 0.525 | -0.425 | 327 | (Dominated) |
| AmB | 197.9 | 44.5 | 0.873 | -0.077 | 227 | (Dominated) |
| L-AmB 20 | 311.6 | 158.2 | 0.949 | -0.001 | 328 | (Dominated) |

The least costly treatment is the co-administration of MF with PM (strategy C) with a cost of \$72.9 per patient treated. The cost per patient treated for the other strategies varied from \$96.6 (strategy F) to \$311.6 (strategy I). The breakdown of costs for each strategy is shown in table 3. The drug cost as a proportion of total costs is the highest for strategies including L-AmB. The price of a 50mg vial of L-AmB at the time of this analysis was \$ 20 and is the most expensive VL drug. Obviously, the higher the dosage, the more expensive the treatment. For example the drug cost of strategy I (20mg/kg of L-AmB for 4 days) is \$280. The highest “other” direct medical costs were found for strategies requiring prolonged treatment, with treatment on an inpatient basis (strategy G) being the most expensive. Similarly, strategies with long treatment duration and/or requiring many visits to the health facility for administration of the drug have the highest indirect cost.

Cost-effectiveness results are illustrated in figure 2 and reported in table 4. All treatment strategies on the left of the line are dominated by strategies C, B and H because they are equally or less effective, and either cost more (strong dominance) or have an incremental cost-effectiveness ratio that is higher than the next more effective strategy (extended dominance). The incremental cost, incremental effectiveness, cost-effectiveness ratio (CER) and incremental CER without the dominated strategies are reported in table 5.




Figure 2: Cost-effectiveness ratio (US\$/death averted) of 10 treatment strategies for visceral leishmaniasis.

Line CBH shows dominance. All strategies left of this line are dominated by C, B and H, meaning they are equally or less effective and more costly.

Table 5: Baseline results without dominated options (simple or extended).

| Strategy | Cost (C) | Incremental Cost* | Effectiveness (E) | Incremental Effectiveness* | C/E | Incremental C/E (ICER)** |
|----------|----------|-------------------|-------------------|----------------------------|-----|--------------------------|
| MF+PM | 81.9 | | 0.900 | | 91 | |
| L-AmB+PM | 132.9 | 51.0 | 0.948 | 0.047 | 140 | 1079 |
| L-AmB10 | 153.4 | 20.6 | 0.950 | 0.002 | 162 | 8224 |

The most cost-effective strategy appears to be strategy C whereby MF and PM are co-administered. Compared with this strategy, the next most cost-effective strategy is the combination of L-AmB with PM (strategy B), followed by a single dose of 10mg/kg of L-AmB (strategy H). While L-AmB combined with MF (strategy A) is more effective than strategy C, it is also more costly and has a higher (incremental) cost-effectiveness ratio than the next best alternative (i.e. strategy B). In other words the additional cost per death averted is lower for strategy B than strategy A.

Sensitivity analysis

The study findings were robust to most changes in the input variables. Varying the values of the drug efficacy and compliance over their plausible range did not affect the ranking of strategies. A sensitivity analysis assuming that all non-adherent patients would effectively be cured did not change the ranking of strategies either. With regard to costs, while varying the price of MF did not alter results, the cost-effectiveness results were sensitive to a change in the price of L-AmB. If the price of a vial is decreased by more than 51% to less than \$ 9.8, then strategy H becomes the most cost-effective strategy. The relationship between the price of MF and the price of L-AmB and their impact on the ranking of strategies according to their cost-effectiveness, keeping all other variables at their baseline values is shown in figure 3. If MF is purchased at market price (\$ 2.68 per capsule), the price per vial of L-AmB would need to decrease to below \$12.5 for strategy H to become the most cost-effective strategy. Varying the assumptions regarding indirect cost did not change conclusions.




Figure 3: Two-way sensitivity analysis on price of AmBisome and miltefosine.

DISCUSSION

The current first line regimen in the Indian subcontinent is MF for 28 days. There are concerns however that the uncontrolled provision of the drug may increase the likelihood of development of parasite resistance (Sundar & Murray 2005). Even when monitored, patient compliance is not optimal (Bhattacharya *et al.* 2007; Sundar & Murray 2005) and the risk remains that women in childbearing age receiving MF do not (or only partially) take contraceptives. Our analysis shows that combination therapies for the treatment of VL are a cost-effective alternative to the current strategy in the Indian subcontinent; this finding may be of interest to control programmes regarding the cost-effectiveness of the currently recommended option. The co-administration of MF with PM for 10 days seems to be the most cost-effective option because of the combined effect of low cost, especially drug cost, and high effectiveness. Also, one would expect that the parenteral intramuscular injection of PM by health workers ensures that patients also take the oral MF as it would be directly observed. With the short treatment duration this is likely to result in high patient compliance, increasing the overall effectiveness of the strategy. Although strategies with L-AmB were the most effective, the high drug cost results in a higher average cost-effectiveness. The next best alternative compared

to the combination MF/PM was a combination of L-AmB with PM with an incremental cost-effectiveness of \$652 per death averted. All other strategies, with the exception of a single dose of 10mg per kg of liposomal amphotericin B were dominated. The relatively poor effectiveness for MF monotherapy in our model is linked to the estimated low adherence when using self-administration with 1-weekly drug supplies. However, alternative drug delivery strategies for MF monotherapy are possible. A strategy where intake of MF would be directly observed would result in significantly higher effectiveness, although at higher direct and indirect costs. When we ran a sensitivity analysis with MF compliance put at 100%, this did not change the ranking of strategies or conclusions.

This study is the most comprehensive cost-effectiveness analysis of alternative strategies for the treatment of VL for the Indian subcontinent to date. We used a simple decision analytical model to compare from a societal perspective the cost and outcome of all possible treatment strategies identified through consultation with experts (Delphi method). The demographic probabilities used in the model, as well as the calculation of the average drug cost was based on real patient data on sex, age and weight obtained from a charitable medical facility in India (Olliaro and Sundar 2009), instead of calculating the drug cost of an “average” 35kg patient as done in other studies. In addition, all cost data in the baseline analysis were based on primary data collected from various sites in Nepal and India. Extensive sensitivity analysis was also done to evaluate the robustness of our estimations and conclusions. The analysis has several limitations. *First* while the use of the anthropometric data can be a strength, the VL treatment centre in Muzaffarpur (Bihar) might not be entirely representative for all VL cases, especially with regard to the male to female ratio. Reassuringly, in a larger series of 4170 patients from two locations in Bihar and 1311 from Nepal the male to female ratio was similar (57:43, Olliaro *et al.*, manuscript submitted). Although various studies have reported a higher proportion of male patients to be affected by kala-azar (Singh *et al.* 2006; Thakur 2000) there may be under-reporting of kala-azar in women (Bora 1999) due to sex-selective treatment seeking whereby “fewer women may seek treatment because of its expense” (Thakur 2000). A M/F ratio in the VL population closer to unity could lower the effectiveness of strategies including MF. *Second*, the drug efficacy estimates for the combination treatments, and the monotherapies with L-AmB were based on input from a Delphi survey of VL experts. The uncertainty surrounding these subjective estimates was minimized by including experts that were clinicians and/or involved in clinical trials of combination treatments and dose-finding studies. The uncertainty was also analysed extensively in sensitivity analysis. *Third* the effectiveness estimates are heavily influenced by the parameters of patient compliance to treatment. Experts assumed treatments with parenteral or intramuscular administration to lead to high compliance and oral treatment to result in lower patient compliance. Given the evidence from the international

literature for other diseases, and the limited information available for kala-azar (Bhattacharya *et al.* 2007; Sundar *et al.* 2009), these assumptions seem plausible. As more evidence becomes available from clinical trials (especially Phase IV and other operational studies) and future studies assessing patient compliance, we will update the input parameters and ranges from our model. *Finally*, some cost variables were not included in the analysis. L-AmB requires a cold chain. Because it was difficult to quantify the cost of the cold chain, we did not include it in our calculations and the cost per patient treated and cost per death averted may therefore be an underestimation. There is, moreover, also a substantial risk of breakdowns in the cold chain system, which may impact on the efficacy of the drug. Governments may want to adapt the drug policy choice to the technology constraints in each level of the health system.

Few other studies have investigated the cost-effectiveness of VL treatment strategies. Vanlerberghe *et al.* (2007) (Vanlerberghe *et al.* 2007) compared various monotherapies from a health service perspective (not including paromomycin) and found a strategy with miltefosine to be the most cost-effective with US\$328 per death averted. Although the study uses a similar decision tree model and sensitivity analysis to account for uncertainty in the input parameters, the results from this study cannot be compared with ours. The model by Vanlerberghe *et al.* starts with a clinical suspect going through diagnosis and then treatment. Treatment effectiveness is therefore defined by probabilities other than those directly related to treatment such as the prior probability of disease, and the sensitivity and specificity of the diagnostic test. Patient compliance was not modelled either. A more recent study by Olliaro *et al.* (2009) (Olliaro *et al.* 2009) compared various monotherapies and a combination of L-AmB with MF with different total dosages for MF from a health systems perspective. Similar to our findings, Olliaro *et al.* show that the combination L-AmB+MF (for 8 days) with a cost of \$124–160 per death averted is more cost-effective than most monotherapies (the exception being PM delivered in an outpatient setting and a 5mg/kg single dose formulation of L-AmB). However this study did not include indirect costs (i.e. productivity losses) underestimating the effect of strategies with a short treatment duration that are beneficial to the patient and household.

Our results highlight that several possible therapeutic options may exist for the South Asian context - especially in light of the ongoing VL elimination campaign in the Indian subcontinent - but combination regimens are efficient options compared to monotherapy. The analysis should be repeated in other VL-endemic areas such as East Africa and Brazil where efficacy outcomes, treatment regimens, direct and indirect costs may differ considerably. Critical elements of importance to national and international policy makers are the cost of drugs, the level of out-of-pocket expenditures by VL patients and compliance to treatment. An obstacle to the introduction of strategies with L-AmB in national control programmes is the cost of the drug.

Despite substantial reductions in the price of AmBisome over the past years (more recently to \$18 for a 50mg vial), the threshold analysis showed this to be not enough to make strategies with L-AmB a cost-effective alternative. In addition, the capacity of VL patients and their family to cover the costs of treatment is very limited. VL is a disease that affects the poorest of the poor Boelaert *et al.* 2009 and places a considerable economic burden on households (Meheus *et al.* 2006; Rijal *et al.* 2006; Sharma *et al.* 2006). Especially in India and Bangladesh, the combination of frequent drug shortages and poor quality of care in public health facilities pushes many patients to buy drugs from private pharmacies or to seek care in the private sector. Unless the government or a donation programme covers the cost of drugs, strategies including expensive drugs such as L-AmB will be a barrier to patients and reduce access to appropriate and effective care resulting in increased mortality.

REFERENCES

Bern C, Adler-Moore J, Berenguer J, et al. (2006) Liposomal amphotericin B for the treatment of visceral leishmaniasis. *Clin Infect Dis* 43: 917–24.

Bhattacharya SK, Jha TK, Sundar S, et al. (2004) Efficacy and tolerability of miltefosine for childhood visceral leishmaniasis in India. *Clin Infect Dis* 38: 217–21.

Bhattacharya SK, Sinha PK, Sundar S, et al. (2007) Phase 4 trial of miltefosine for the treatment of Indian visceral leishmaniasis. *J Infect Dis* 196: 591–98.

Boelaert M, Meheus F, Sanchez A, et al. (2009) The poorest of the poor: a poverty appraisal of households affected by visceral leishmaniasis in Bihar, India. *Trop Med Int Health* 14: 639–44.

Bora D (1999) Epidemiology of visceral leishmaniasis in India. *Natl Med J India* 12: 62–68.

Bryceson A (2001) A policy for leishmaniasis with respect to the prevention and control of drug resistance. *Trop Med Int Health* 6: 928–34.

Jha TK, Olliaro P, Thakur CP, et al. (1998) Randomised controlled trial of aminosidine (paromomycin) v sodium stibogluconate for treating visceral leishmaniasis in North Bihar, India. *BMJ* 316: 1200–05.

Meheus F, Boelaert M, Baltussen R, Sundar S (2006) Costs of patient management of visceral leishmaniasis in Muzaffarpur, Bihar, India. *Trop Med Int Health* 11: 1715–24.

Mulligan JA, Fox-Rushby JA, Johns B, Mills A (2003) Unit Costs of Health Care Inputs in Low and Middle Income Regions. Disease Control Priorities Project (DCPP) Working Paper n°9.

Olliaro P, Darley S, Laxminarayan R, Sundar S (2009) Cost-effectiveness projections of single and combination therapies for visceral leishmaniasis in Bihar, India. *Trop Med Int Health* 14: 918–25.

Olliaro P, Sundar S (2009) Anthropometrically derived dosing and drug costing calculations for treating visceral leishmaniasis in Bihar, India. *Trop Med Int Health* 14: 88–92.

Rijal S, Chappuis F, Singh R, et al. (2003) Treatment of visceral leishmaniasis in south-eastern Nepal: decreasing efficacy of sodium stibogluconate and need for a policy to limit further decline. *Trans R Soc Trop Med Hyg* 97: 350–54.

Rijal S, Koirala S, Van der Stuyft P, Boelaert M (2006) The economic burden of visceral leishmaniasis for households in Nepal. *Trans R Soc Trop Med Hyg* 100: 838–41.

Seifert K, Matu S, Javier Perez-Victoria F, Castanys S, Gamarro F, Croft SL (2003) Characterisation of *Leishmania donovani* promastigotes resistant to hexadecylphosphocholine (miltefosine). *Int J Antimicrob Agents* 22: 380–87.

Sharma SD, Bern C, Varghese B, et al. (2006) The economic impact of visceral leishmaniasis on households in Bangladesh. *Trop Med Int Health* 11: 757–64.

Singh SP, Reddy DC, Rai M, Sundar S (2006) Serious underreporting of visceral leishmaniasis through passive case reporting in Bihar, India. *Trop Med Int Health* 11: 899–905.

Sundar S, Agrawal G, Rai M, Makharia MK, Murray HW (2001) Treatment of Indian visceral leishmaniasis with single or daily infusions of low dose liposomal amphotericin B: randomised trial. *BMJ* 323: 419–22.

Sundar S, Agrawal N, Arora R, Agarwal D, Rai M, Chakravarty J (2009) Short-course paromomycin treatment of visceral leishmaniasis in India: 14-day vs 21-day treatment. *Clin Infect Dis* 49: 914–18.

Sundar S, Chakravarty J, Agarwal D, Rai M, Murray HW (2010) Single-dose liposomal amphotericin B for visceral leishmaniasis in India. *N Engl J Med* 362: 504–12.

Sundar S, Jha TK, Thakur CP, et al. (2002a) Oral miltefosine for Indian visceral leishmaniasis. *N Engl J Med* 347: 1739–46.

Sundar S, Jha TK, Thakur CP, Mishra M, Singh VR, Buffels R (2002b) Low-dose liposomal amphotericin B in refractory Indian visceral leishmaniasis: a multicenter study. *Am J Trop Med Hyg* 66: 143–46.

Sundar S, Jha TK, Sindermann H, Junge K, Bachmann P, Berman J (2003a) Oral miltefosine treatment in children with mild to moderate Indian visceral leishmaniasis. *Pediatr Infect Dis J* 22: 434–38.

Sundar S, Jha TK, Thakur CP, Mishra M, Singh VP, Buffels R (2003b) Single-dose liposomal amphotericin B in the treatment of visceral leishmaniasis in India: a multicenter study. *Clin Infect Dis* 37: 800–04.

Sundar S, Jha TK, Thakur CP, Sinha PK, Bhattacharya SK (2007) Injectable paromomycin for Visceral leishmaniasis in India. *N Engl J Med* 356: 2571–81.

Sundar S, Makharia A, More DK, et al. (2000a) Short-course of oral miltefosine for treatment of visceral leishmaniasis. *Clin Infect Dis* 31: 1110–13.

Sundar S, Mehta H, Suresh AV, Singh SP, Rai M, Murray HW (2004) Amphotericin B treatment for Indian visceral leishmaniasis: conventional versus lipid formulations. *Clin Infect Dis* 38: 377–83.

Sundar S, Mondal D, Rijal S, et al. (2008a) Implementation research to support the initiative on the elimination of kala azar from Bangladesh, India and Nepal—the challenges for diagnosis and treatment. *Trop Med Int Health* 13: 2–5.

Sundar S, More DK, Singh MK, et al. (2000b) Failure of pentavalent antimony in visceral leishmaniasis in India: report from the center of the Indian epidemic. *Clin Infect Dis* 31: 1104–07.

Sundar S, Murray HW (2005) Availability of miltefosine for the treatment of kala-azar in India. *Bull World Health Organ* 83: 394–95.

Sundar S, Rai M, Chakravarty J, et al. (2008b) New treatment approach in Indian visceral leishmaniasis: single-dose liposomal amphotericin B followed by short-course oral miltefosine. *Clin Infect Dis* 47: 1000–06.


Sundar S, Singh VP, Sharma S, Makharia MK, Murray HW (1997) Response to interferon-gamma plus pentavalent antimony in Indian visceral leishmaniasis. *J Infect Dis* 176: 1117–19.

Tan Torres T, Baltussen R, Adam T, et al. (2003) *Making choices in health: WHO guide to cost-effectiveness analysis*. Geneva: World Health Organization.

Thakur CP (2000) Socio-economics of visceral leishmaniasis in Bihar (India). *Trans R Soc Trop Med Hyg* 94: 156–57.

Thakur CP (2001) A single high dose treatment of kala-azar with Ambisome (amphotericin B lipid complex): a pilot study. *Int J Antimicrob Agents* 17: 67–70.

- Thakur CP, Kanyok TP, Pandey AK, Sinha GP, Messick C, Olliaro P (2000a) Treatment of visceral leishmaniasis with injectable paromomycin (aminosidine). An open-label randomized phase-II clinical study. *Trans R Soc Trop Med Hyg* 94: 432–33.
- Thakur CP, Kanyok TP, Pandey AK, et al. (2000b) A prospective randomized, comparative, open-label trial of the safety and efficacy of paromomycin (aminosidine) plus sodium stibogluconate versus sodium stibogluconate alone for the treatment of visceral leishmaniasis. *Trans R Soc Trop Med Hyg* 94: 429–31.
- Thakur CP, Narayan S (2004) A comparative evaluation of amphotericin B and sodium antimony gluconate, as first-line drugs in the treatment of Indian visceral leishmaniasis. *Ann Trop Med Parasitol* 98: 129–38.
- Thakur CP, Olliaro P, Gothoskar S, et al. (1992) Treatment of visceral leishmaniasis (kala-azar) with aminosidine (= paromomycin)-antimonial combinations, a pilot study in Bihar, India. *Trans R Soc Trop Med Hyg* 86: 615–16.
- Thakur CP, Singh RK, Hassan SM, Kumar R, Narain S, Kumar A (1999) Amphotericin B deoxycholate treatment of visceral leishmaniasis with newer modes of administration and precautions: a study of 938 cases. *Trans R Soc Trop Med Hyg* 93: 319–23.
- Thakur CP, Sinha GP, Pandey AK, Kumar N, Kumar P, Hassan SM (1998) Do the diminishing efficacy and increasing toxicity of sodium stibogluconate in the treatment of visceral leishmaniasis in Bihar, India, justify its continued use as a first-line drug? An observational study of 80 cases. *Ann Trop Med Parasitol* 92: 561–69.
- Thakur CP, Thakur S, Narayan S, Sinha A (2008) Comparison of treatment regimens of kala-azar based on culture & sensitivity of amastigotes to sodium antimony gluconate. *Indian J Med Res* 127: 582–88.
- van Griensven J, Balasegaram M, Meheus F, Alvar J, Lynen L, Boelaert M (2010) Combination therapy for visceral leishmaniasis. *Lancet Infect Dis* 10: 184–94.
- Vanlerberghe V, Diap G, Guerin PJ, et al. (2007) Drug policy for visceral leishmaniasis: a cost-effectiveness analysis. *Trop Med Int Health* 12: 274–83.
- World Health Organization (2005) *Regional strategic framework for elimination of Kala-azar from the South-East Asia region (2005–2015)*. New Delhi: WHO Regional Office for South-East Asia.



CHAPTER 9

General discussion and Conclusion

The general objective of the thesis was to improve our understanding of the economic aspects of visceral leishmaniasis and to provide evidence for more rational decision making on diagnosis, case management and VL control in the Indian subcontinent and East Africa. To meet this objective, we developed four research questions: (i) What is the burden of visceral leishmaniasis? (ii) What is the health seeking behaviour and costs of VL from the household and provider perspective? (iii) What is the cost and cost-effectiveness of VL treatment strategies? and (iv) What are the policy recommendations that can be formulated on the basis of our findings?

In the following sections I will discuss the findings from the studies presented in this thesis in conjunction with the evolving scientific literature along these research questions, and discuss ongoing research and provide recommendations for future work.

THE BURDEN OF VISCERAL LEISHMANIASIS

One objective of the thesis was to improve our understanding about the burden of VL. The burden of VL can be thought of in terms of the morbidity, mortality and disability as well as the economic and social impact of the disease. In 2010 we published a scientific supplement on the burden of VL in South Asia (Meheus *et al.* 2010) in support of the VL elimination initiative in Nepal, India and Bangladesh with a collection of studies investigating the epidemiological and economic impact of the disease. To build further on the information that was provided in these studies, we highlighted the need for more reliable and in particular routinely collected data on VL incidence and mortality. Recently, Alvar *et al.* 2012 provided updated figures and estimated that between 20 and 40 thousand people die from VL each year. This makes VL one of the most severe NTDs with the highest mortality after schistosomiasis and hookworm infection (Hotez *et al.* 2006). However, unlike the other NTDs, the case fatality rate for untreated VL is 100% and death may occur within a matter of months if the affected individual is left untreated.

The disease burden is now often expressed in terms of disability adjusted life years (DALYs), an indicator that combines (premature) mortality, morbidity and disability in a single metric (Murray and Lopez 1994; WHO 1996). DALYs have become a key measure in the WHO Global Burden of Disease project to determine the disease burden of a wide range of health conditions and are frequently employed by international donors in priority setting and resource allocation. Leishmaniasis is estimated to cause the loss of about 2.1 million DALYs annually (WHO 2004; Mathers *et al.* 2007). However the use of DALYs in priority setting for neglected tropical diseases (NTDs), including leishmaniasis, has been heavily criticized (Lutumba *et al.* 2007; Bern *et al.* 2008; King & Bertino 2008; Reithinger 2008) and estimates are generally

considered too low (Engels & Savioli 2006). Besides the problems discussed in the introduction of the thesis of not having accurate data on mortality and morbidity that are needed for the DALY calculation, DALYs do not take account of the spatial clustering of VL. While at the global, national or sub-national level the burden of VL in terms of incidence, mortality and DALYs may be low, the burden at the community level can be very high. Due to the focal distribution of VL, incidence rates (and underreporting) vary greatly between countries (Mondal *et al.* 2009) as well as between districts and within districts as shown by Das *et al.* (2010). Moreover, VL cases tend to cluster within certain sections of the village at hamlet level and further at household level (Ahluwalia *et al.* 2003; Hasker *et al.* 2012). Incidence rates can be ten times greater at hamlet level (in the range of 1-2% per year) than the aggregate figures recorded at district level (in the range of 1-2 per 1000 per year). Because of the inadequate representation of the burden of disease of VL in terms of DALYs and since VL treatment is a life-saving intervention, we have chosen deaths averted as the unit of outcome to compare treatment alternatives in chapter 8. However, because DALYs are increasingly used in priority setting for health, a re-estimation of the DALYs for VL is necessary as has been done for other NTDs (Luz *et al.* 2009).

In chapter 3 we studied the socio-economic profile of households living in areas with high VL endemicity relative to the general population of Bihar, using data collected from a community intervention trial (i.e. the KALANET trial) and the third Indian National Family Health Survey (NFHS-3). With 83% of households in VL endemic areas belonging to the poorest two quintiles in Bihar, we showed that in areas where VL occurs, the socio-economic status of households is considerably lower than the rest of Bihar. Bihar, in turn, is also one of the poorest states in India. The study confirmed the general notion that leishmaniasis is strongly associated with poverty and affects the “poorest of the poor”. Numerous studies have shown that poverty increases the risk of VL infection or disease, mainly through poor living conditions (Bern *et al.* 2000; Ranjan *et al.* 2005; Schenkel *et al.* 2006; Saha *et al.* 2008; Bern *et al.* 2010; Singh *et al.* 2010a; Hasker *et al.* 2012; Pascual Martinez *et al.* 2012). Killick-Kenrick coined the term ‘paradise for sandflies’ to describe the housing conditions in endemic villages. Damp earthen floors, mud-plastered walls with cracks, proximity to water bodies, soil littered with organic matter, sleeping on the ground etc. are all linked to poverty as well as to increased exposure to peri-domestic sandflies. VL also seems to occur more frequently among scheduled castes in India (e.g. the Mushar caste in Bihar) (Singh *et al.* 2000; Sharma *et al.* 2004; Shah 2005; Hasker *et al.* 2010; Pascual Martinez *et al.* 2012), even after controlling for socio-economic status (Hasker *et al.* 2012). The Indian government rural housing scheme, Indira Awaas Yojana, that provides incentives to people living below the poverty line to construct permanent housing, may help in reducing VL incidence (Government of India 2012).

In summary, VL is one of the most severe NTDs with a 100% case fatality rate for untreated VL. The current estimations of the general burden of disease of VL using DALYs are too low; a re-estimation of DALYs for VL is therefore needed. This also includes revising the disability weight attributed to VL. However an important stumbling block to estimate the general burden of disease of VL remains the uncertainty around incidence and mortality data. We have also shown that VL is strongly associated with poverty and affects the “poorest of the poor”.

THE HEALTH SEEKING BEHAVIOUR AND COSTS OF VL PATIENT MANAGEMENT

In part II of the thesis we presented three chapters that analysed the economic burden of VL on the household (i.e. household perspective) in India (chapter 4), Nepal (chapter 5) and Sudan (chapter 6). These studies explored issues related to the health care seeking behaviour of households, direct and indirect costs, and mechanisms to cope with the cost of illness (research question #2). Chapters 4 (India) and 6 (Sudan) also present information on the cost of VL case management from the perspective of the health provider. I will first discuss the results from the provider’s perspective followed by a discussion on the economic burden of VL from the household perspective. The cost data presented below was converted to 2010 US dollars using the consumer price index of each respective country to facilitate comparisons between studies.

The aim of the analysis from the providers’ perspective was to estimate how much it costs to public (and not-for-profit) health facilities to provide VL treatment for one episode of VL. The evidence on the cost of VL patient management from the perspective of the health care provider is very limited. Only two studies have been conducted so far and were presented in this thesis in chapters 4 and 6 for India and Sudan respectively. Besides informing policy makers about the cost per case treated, the information provided by these studies has been used in economic evaluation studies to compare treatment alternatives for VL (see chapter 8), assess the value of a hypothetical VL vaccine (Lee *et al.* 2012) and can be used to estimate the resource requirements needed for VL control, as is commonly done for other infectious diseases such as HIV/AIDS (Vassall & Compernelle 2006).

In our studies we estimated the medical cost of VL care (drugs, diagnosis, medical supplies, laboratory investigations) and the cost of hospitalization separately by combining step-down cost accounting with an ingredients approach (i.e. bottom-up microcosting). These methods, described in detail in chapter 6, are frequently employed in countries where health information systems and health facility financial records are weak. The estimation in chapter 4 was relatively straightforward since the study was conducted at a non-governmental hospital

in Muzaffarpur in Bihar that provides care for VL only. All costs made by the hospital could therefore be directly allocated to VL patient management. In Sudan (chapter 6) estimations were more complex as VL care represented only a small proportion of overall activities in two out of the three hospitals that were included in the study. Despite the methodological challenges, we believe the cost data to be accurate since most of the data related to the medical costs of VL care were collected using microcosting whereby the actual resource consumption by patients was examined. Although time consuming, microcosting is generally considered to provide the most precise estimates of the costs of care (Drummond *et al.* 2005).

The study in India was conducted at a time when amphotericin B was provided at not-for-profit hospitals, while sodium stibogluconate (SSG) was still the first line treatment at public health facilities, despite the reports about unresponsiveness to the drug in the Indian subcontinent (Sundar *et al.* 2000; Rijal *et al.* 2003). In India, non-governmental health facilities, alongside the private sector, are common sources of treatment for VL patients (Mondal *et al.* 2009; Sundar *et al.* 2010). Although a different first line drug is now used, the daily cost of hospitalization, or unit cost per inpatient day, is likely to remain about the same since medical costs were excluded in its calculation. Our definition of unit cost was also similar to the WHO definition of a “hotel” unit cost (WHO-Choice 2012) and would now be equal to 4.6 US dollars in 2010 values. Obviously, the longer the treatment duration, the higher the cost of hospitalization will be (i.e. duration of treatment multiplied by the unit cost per inpatient day). The cost of hospitalization represented a third of the cost of VL patient management, with drugs and laboratory investigations (i.e. medical costs) each representing a third as well (VL drugs accounted for 71% of the drug costs). These medical costs were very specific to the type of VL drug used. With amphotericin B, patients were regularly submitted to laboratory investigations to test for toxicity and side-effects related to the drug, leading to a high cost for the hospital. Since not only the type of VL drug used but also the prices of these drugs may change over time, regular updates of the cost estimations are needed.

In Sudan, the cost of hospitalization was the largest cost driver in the hospitals included in the study with nearly 80% of the total cost of VL patient management. This was caused by the fact that patients were hospitalized for 30 days of treatment with SSG. The remaining 20% of the cost of VL patient management consisted mainly of the cost of SSG; few laboratory investigations were done. While the average unit cost per inpatient day across health facilities was 5.7 US dollars, we observed a large variation between facilities. Unit cost estimates are particularly sensitive to (inaccurate) activity data and there were some concerns regarding the activity statistics of the hospitals. Only crude information was available on patient throughput needed to calculate the cost per inpatient day. Although information on the number of admissions was

available, statistics on the length of stay by disease category was not centrally available and had to be derived from a sample of medical records.

In terms of the economic burden of VL to households (i.e. household perspective), the following conceptual framework can be derived from our studies:




Figure 9.1: Conceptual framework on the economic burden of VL to households

Our studies find that patients usually presented to a qualified health provider with a delay of one to two months. The type of provider first visited varied between countries. This was not unexpected given the important differences in the organization of the health care system between countries.

In India, where the private sector plays an important role in the provision of health care, our findings showed that patients most often visited either a private qualified or unqualified provider first; less than a third of patients that visited a private qualified provider were diagnosed with VL on their first visit. Several studies published in recent years confirmed our findings and showed that the health seeking behaviour of patients had not changed (Sundar *et al.* 2010; Sarnoff *et al.* 2010; Hasker *et al.* 2010) and private providers continued to be an important primary source of care for affected households.

In Nepal and Sudan the majority of patients had first consulted a public provider. In Nepal, our study showed some differences with previous studies conducted in the country that had shown that traditional providers were most commonly the patient's first choice of provider (Sharma *et al.* 2004; Adhikari and Maskay 2005). These studies were carried out nearly a decade ago; a knowledge attitudes and practices survey carried out in 2006/07 had shown knowledge of

VL to be reasonable in VL endemic areas in Nepal with patients mainly using the public sector, similar to our study (Mondal *et al.* 2009). In addition, the presence of the BP Koirala Institute of Health Sciences (BPKIHS) in our study area may have influenced the health seeking behaviour of households and referral practices of health providers. The BPKIHS is a tertiary level hospital situated in Sunsari district that has a widespread reputation in surrounding areas as a VL treatment and research centre.

In the case of Sudan there are not many qualified private practitioners in rural areas (WHO-EMRO 2006) and nearly all patients had visited a public provider, either a village health worker, primary health care centre or public hospital. We found that few patients in Sudan were diagnosed with VL on their first visit, also among those visiting a public hospital. While Gerstl *et al.* (2006) had shown that VL knowledge was limited among community health workers, our data seems to indicate that with the exception of VL dedicated treatment centres, the knowledge of VL is also low at public hospitals and primary health care centres. However our findings on health seeking behaviour in Sudan need to be interpreted with caution because investigating health seeking behaviour was not the primary objective of the study and only quantitative data was collected. Despite this limitation, it is clear that more training is needed among health workers at all levels in highly endemic areas to strengthen the early diagnosis of VL, especially since diagnosis and treatment of VL is the only control strategy currently implemented in Sudan.

At the time of the review presented in chapter 2, there had been seven studies that estimated the financial and/or economic costs of VL in India (Meheus *et al.* 2006 (chapter 4); Sarnoff *et al.* 2010; Sundar *et al.* 2010), Nepal (Sharma *et al.* 2004; Rijal *et al.* 2006; Adhikari *et al.* 2009) and Bangladesh (Sharma *et al.* 2006). There had been no published studies on the economic impact of VL in East Africa. Since then, two additional studies have been published (presented in this thesis), including one study for Sudan (chapter 6). So far, all studies estimated the economic impact of VL at the micro-economic level, i.e. at the level of the household. No studies have assessed the economic impact of VL at the macroeconomic level. Although the severity of disease is high it is unlikely that VL will have an impact on for example gross domestic product as has been shown for malaria or HIV/AIDS (Gallup & Sachs 2001; Dixon *et al.* 2002; Jefferis *et al.* 2008). The majority of VL affected patients are not economically active as primarily young people are affected, and those that are working are mainly employed in the informal sector or subsistence agriculture.

From the perspective of the household, there are two types of costs that can be associated with an episode of VL. Direct costs are (out-of-pocket) expenses incurred by households when seeking treatment. These costs can be broken down into medical (e.g. diagnosis, drugs or

laboratory investigations) or non-medical costs (e.g. food, transportation). Indirect costs were defined as the lost economic productivity due to VL illness to the patient and family members taking care of the patient. We used a uniform definition of costs across the three studies included in this thesis to ensure comparability, which was sometimes an issue with the studies identified in chapter 2.

The studies presented in section 2 (chapters 4-6) of the thesis showed that the economic burden of VL illness is considerable and in most cases catastrophic. We found that indirect costs were larger than direct costs in Nepal and India, while direct costs greatly exceeded indirect costs in Sudan. The duration of illness and the length of treatment were important determinants of costs in all three studies.

In terms of the direct costs, the proportions spent by households on medical and non-medical services varied by type of VL drug used and the patient management process in each country. Expenditure on VL drugs were the largest in India (chapter 4). While most medical services were provided free of charge to patients, VL drugs (at the time amphotericin B) had to be retrieved by the patient or a family member outside the hospital at a local private pharmacy and brought back to the hospital where they were administered intravenously by the medical personnel. In Nepal and Sudan, VL diagnosis and drugs were provided free of charge to patients at public health facilities. Similarly to Nepal, miltefosine is now increasingly being used in India (Banjara *et al.* 2012) as recommended by the VL elimination initiative and both NGO clinics and public health care facilities provide VL drugs and diagnostics for free. The provision of free VL drugs should reduce the economic burden of VL to households. Although two additional studies were published for India in 2010 (i.e. Sundar *et al.* 2010; Sarnoff *et al.* 2010), the data for these studies was collected only two years after our study, prior to the actual implementation of the VL elimination initiative and households still had to pay for the VL drugs in these studies.

In Nepal (chapter 5), the data for the study was collected in 2011 and we examined whether the VL control efforts by the government resulted in a decrease in household costs. The study was not experimental (e.g. pre-post test design) but descriptive and data was collected from patients that were treated for VL at public health care facilities and identified from the medical records of the District Public Health Office (DPHO) of 5 districts and the BPKIHS. We found that overall the costs of a VL episode were one third less compared to the studies carried out prior to the VL elimination initiative. Despite the free provision of VL drugs, we found that the direct costs of an episode of VL were still high due to out-of-pocket expenditures on ancillary drugs, in particular for households that had visited a private provider prior to receiving VL treatment, as well as food costs during treatment.

In Sudan (chapter 6), SSG was provided as first line treatment where it is still efficacious. Because VL drugs are provided free of charge, direct medical costs were small compared to non-medical costs. The largest cost component to households were food costs for the patient and caretakers when hospitalized for VL treatment. Patients are hospitalized for the full duration of treatment of 30 days to ensure 100% adherence and to monitor for adverse events. Due to severe anaemia and malnutrition patients are at high risk of complications and death (Collin *et al.* 2004). Because patients are always accompanied by one or more family members for the entire duration of hospitalization, food costs represent 78% of the median direct cost. These high food costs were also observed in India and Nepal, and are not often taken into account but are an important determinant of the financial burden of VL to households. Since the food costs are caused by the long hospitalization of the patient, alternative treatment regimens with shorter treatment durations will have a considerable impact on the total cost of a VL episode.

Indirect costs were measured according to the human capital method whereby we quantified the loss of productivity in terms of foregone earnings to the patient or other household members taking care of the patient; unpaid household work was excluded. Basically the median number of days of inactivity due to VL illness was multiplied by the median daily income of the individual (either the patient or another household member). We also considered intra-household labour substitution which is a common strategy that households use to mitigate income losses.

Our studies showed that in India and Nepal, indirect costs represented a considerable proportion of total costs (58% and 53% respectively). VL is characterized by prolonged fever, weight loss, anaemia, fatigue and enlargement of the liver and spleen. As a result patients are either severely limited or not able at all to carry out their daily activities and need much support from family members. We found that patients were not able to work for a median number of 51, 57 and 70 days in Sudan, Nepal and India respectively. The long inactivity of patients was caused by a combination of the delay between the onset of symptoms and the time the patient is correctly diagnosed for VL and the long duration of treatment. Most of the economically active patients in India and Nepal were working as daily labourers such as rickshaw driver or farm labourer. When falling ill, they were rarely replaced by other household members. In contrast indirect costs in Sudan represented only 14% of the total cost of a VL episode. Subsistence farming was the main activity of households in Sudan and there were less patients and caretakers economically active in Sudan compared to India and Nepal. However, in households with a working patient reporting income losses due to VL illness, indirect costs represented 40% of the total cost of a VL episode.

In table 9.1 we summarized direct and total costs (direct and indirect) as a proportion of annual household income by all studies that were carried out in the Indian subcontinent and East Africa. Given the variation in methods between studies, both in terms of the classification of costs and summary statistics used (e.g. median vs. mean values), comparing these results across countries need to be done with caution.

With the exception of two studies in Nepal (including our study presented in chapter 5), household expenditure on medical and non-medical items (i.e. direct costs) were catastrophic to households. Combined with the income losses due to VL illness, the burden of an episode of VL was very high in all studies. We determined costs as catastrophic if they exceeded 10% of annual household income (Ranson 2002). It is hypothesized that if costs incurred by households exceed this threshold, they will need to cut expenditure on basic goods and services, incur debt or sell productive assets which may lead to impoverishment. This is obviously a somewhat arbitrary cut-off point; as we have seen in chapter 3, VL affected households are very poor and have therefore a limited ability to cope with the costs of illness. The catastrophic threshold for these households may therefore be well below this cut-off point.

Table 9.1: Overview of studies investigating the economic burden of visceral leishmaniasis in the Indian subcontinent and East Africa

| Authors | Country | Sample size (HHs) | Direct costs as % of annual HH income ² | Total costs as % of annual HH income ^{2,3} |
|---------------------------------------------------------------------|------------|-------------------|----------------------------------------------------|-----------------------------------------------------|
| (Ch. 4) Meheus <i>et al.</i> 2006 | India | 77 | 14.8 | 21.7 |
| Sarnoff <i>et al.</i> 2010 | India | 182 | 11.3 | 16.0 |
| Sundar <i>et al.</i> 2010 | India | 171 | 27.8 | na |
| Adhkari & Maskay 2003; 2005 ¹ | Nepal | 18 | 15.5 | 35.6 |
| Rijal <i>et al.</i> 2006 | Nepal | 20 | 7.2 | 17.1 |
| Sharma <i>et al.</i> 2004; Adhikari <i>et al.</i> 2009 ¹ | Nepal | 61 | 14.1 | 36.9 |
| (Ch. 5) Uranw <i>et al.</i> 2012 | Nepal | 168 | 3.9 | 11.0 |
| Sharma <i>et al.</i> 2006 | Bangladesh | 87 | 16.1 | 23.3 |
| (Ch. 6) Meheus <i>et al.</i> 2012 | Sudan | 75 | 16.6 | 23.0 |

HH = household; na = not available
¹ These studies used the same dataset of households
² Median values except studies by Adhkari & Maskay 2003; 2005 that presented mean values. These figures are likely to overestimate the median values.
³ Total costs include direct and indirect costs.

In our studies, total household costs per episode of VL were as much as 23% of annual household income in Sudan and nearly 22% in India. In Nepal the economic burden of VL was lower but still catastrophic at 11% of household income. However, in our studies we analysed costs for the last episode of VL. Because of the clustering of VL in communities and villages, several cases of VL often occur in the same household. For example , 31% and 44% of households in Sudan and Nepal respectively had experienced a case of kala-azar before, often within a few years. More than one episode of VL in the household will significantly increase the burden of VL. Even if these cases do not occur in the same year, the risk of impoverishment and indebtedness would still be much higher.

The economic impact of a VL episode is not limited to direct and indirect costs, but also the strategies used by households to cope with the costs of illness add to the burden of VL. These strategies address direct costs (e.g. using savings or borrowing) and/or indirect costs (intra-household labour substitution) (Sauerborn *et al.* 1996). We observed some differences in the coping strategies employed between the Indian subcontinent (India and Nepal) and Sudan that were related to the capacity of households to raise money to pay for the costs of care. In Nepal and India, the first and most common strategy used by households was the mobilization of savings or use of available cash. These “savings” or cash were usually intended to buy food. For the majority of households this was not sufficient to cover the costs of care and 56% and 81% of households in Nepal and India respectively had also to take a loan. The terms of the loan varied depending on the source of the loan (e.g. from friends/family or a money lender). For instance in Nepal, households had to repay on average 140% of the original amount borrowed when borrowing from either an informal money lender or a village member. In contrast in Sudan, few households had savings and relied on cash gifts from village members (59%) or sold crops (57%) or livestock (38%) to raise money. The timing of the survey in Sudan may have influenced the coping strategies as the survey was taken after the harvesting period. About 5% of households in Sudan sold part of their land to pay for the costs of care. Strategies to cope with the indirect costs of illness included intra-household labour substitution and hiring labour to replace the ill household member. In 24% of households in Nepal, children were involved in preserving the income of the household and were removed from school for the duration of illness to replace the VL patient.

In summary, the free provision of VL diagnosis and drugs has been an important policy measure in reducing financial barriers and improving access to VL care. Without this policy the economic burden of VL would have been much higher. However, given the extreme poverty of VL affected households, a VL episode is still catastrophic to most households. One of the main cost determinants to both public hospitals providing VL treatment and households is

the duration of treatment and length of hospitalization. In addition households also incur substantial costs during the health seeking phase (prior to correct VL diagnosis) as well as through the strategies they use to cope with the cost of illness such as loans.

THE COST AND COST-EFFECTIVENESS OF VL TREATMENT STRATEGIES

Part III of the thesis included two studies and investigated the third research question: “*What is the cost and cost-effectiveness of VL treatment strategies?*”. In this section, I will also place the cost and cost-effectiveness findings within a context of priority setting for VL research and control.

Early diagnosis and effective treatment is the main control strategy for VL; in East Africa it is the only strategy that is currently available. Against the background of limited treatment options for VL, and the importance of preserving existing drugs against the development of resistance (see introductory chapter), we reviewed in chapter 7 the evidence and explored the potential of combination therapy for the treatment of VL. With no new compounds expected to come to the market in the near future and the risk of resistance to new medicines, combination therapy is brought forward as a possible alternative. Combining drugs from different chemical classes could also reduce treatment duration or total drug doses, resulting in fewer toxic effects, higher compliance, and less burden on the health system. Combination therapies may also reduce the overall costs (direct and indirect) and be more cost-effective which we will discuss in the next paragraphs.

In chapter 8 we assessed the cost-effectiveness of all possible mono- and combination therapies for the treatment of visceral leishmaniasis in the Indian subcontinent using a decision analytical model based on a decision tree. The analysis was done from a societal perspective and included all costs irrespective of whether they were borne by the government or the patient/household (i.e. health provider and household perspective). We included the household perspective in the analysis since treatments with shorter duration may result in important cost savings to the household. The values related to drug efficacy for therapies that were still in clinical trials at the time of the study were derived through an iterative consultation procedure with a group of VL experts (i.e. Delphi process). These experts were clinicians and/or involved in clinical trials of combination treatment and dose-finding studies. The estimates provided by VL experts proved to be very accurate and were confirmed recently by a phase 3 trial published in 2011. In this study Sundar *et al.* (2011) reported definitive cure rates of at least 97% for the three combination therapies included in our cost-effectiveness study (i.e. 10 days each of miltefosine

and paromomycin (MF+PM), single -dose liposomal amphotericin B and 10-day paromomycin (L-AmB+PM); and single-dose liposomal amphotericin B and 7-day miltefosine (L-AmB+MF)). The treatment durations of these combination therapies were also considerably shorter (range 8-11 days) compared to current monotherapies (range 21-28 days) and corresponded to the values used in our analysis.

Our results suggested that a combination of miltefosine and paromomycin (MF+PM) provided the lowest cost-effectiveness ratio at US\$92 per death averted. The next best alternative was L-AmB+PM with an incremental cost-effectiveness ratio of US\$ 652. Due to a high drug efficacy and expected compliance, the most effective treatment option was a single dose of 10mg per kg of liposomal amphotericin B (L-AmB10). This treatment is also recommended by WHO as the preferred monotherapy (WHO 2010; Matlashewski *et al.* 2011) as it is safe and effective and only a single dose is needed allowing point-of-care diagnosis and treatment on the same day (Van Griensven & Boelaert 2011). Such a strategy would significantly reduce the economic burden to households and would relieve public health care facilities in VL endemic areas particularly during the peak transmission periods. However, L-AmB needs to be administered intravenously and stored below 25°C. In addition, the incremental cost-effectiveness ratio of L-AmB was US\$ 8,224 due to the cost of the drug which is its main barrier to widespread use. Although price negotiations between the WHO and the manufacturer reduced the price of L-AmB to US\$ 20 per vial at the time of the study, as several vials are needed per treatment course the total cost of L-AmB was higher than any of the combination therapies. For L-AmB to become the most cost-effective strategy, we showed in our study that the price would need to be reduced to under US\$ 10 per vial. Since our study, the price of L-AmB has further decreased to US\$ 18 per vial, which will lower the cost of L-AmB strategies but will not change the conclusions of our study.

What are the implications of our findings from the Indian subcontinent to East Africa? The results from chapter 8 cannot be extrapolated to East Africa because of the geographical heterogeneity of VL (van Griensven & Boelaert 2011) and differences in the cost structure between countries (Hutton & Baltussen 2005). For example a phase III study in Africa showed that the cure rate of 15 mg/kg per day of paromomycin for 21 days was inadequate, in particular in Sudan (Hailu *et al.* 2010). Several other clinical trials evaluating combination therapies are currently underway in East Africa (Omollo *et al.* 2011). Similar to the study that was done in the Indian subcontinent, we are currently evaluating the cost-effectiveness of various treatment regimens for Sudan and pending the results from the clinical trials I present here some findings on the costs of the various VL treatment strategies that are currently being evaluated in Sudan. These include 3 combination therapies (strategies 1-3) and 4 monotherapies (strategies 4-7) (table 9.2).

Table 9.2: Overview of possible treatment strategies in Sudan

| | Strategies | Length of treatment (days) |
|---|-------------------------------|----------------------------|
| 1 | L-AmB (10MK) + MF (50/100 MD) | 11 |
| 2 | L-AmB (10MK) + SSG (20 MKD) | 11 |
| 3 | SSG (20 MKD) + PM (15 MKD) | 17 |
| 4 | L-AmB (30 MKD) | 10 |
| 5 | SSG (20 MKD) | 30 |
| 6 | MF (50/100 MD) | 28 |
| 7 | Glucantime (20 MKD) | 20 |

L-AmB : Liposomal Amphotericin B; MF : miltefosine; SSG: Sodium stibogluconate; PM : paromomycin
 MK = mg/kg single dose; MD = mg per day; MKD = mg/kg body weight per day.
 Miltefosine if given at 50 mg/day if body weight is <25 or 100 mg/day if body weight ≥ 25 kg.

The costs of the various treatment alternatives are as follows (table 9.3):

Table 9.3: Cost estimates of each treatment strategy per patient treated (US\$ 2010)

| | Strategies | Drug cost | Other direct medical ¹ | Non-medical & indirect | Total cost ² |
|---|-------------|-----------|-----------------------------------|------------------------|-------------------------|
| 1 | L-AmB + MF | 139.6 | 64.9 | 41.1 | 245.6 |
| 2 | L-AmB + SSG | 148.0 | 64.3 | 41.1 | 253.4 |
| 3 | SSG + PM | 41.5 | 99.1 | 63.5 | 204.0 |
| 4 | L-AmB | 378.0 | 64.1 | 37.3 | 479.5 |
| 5 | SSG | 51.3 | 173.5 | 112.0 | 336.9 |
| 6 | MF | 38.1 | 163.5 | 104.5 | 306.1 |
| 7 | Glucantime | 42.0 | 115.7 | 74.7 | 232.4 |

¹ Calculated using the average hotel unit cost of US\$ 5.74 across the three hospitals

² The cost estimations are preliminary as more definite information on resource usage may become available from the clinical trials.

The category “non-medical and indirect” costs are entirely borne by the households while the other two categories (“drug” and “other direct medical”) are mainly borne by the health care provider. With the exception of treatment with L-AmB, all other alternatives are cheaper than the current first-line treatment SSG. While a MF-strategy has the lowest drug cost, these are largely offset by the high cost of hospitalization and the household costs. As explained in chapter 6, VL patients in Sudan are hospitalized for the entire duration of treatment. This is also likely to be the case with the oral drug miltefosine. In contrast the shorter treatment durations

and resulting low cost of hospitalization and household costs of combination therapies with a low dosage of L-AmB (requiring fewer vials than the monotherapy) “compensates” for the higher cost of the drug. The cheapest strategy was a combination of SSG and PM at US\$ 204.

Priority setting for neglected tropical diseases (NTDs) – making a case for investment in visceral leishmaniasis control

In the past decade, a renewed interest in NTD research and control has emerged in both the public and private sectors leading to the establishment of various initiatives and public-private partnerships (PPPs), such as the World Health Organization NTD programme, the USAID programme on NTDs, the Global Network for NTDs (GNNTD), the Drugs for Neglected Diseases Initiative (DNDi) and various vertical initiatives such as the Schistosomiasis Control Initiative (SCI). An important milestone was reached end January 2012 in London where global health leaders, national and international policy makers and the heads of major pharmaceutical companies committed to control or eliminate 10 neglected tropical diseases (NTDs) by the end of 2020. The London Declaration on Neglected Tropical Diseases endorsed during this meeting followed years of intense advocacy efforts conducted to increase public awareness and funding for the NTDs.

Advocates were particularly successful in raising the profile of seven NTDs described as “tool-ready” by WHO by demonstrating the low cost and cost-effectiveness of interventions to control these diseases. The seven NTDs include six helminthic infections (soil-transmitted helminths (ascariasis, trichuriasis, hookworm), lymphatic filariasis, onchocerciasis and schistosomiasis) and a bacterial infection, trachoma. The main control strategy consists of providing single doses of one or more drugs (albendazole, ivermectin, praziquantel and/or azithromycin) targeting the seven NTDs once or twice per year through mass drug administration (MDA) based on the WHO guidelines for preventive chemotherapy (WHO 2006). Strategies are low-cost (and safe and effective, hence the term “tool-ready) because most of the drugs are donated by private pharmaceutical companies (Fenwick *et al.* 2005) or purchased at low prices. Diseases such as VL or human African trypanosomiasis (HAT) were not considered among these “tool-ready” diseases.

While the emphasis on low-cost and cost-effectiveness were strong arguments to make a case for investment in the “tool-ready” NTDs (Molyneux *et al.* 2005; Canning 2006; Hotez *et al.* 2009; Ruxin 2009; Conteh *et al.* 2010), it was less “beneficial” for other NTDs that currently lack (known) cost-effective interventions and do not receive similar support from private pharmaceutical companies (the “tool-deficient” diseases). VL or HAT therefore risk being labelled low priority by national programmes and international donors.

With the evidence presented in this thesis and elsewhere, can we make a case for investment in VL? How cost-effective is VL treatment and how cost-effective is VL treatment relative to other neglected tropical diseases?

The cost-effectiveness study presented in chapter 8 showed that average cost-effectiveness ratios for VL treatment ranged from US\$ 91 to US\$ 162 per death averted depending on the type of drug regimen. With a cost less than US\$ 100 to save a life (for treatment with MF+PM), this seems to be very good “value for money”. There is however no universal threshold value below which an intervention is generally considered cost-effective. Suggested thresholds consider an intervention to be very cost-effective when the cost-effectiveness ratio is lower than gross domestic product (GDP) per capita and cost-effective if it is between 1 and 3 times GDP per capita (Commission on Macroeconomics and Health 2001; WHO 2012). In the case of our study in chapter 8, relative to the country with the lowest GDP in the Indian subcontinent (i.e. Nepal with US\$ 620 in 2011; World Bank 2012) VL treatment can be judged a highly cost-effective intervention. Others suggested an intervention to have a “highly attractive” cost-effectiveness if the cost per DALY averted is less than US\$ 25 and “attractive” if it is less than US\$ 150 (World Bank 1993). As explained in the first section of this discussion, we had chosen deaths averted as an outcome measure since VL treatment is a lifesaving intervention. Because comparative assessments are now often done in terms of disability adjusted life years (DALYs) (e.g. the Disease Control Priorities in Developing Countries project (Jamison *et al.* 2006), I provide here an approximation of DALYs for VL. DALYs are calculated as the sum of the years of life lost due to premature mortality (YLL) and the years of life lived in disability (YLD) (Murray & Lopez 1996; Fox-Rushby & Hanson 2001). DALYs were calculated following the approach by Fox-Rushby & Hanson (2001) using a number of simplifying assumptions¹. Firstly I assumed a 100% case fatality rate for untreated VL and instantaneous death once clinical symptoms occur although the time period between the onset of symptoms and death for untreated VL is about 3 to 5 months. Secondly, given this brief time period between the onset of symptoms and death for untreated VL, YLDs as a share of total DALYs is likely to be small. Since long-term disability after VL treatment is also infrequent, YLDs were not included in the DALY calculation². The

¹ The formula used to estimate YLLs is as follows (Murray & Lopez 1996; Fox-Rushby & Hanson 2001):

$$YLL[r, K, \beta] = \frac{KCe^{ra}}{(r + \beta)^2} \{e^{-(r+\beta)(L+a)}[-(r + \beta)(L + a) - 1] - e^{-(r+\beta)a}[-(r + \beta)a - 1]\} + \frac{1 - K}{r}(1 - e^{-rL})$$

Where: K = age weighting modulation factor; C = Constant; r = discount rate; a = age of death; = parameter from the age weighting function; and L = standard expectation of life at age a .

² There are also some important methodological issues with the estimation of YLDs for VL, in particular related to the disability weight (DW), which is a measure of disease severity. The DW for VL is 0.243 (Mathers *et al.* 2007) and is too low (a score of 0 represents perfect health and 1 equals death). For example, the weight for an uncomplicated episode of malaria is much higher at 0.471, while we found from our surveys that the vast majority of VL patients are not able to engage in productive activities at all when ill.

data on life expectancy at age a were derived from life tables for Nepal (Lopez *et al.* 2001) and DALYs were calculated in five-year age intervals using data on age distribution from our household survey in Nepal (chapter 5).

In the absence of VL treatment, the total number of DALYs [0.03,1,0.04] lost per person equals 30.1. This number is relatively high because (i) untreated VL is fatal and (ii) many cases of VL occur among young adults that are given more weight in the DALY estimation. Without age weighting ($K=0$), the total number of DALYs [0.03,0,0] lost equals 24.6. Carrying out the analysis for the strategies MF+PM, L-AmB+PM and L-AmB (10mg/kg) of chapter 8, we find that cost-effectiveness ratios vary from US\$ 2.9 to US\$ 5.1 per DALY averted. The only other estimation available in the literature on DALYs for VL was made by Griekspoor *et al.* (1999). They estimated the cost-effectiveness of VL treatment in an emergency situation in Sudan and obtained a cost-effectiveness ratio of US\$ 18.4 per DALY averted. This estimate is higher than ours because the average cost per patient treated was higher (i.e. US\$ 394) and the authors assumed a case fatality rate for untreated VL below 100%.

Compared to other neglected tropical diseases, our estimates are very similar to those for mass drug administration for lymphatic filariasis (US\$ 4 to US\$ 8 per DALY averted) (Jamison *et al.* 2006; Laxminarayan *et al.* 2006; Ottesen *et al.* 2008) and combined school-based treatment of intestinal worms and schistosomiasis (US\$ 10 to US\$23 per DALY averted) (Miguel & Kremer 2004; Conteh *et al.* 2010).

In summary, we have shown here that interventions for VL treatment are within the range of highly cost-effective public health interventions (Laxminarayan *et al.* 2006). A number of observations need to be made regarding this finding. First, as we have seen throughout our studies, both the measurement of the general burden of disease through DALYs and cost-effectiveness estimates (cost per DALY or death averted) “hide” the economic burden of VL to patients and their family. While interventions may be cost-effective, they ignore the substantial hardship that is placed on households. As we have shown in our studies in Nepal and Sudan even in situations whereby patients are not charged for diagnosis or VL drugs, households still incur high costs mainly as a result of the long duration of hospitalization and the delay to correct VL diagnosis. Implementing treatment interventions with shorter treatment duration, such as combination therapies, are therefore a crucial component in reducing the cost to households. A limitation of current studies is that they only captured the short-term effects of VL illness. In the studies presented in section 2 of the thesis, we have mainly documented household direct and indirect costs, as well as the strategies used by these households to cope with these costs. No studies to date have examined the long term effects of a VL episode and

the coping strategies used on the wealth and livelihood of affected households. Better insights into these processes may provide additional arguments for investing in VL treatment and in particular prevention, and find ways to mitigate the economic burden of VL.

Secondly while VL treatment may be very cost-effective according to the thresholds mentioned in previous paragraphs, this does not mean however that VL treatment is affordable to national VL control programmes. The issue of affordability is an often forgotten criterion in health-care priority setting (Cleary and McIntyre 2009). While the cost per death or DALY averted is relatively low, VL occurs in countries facing many competing health priorities and constrained health resources. It is therefore necessary to not only estimate the cost and cost-effectiveness of interventions, but also the budgetary impact of VL control. With the recent updated estimates of VL incidence, such an estimation is now possible although the authors of the study acknowledge that uncertainties inherent in their data remain (Alvar *et al.* 2012). In addition, continued advocacy is needed to bring the price of VL drugs further down, in particular the price of liposomal amphotericin B. End 2011, Gilead Sciences, the manufacturer of AmBisome® (i.e. liposomal Amphotericin B), signed an agreement with WHO to donate drugs that will allow 50,000 patients to be treated in the next 5 years. With 10,000 patients per year, this donation only covers a small proportion of the annual number of VL cases. While drug donations are essential during emergency relief situations, such as a VL epidemic, more sustainable solutions should be promoted such as reducing the price of AmBisome® rather than relying on drug donations. According to a press statement released as part of the London Declaration on NTDs (United to Combat NTDs press release 30/01/20123), “Gilead, ..., will continue its program to offer VL at cost and commit to investigate and invest in technologies and processes that could reduce that cost in resource-limited countries”. The “no-profit” price of AmBisome® is US\$ 18 per vial (Gilead press release 20114); as we have shown, the price per vial of AmBisome® needs to be reduced below US\$ 10 to become the most cost-effective strategy to treat the disease.

RECOMMENDATIONS FOR VL POLICY

The free provision of VL diagnosis and drugs have been important policy measures to lower financial barriers and improve access to VL diagnosis and care. Without this policy the economic burden of VL would have been much higher. However, the economic impact of VL is still considerable and intensified efforts are needed to further reduce the burden of VL for

affected households. The implications of the studies presented in this thesis conducted from various perspectives call for a change in current policies.

On the basis of our findings, the following recommendations can be formulated:

→ Reduce delays between onset of symptoms and correct VL diagnosis.

The costs incurred by patients can basically be divided in two time periods: (i) costs incurred prior to correct VL diagnosis during the health seeking phase and (ii) the costs incurred when receiving treatment (discussed in the next paragraph).

We have shown that households incur substantial expenses when seeking care. The longer the delay between symptoms onset and access to VL care, the higher the household costs are likely to be. We found that the median lag time between symptoms onset and presentation to a qualified provider was 1-2 months; in particular at private providers in India the patient was not often diagnosed with VL on the first visit. There are a number of lessons that can be drawn here. In India, given the importance of the private sector, there is a need to involve private health practitioners in VL control efforts by ensuring compliance to diagnosis and treatment practices as defined in the national guidelines as well as regular reporting of all diagnosed and treated cases to district health authorities. Since our study, a rural outreach programme has been rolled out in 18 states, including Bihar, with the introduction of voluntary community health workers (ASHA) in every village and additional auxiliary nurse midwives at sub-centres (Ministry of Health and Family Welfare). Similar to malaria control activities at the community level (Sharma and Dutta 2011), ASHAs can play a role in VL control activities. The rK39 rapid diagnostic test (RDT) is easy to perform and can be used by ASHAs in villages and provide an important contribution to the early detection of VL cases with subsequent referral to primary health care centres for treatment. It seems however that ASHAs and other community workers are not yet involved in the diagnosis of VL (Srivastava *et al.* 2009) and according to a study by Hasker *et al.* (2010), only few VL patients had first reported to ASHAs. Additional investigation into the role of community workers is needed, including studies on the current knowledge, attitude and practices of ASHAs and other health workers active at the community level in India as was done for tuberculosis and the DOTS programme (Sagare *et al.* 2012; Malaviya *et al.* 2012). Unlike the Indian subcontinent, VL diagnosis at the community level by village health workers may not be a feasible strategy in Sudan. Not only does the overall lack of human and financial resources allocated to the VL programme hamper the development of decentralized control policies, recent studies showed varying sensitivity between different brands of the rK39 RDT in East Africa with several RDTs inadequate for excluding a VL diagnosis when used alone (WHO 2011; Cunningham *et al.* 2012).

³ United to Combat Neglected Tropical Diseases Press Release, 30 January 2012: http://www.unitingtocombatntds.org/downloads/press/ntd_event_press_release.pdf

⁴ Gilead Sciences Press Release, 08 December 2011: http://www.gilead.com/pr_1637695

→ Change the first line treatment

An important determinant of household and hospital (i.e. provider) costs is the duration of treatment of current first line regimens; respectively 28 and 30 days in the Indian subcontinent and East Africa. The extended length of stay poses a significant burden to households and hospitals. As shown in this thesis, households incur substantial non-medical costs, in particular on food, and income losses while the length and cost of hospitalization are a heavy burden to the public health care system especially during the peak transmission period.

Although making strong recommendations about the best treatment options is beyond the scope of this thesis, we have shown that, from a societal perspective, more effective and cheaper alternatives are available. Combination therapies and L-AmB monotherapy are possible alternatives for the current first-line treatment with miltefosine in the Indian subcontinent, while a preliminary cost analysis for East Africa presented above suggest that a combination of SSG and PM is the cheapest and has a similar efficacy to the current first line (Musa *et al.* 2012). The latter strategy is already successfully implemented in East Africa by non-governmental organizations (NGO's) such as Médecins sans Frontières; similarly L-AmB based strategies are also used on a small scale by NGO's in India at the primary health care level and have overcome the cold chain constraints.

The findings on cost and cost-effectiveness presented in this thesis support the clinical evidence to integrate these treatment strategies into national guidelines for the management of VL and are also in line with the recommendations of the WHO expert committee on leishmaniasis. This should be done as soon as possible given the demonstrated economic impact of VL illness and treatment costs on household wealth.

→ Bring the price of L-AmB further down

An important bottleneck to the implementation of L-AmB based treatment strategies and the inclusion of the drug in national guidelines in the Indian subcontinent is its price. Since price and performance are often the main criteria in tender processes and with cheaper alternatives available, it is unlikely that governments will adopt L-AmB treatment strategies at the current price per vial (i.e. US\$ 18 per vial; about US\$ 280 for a full treatment course). It is therefore necessary to bring the price per L-AmB vial further down. We showed in this thesis that the price of a L-AmB vial needs to be reduced to below US\$10 to become cost-effective. However obtaining further price reductions from the manufacturer is not going to be evident. A possible solution could be the use of so called “pull” mechanisms whereby a specified level of demand for the drug is ensured and thereby mitigating the unpredictability in market demand to the manufacturer. Establishing a larger and more credible market for the drug may also stimulate

generic manufacturers to develop generic L-AmB formulations which may in turn bring the price further down.

An example of such a pull mechanism is the advance market commitment (AMC) whereby a financial commitment is made for future purchases at a pre-agreed price (Batson *et al.* 2006). Currently AMCs have mainly been used to accelerate the introduction of vaccines in developing countries (e.g. the new pneumococcal vaccines). The potential of these kinds of innovative financing mechanisms to improve access to VL drugs in endemic countries should be further investigated.

→ Reduce household non-medical costs

To mitigate the impact of direct costs to households (medical and non-medical), national control programmes ought to consider introducing demand side financing mechanisms. These financing arrangements are more realistic on the short term than health insurance since coverage of the latter is currently very low or mostly non-existent in the three study countries. Furthermore, insurance primarily covers direct medical costs at accredited formal health care practitioners while we have seen in part II of the thesis that non-medical costs on transportation and in particular food are as important or even greater than direct medical costs.

With demand side-financing households receive a subsidy (i.e. public funds) to purchase defined goods and services in order to improve access to these services. Demand side financing mechanisms include for example voucher schemes or conditional cash transfers. These mechanisms have already been used in the study countries as humanitarian cash transfers on nutrition in Sudan or to improve access to child and maternal health services in India. Demand side financing is feasible for VL since public health authorities in Nepal now provide a one-time payment of Rs 1,000 to households that completed VL treatment to cover transportation costs (Adhikari *et al.* 2011) and is implemented through the district public health offices. Given the importance of food costs, authorities in Nepal should consider expanding the programme to include these costs as well. There is however little information available on the programme in terms of coverage or beneficiaries.

CONCLUDING REMARKS

With a renewed interest in neglected tropical diseases and the implementation of a VL control initiative in the Indian subcontinent, the studies presented in this thesis contributed to ongoing VL policies and research efforts in various ways. With new technologies and treatments that have become available, there was an urgent need to provide national and international policy

makers with sound and systematic economic evidence that enable the design of more rational and effective case management policies. This was particularly important given the resource constraints VL endemic countries face, and the severe economic impact of VL to households.

Equally important was to raise awareness about the economic burden of VL to affected households in order to draw the attention of national and global health policy makers to the need for further action and support in both research and policy. This effort will need to be intensified as VL has received relatively little attention compared to other neglected tropical diseases. Here, economic arguments about cost and cost-effectiveness can contribute to making a compelling case for investment in the control of VL.

Finally, I have studied for my thesis the economic burden of VL and the cost and cost-effectiveness of VL patient management. Improved treatment policies are one side of the story. Successful VL control or elimination will only be possible if sufficient resources are also invested in the prevention of VL through vector control and ensuring access to rapid diagnosis and treatment, removing financial barriers and improving the quality of VL care.

REFERENCES

- Adhikari SR, Maskay NM (2003) The economic burden of Kalaazar in households of the Danusha and Mahottari districts of Nepal. *Acta Trop* 88: 1–2.
- Adhikari SR, Maskay NM (2005) Economic cost and consequences of kala-azar in Dhanusha and Mahottari districts of Nepal. *Indian J Community Med* 30: 121–125.
- Adhikari SR, Maskay NM, Sharma BP (2009) Paying for hospital-based care of kala-azar in Nepal: assessing catastrophic impoverishment and economic consequences. *Health Policy Plann* 24: 129–139.
- Adhikari SR, Prasai DP, Sharma SK (2011) *A review of demand side financing schemes in the health sector in Nepal*. Nepal Health Sector Support Programme.
- Ahluwalia IB, Bern C, Costa C, et al. (2003) Visceral leishmaniasis: consequences of a neglected disease in a Bangladeshi community. *Am J Trop Med Hyg* 69: 624–628.
- Alvar J, Vélez ID, Bern C, et al. (2012) Leishmaniasis Worldwide and Global Estimates of Its Incidence. *PLoS ONE* 7: e35671.
- Banjara MR, Hirve S, Siddiqui NA, et al. (2012) Visceral leishmaniasis clinical management in endemic districts of India, Nepal, and Bangladesh. *J Trop Med*. 2012: 126093.
- Batson A, Meheus F, Brooke S (2006) Innovative financing mechanisms to accelerate the introduction of HPV vaccines in developing countries. *Vaccine*, S3: S219–25.
- Bern C, Courtenay O, Alvar J (2010) Of Cattle, Sand Flies and Men: A Systematic Review of Risk Factor Analyses for South Asian Visceral Leishmaniasis and Implications for Elimination. *PLoS Negl Trop Dis* 4: e599.
- Bern C, Joshi AB, Jha SN, et al. (2000) Factors associated with visceral leishmaniasis in Nepal: bed-net use is strongly protective. *Am J Trop Med Hyg* 63: 184–188.
- Bern C, Maguire JH & Alvar J (2008) Complexities of assessing the disease burden attributable to leishmaniasis. *PLoS Negl Trop Dis* 2: e313.
- Canning D (2006) Priority setting and the ‘neglected’ tropical diseases. *Trans R Soc Trop Med Hyg* 100: 499–504.
- Cleary SM, McIntyre D (2009) Affordability - the forgotten criterion in health-care priority setting. *Health Econ* 18: 373–5
- Collin S, Davidson R, Ritmeijer K, et al. (2004) Conflict and kala-azar: determinants of adverse outcomes of kala-azar among patients in southern Sudan. *Clin Infect Dis* 38: 612–619.
- Commission on Macroeconomics and Health (2001) Macroeconomics and Health: Investing in Health for Economic Development. Chaired by Jeffrey D. Sachs. Report of the Commission on Macroeconomics and Health. Geneva: World Health Organization.
- Conteh L, Engels T, Molyneux DH (2010) Socioeconomic aspects of neglected tropical diseases. *Lancet* 375: 239–247.

- Cunningham J, Hasker E, Das P, et al. (2012) A Global Comparative Evaluation of Commercial Immunochromatographic Rapid Diagnostic Tests for Visceral Leishmaniasis. *Clin Infect Dis* 55: 1312-1319.
- Das P, Samuels S, Desjeux P, et al. (2010) The annual incidence of visceral leishmaniasis in an endemic area of Bihar, India. *Trop Med Int Health* 15(S2): 4-11.
- Dixon S, McDonald S, Roberts J (2002) The impact of HIV and AIDS on Africa's economic development. *BMJ* 324: 232-4.
- Drummond M, Sculpher MJ, Torrance GW, O'Brien BJ, Stoddart GL (2005) *Methods for the economic evaluation of health care programmes*. 3rd ed. Oxford: Oxford University Press.
- Engels D, Savioli L (2006) Reconsidering the underestimated burden caused by neglected tropical diseases. *Trends Parasitol* 22: 363-6.
- Fenwick A, Molyneux D, Nantulya V (2005) Achieving the millennium development goals. *Lancet* 365: 1029-1030.
- Fox-Rushby JA, Hanson K (2001) Calculating and presenting disability adjusted life years (DALYs) in cost-effectiveness analysis. *Health Policy Plan* 16: 326-31.
- Gallup J, Sachs J (2001) The economic burden of malaria. *Am J Trop Med Hyg* 64: 85-96.
- Gerstl S, Amsalu R, Ritmeijer K (2006) Accessibility of diagnostic and treatment centres for visceral leishmaniasis in Gedaref State, northern Sudan. *Trop Med Int Health* 11: 167-175.
- Government of India, Ministry of Health and Family Welfare. *National Rural Health Mission: Meeting people's needs in rural health areas*. Framework for implementation 2005-2012.
- Available online at: http://mohfw.nic.in/NRHM/Documents/NRHM_Framework_Latest.pdf. Accessed on August 23, 2012.
- Griekspoor A, Sondorp E, Vos T (1999) Cost-effectiveness analysis of humanitarian relief interventions: visceral leishmaniasis treatment in the Sudan. *Health Policy Plan* 14: 70-76.
- Hailu A, Musa A, Wasunna M, et al. (2010) Geographical Variation in the Response of Visceral Leishmaniasis to Paromomycin in East Africa: A Multicentre, Open-Label, Randomized Trial. *PLoS Negl Trop Dis* 4: e709.
- Hasker E, Singh SP, Malaviya P, et al. (2012) Visceral leishmaniasis in rural Bihar, India. *Emerg Infect Dis* 18: 1662-4
- Hasker E, Singh SP, Malaviya P, et al. (2010) Management of visceral leishmaniasis in rural primary health care services in Bihar India. *Trop Med Int Health* 15 (S2): 55-62
- Hotez PJ, Ottesen E, Fenwick A, Molyneux D (2006) The neglected tropical diseases: the ancient afflictions of stigma and poverty and the prospects for their control and elimination. *Adv Exp Biol Med* 582: 22-33.
- Hotez PJ, Fenwick A, Savioli L, Molyneux DH (2009) Rescuing the bottom billion through control of neglected tropical diseases. *Lancet* 373: 1570-5.
- Hutton G, Baltussen R (2005) Cost valuation in resource-poor settings. *Health Pol Plann* 20: 252-259.
- Jamison DT, Breman JG, Measham AR, et al. (eds.) (2006) *Disease control priorities in developing countries*. 2nd edn. New York: Oxford University Press.
- Jefferis K, Kinghorn A, Siphambe H, Thurlow J (2008) Macroeconomic and household-level impacts of HIV/AIDS in Botswana. *AIDS* 22(S1): S113-9.
- King CH & Bertino AM (2008) Asymmetries of poverty: why global burden of disease valuations underestimate the burden of neglected tropical diseases. *PLoS Negl Trop Dis* 2: e209.
- Laxminarayan R, Mills AJ, Breman JG, et al. (2006) Advancement of global health: key messages from the Disease Control Priorities Project. *Lancet* 367: 1193-208.
- Lee BY, Bacon KM, Shah M, Kitchen SB, Connor DL, Slayton RB (2012) The economic value of a visceral leishmaniasis vaccine in Bihar state, India. *Am J Trop Med Hyg* 86: 417-25.
- Lopez AD, Ahmad OB, Guillot M, Inoue M, Ferguson BD, Salomon JA (2001) Life tables for 191 countries for 2000: data, methods, results. GPE Discussion Paper 40. Geneva: World Health Organization.
- Lutumba P, Makieya E, Shaw A, Meheus F & Boelaert M (2007) Human African Trypanosomiasis in a Rural Community, Democratic Republic of Congo. *Emerg Infect Dis* 13: 248-254.
- Luz PM, Grinsztejn B, Galvani AP (2009) Disability adjusted life years lost to dengue in Brazil. *Trop Med Int Health* 14 :237-46.
- Malaviya P, Hasker E, Singh R, Van Geertruyden JP, Boelaert M, Sundar S (2012) Village health workers in Bihar, India: an untapped resource in the struggle against kala azar. *Trop Med Int Health*. Accepted.
- Mathers CD, Ezzati M, Lopez AD (2007) Measuring the Burden of Neglected Tropical Diseases: The Global Burden of Disease Framework. *PLoS Negl Trop Dis* 1: e114.
- Matlashewski G, Arana B, Kroeger A, et al. (2011) Visceral leishmaniasis: elimination with existing interventions. *Lancet Infect Dis* 11: 322-5.
- Meheus F, Abuzaid AA, Baltussen R, Brima M, Boelaert M, Musa M (2012) The economic burden of visceral leishmaniasis in Sudan: an assessment of hospital and household costs. *Submitted for publication*.
- Meheus F, Boelaert M (2010) The burden of visceral leishmaniasis in South Asia. *Trop Med Int Health* 15: 1-3.
- Meheus F, Boelaert M, Baltussen R, Sundar S (2006) The costs of patient management of visceral leishmaniasis in Muzaffarpur, India. *Tropical Medicine and International Health*; 2006. 11: 1715-1724
- Miguel EA, Kremer M (2004) Worms: identifying impacts on education and health in the presence of treatment externalities. *Econometrica* 72: 159-217.
- Molyneux DH, Hotez PJ, Fenwick A (2005) "Rapid-impact interventions": how a policy of integrated control for Africa's neglected tropical diseases could benefit the poor. *PLoS Med* 2005 2: e336.
- Mondal D, Singh SP, Kumar N, et al. (2009) Visceral leishmaniasis elimination programme in India, Bangladesh, and Nepal: reshaping the case findings/case management strategy. *PLoS Negl Trop Dis* 3: e355.
- Murray CJ, Lopez AD (1994) Quantifying disability: data, methods and results. *Bull World Health Organ* 72: 481-494.

- Murray JL, Lopez AD (1996) The global burden of disease. A comprehensive assessment of mortality and disability from diseases, injuries, and risk factors in 1990 and projected to 2020. Cambridge: Harvard University Press.
- Musa A, Khalil E, Hailu A, et al. (2012) Sodium Stibogluconate (SSG) & Paromomycin Combination Compared to SSG for Visceral Leishmaniasis in East Africa: A Randomised Controlled Trial. *PLoS Negl Trop Dis* 6: e1674.
- Omollo R, Alexander N, Edwards T, et al. (2011) Safety and efficacy of miltefosine alone and in combination with sodium stibogluconate and liposomal amphotericin B for the treatment of primary visceral leishmaniasis in East Africa: study protocol for a randomized controlled trial. *Trials* 12:166.
- Ottesen EA, Hooper PJ, Bradley M, Biswas G (2008) The Global Programme to Eliminate Lymphatic Filariasis: Health Impact after 8 Years. *PLoS Negl Trop Dis* 2: e317.
- Pascual Martínez F, Picado A, Roddy P, Palma P (2012) Low castes have poor access to visceral leishmaniasis treatment in Bihar, India. *Trop Med Int Health* 17: 666-73
- Ranjan A, Sur D, Singh VP, et al. (2005) Risk factors for Indian kala-azar. *Am J Trop Med Hyg* 73: 74–78.
- Ranson MK (2002) Reduction of catastrophic health care expenditures by a community-based health insurance scheme in Gujarat, India: current experiences and challenges. *Bull World Health Organ* 80: 613-621.
- Reithinger R (2008) Leishmaniasis' burden of disease: ways forward for getting from speculation to reality. *PLoS Negl Trop Dis* 2: e285.
- Rijal S, Chappuis F, Singh R, et al. (2003) Treatment of visceral leishmaniasis in south-eastern Nepal: decreasing efficacy of sodium stibogluconate and need for a policy to limit further decline. *Trans R Soc Trop Med Hyg* 97: 350-354.
- Rijal S, Koirala S, van der Stuyft P, Boelaert M (2006) The economic burden of visceral leishmaniasis for households in Nepal. *Trans R Soc Trop Med Hyg* 100: 838–841
- Ruxin J (2009) Case for investment in neglected tropical diseases. *Journal of Lymphoedema* 4: 99-101
- Sagare SM, Bogam RR, Murarkar SK, Patil UP, Ghatte MM (2012) Knowledge, attitude and practices of ASHAs regarding tuberculosis and DOTS. *Indian Journal of Science and Technology* 5: 2401-2404
- Saha S, Ramachandran R, Hutin YJ, Gupte MD (2008) Visceral leishmaniasis is preventable in a highly endemic village in West Bengal, India. *Trans R Soc Trop Med Hyg* 103: 737-42
- Sarnoff R, Desai J, Desjeux P, et al. (2010) The economic impact of visceral leishmaniasis on rural households in one endemic district of Bihar India. *Trop Med Int Health* 15(S2): 42–49
- Sauerborn R, Adams A & Hien M (1996) Household strategies to cope with the economic costs of illness. *Soc Sci Med* 43: 291–301.
- Schenkel K, Rijal S, Koirala S, et al. (2006) Visceral leishmaniasis in southeastern Nepal: a cross-sectional survey on Leishmania donovani infection and its risk factors. *Trop Med Int Health* 11: 1792–1799.
- Shah H (2005) Prevalence of visceral leishmaniasis associated with the behaviors of the people in rural areas. *J Nepal Med Assoc* 44: 116-20.
- Sharma AD, Bern C, Varghese B, et al. (2006) The economic impact of visceral leishmaniasis on households in Bangladesh. *Trop Med Int Health* 11: 757–764
- Sharma BP, Maskay NM, Adhikari SR, et al. (2004) Socio-economic determinants of kala-azar in Nepal. *J Nepal Health Res Counc* 2: 35–42
- Sharma R, Dutta AK (2011) Malaria and National Vector Borne Disease Control Programme. *Indian J Pediatr* 78: 1527-35.
- Singh S, Kumar J, Singh R, Dwivedi SN (2000) Hepatitis B and C viral infections in Indian kala-azar patients receiving injectable anti-leishmanial drugs: a community-based study. *Int J Infect Dis* 4: 203-8.
- Singh SP, Hasker E, Picado A, et al. (2010) Risk factors for visceral leishmaniasis in India: further evidence on the role of domestic animals. *Trop Med Int Health* 15 (S2): 29-35.
- Srivastava DK, Prakash S, Adhish V, Nair KS, Gupta S, Nandan D (2009) A study of interface of ASHA with the community and the service providers in Eastern Uthar Pradesh. *Indian J Public Health* 53: 133-136.
- Sundar S, Arora R, Singh SP, Boelaert M, Varghese B (2010) Household cost-of-illness of visceral leishmaniasis in Bihar, India. *Trop Med Int Health* 15 (S2): 50-54.
- Sundar S, More DK, Singh MK, et al. (2000) Failure of pentavalent antimony in visceral leishmaniasis in India: report from the center of the Indian epidemic. *Clin Infect Dis* 31: 1104-1107.
- Sundar S, Sinha PK, Rai M, et al. (2011) Comparison of short-course multidrug treatment with standard therapy for visceral leishmaniasis in India: an open-label, non-inferiority, randomised controlled trial. *Lancet* 377: 477-86.
- Uranw S, Meheus F, Baltussen R, Rijal S, Boelaert M (2012) The economic burden of visceral leishmaniasis on households in eastern Nepal. *BMC Infect Dis*. In press.
- van Griensven J, Boelaert M (2011) Combination therapy for visceral leishmaniasis. *Lancet* 377: 443-4
- Vassall A, Compennolle P (2006) Estimating the resource needs of scaling-up HIV/AIDS and tuberculosis interventions in sub-Saharan Africa: a systematic review for national policy makers and planners. *Health Policy* 79: 1-15.
- World Health Organization (1996) Global health statistics. *A compendium of incidence, prevalence, and mortality estimates for over 200 conditions*. Murray CJ, Lopez LD (eds.) Global Burden of Disease and Injury Series, Volume II. Atlanta, GA: World Health Organization.
- World Health Organization (2004) The global burden of disease: 2004 update. Geneva: World Health Organization
- World Health Organization (2006) *Preventive chemotherapy in human helminthiasis. Coordinated use of anthelmintic drugs in control interventions: a manual for health professionals and programme managers*. Geneva: World Health Organization.
- World Health Organization (2010) *Control of the Leishmaniases. Report of a meeting of the WHO Expert Committee on the Control of the Leishmaniases*. Geneva: World Health Organization.
- World Health Organization (2012) *Cost-effectiveness thresholds*. Available at: http://www.who.int/choice/costs/CER_thresholds/en/index.html. Accessed on November 13, 2012.

WHO Regional Office for the Eastern Mediterranean (2006) *Health system profile Sudan*. Egypt: Cairo.

World Health Organization and UNICEF/UNDP/World Bank/WHO Special Programme for Research and Training in Tropical Diseases (2011) *Visceral leishmaniasis rapid diagnostic test performance*. Diagnostics Evaluation Series No4. Geneva: World Health Organization.

World Health Organization-CHOICE. *Country specific unit costs*. Available at: http://www.who.int/choice/country/country_specific/en/index.html. Accessed on: July 15, 2012

World Bank (1993) *World development report*. Washington, DC: World Bank.

SUMMARY



SUMMARY

Visceral leishmaniasis (VL) is a vector-borne disease caused by *Leishmania* parasites that are transmitted by the bite of an infected female sandfly. VL is an important public health problem and a significant cause of mortality and morbidity in the Indian subcontinent and East Africa. Since 2005 the governments of India, Nepal and Bangladesh are engaged in a collaborative effort to control and eliminate visceral leishmaniasis from the Indian subcontinent. Approaches to control aim on the one hand at improving the human recovery rate through early diagnosis and case management and on the other hand reducing vector density and biting rate through vector control interventions such as indoor residual spraying (IRS) and the use of bednets. In East Africa such an integrated effort to control VL does not yet exist and control is mainly based on passive case detection and treatment, although bednets are distributed by non-governmental organizations.

VL disproportionately affects the poorest population groups in low-income countries. For the control, or even elimination, of VL to be successful, it will require a great deal of effort that needs to be sustained over the longer term. Ultimately, effective control of VL will depend on a mix of interventions, consisting of early and efficient diagnosis and treatment, with drug regimens that put as little burden as possible on both the patient and the health system. However, the scarcity of reliable data, both epidemiological and economic, contributes to the low visibility and priority given to VL by national disease control programmes and international donors.

Economic analysis provides an important set of tools to underpin the choice for evidence-based policy. Given the limited financial resources available to VL control and the increasing number of possible interventions, especially regarding drug treatment, economic evaluation provides us with methods to systematically compare all relevant options in terms of their costs and consequences and thereby contribute to more rational decision-making and efficient allocation of scarce resources. Moreover it can improve our understanding of factors influencing both the supply of VL interventions and factors related to the behaviour of VL affected households. While economic analysis is increasingly recognized by both national and international policy makers as an important tool in the development of VL control policies, the scarcity of reliable data, in particular on costs, severely hampers efforts to control the disease.

The general objective of this thesis was to improve our understanding on the economic aspects of visceral leishmaniasis, provide evidence for more rational decision making on diagnosis, case management and VL control and justify increased investment in VL research and control in the Indian subcontinent and East Africa.

More specifically, the studies included in this thesis were framed around the following specific objectives: (i) understand the burden of visceral leishmaniasis in India, Nepal and Sudan; (ii) provide a better understanding of the health seeking behaviour and costs of VL illness from the provider and household perspective; and (iii) examine the cost and cost-effectiveness of VL treatment alternatives with a focus on combination therapies.

In terms of the first research question “to understand the burden of visceral leishmaniasis in India, Nepal and Sudan”, we provided in **chapter 2** an overview of studies that investigated the socio-economic aspects of two neglected tropical diseases: visceral leishmaniasis and sleeping sickness. Both of these diseases affect the poorest of the poor in endemic countries, cause considerable direct and indirect costs (even though the national control programmes tend to provide free care) and push affected households deeper into poverty. In **chapter 3**, we studied the socio-economic profile of households living in areas with high VL endemicity relative to the general population of Bihar, using data collected from a community intervention trial (i.e. the KALANET trial) and the third Indian National Family Health Survey (NFHS-3). With 83% of households in VL endemic areas belonging to the poorest two quintiles in Bihar, we showed that in areas where VL occurs, the socio-economic status of households is considerably lower than the rest of Bihar. Bihar, in turn, is also one of the poorest states in India. The study confirmed the general notion that leishmaniasis is strongly associated with poverty and affects the “poorest of the poor”.

The second research question “provide a better understanding of the health seeking behaviour and costs of VL illness from the provider and household perspective” was investigated in **chapter 4** (India), **chapter 5** (Nepal) and **chapter 6** (Sudan). We find that the free provision of VL diagnosis and drugs has been an important policy measure in reducing financial barriers and improving access to VL care. Without this policy the economic burden of VL would have been much higher. However, given the extreme poverty of VL affected households, a VL episode is still catastrophic to most households. One of the main cost determinants to both public hospitals providing VL treatment and households is the duration of treatment and length of hospitalization. In addition, households also incur substantial costs during the health seeking phase (prior to correct VL diagnosis) as well as through the strategies they use to cope with the cost of illness such as loans.

The third research question about the cost and cost-effectiveness of VL treatment alternatives was investigated in chapters 7 and 8. In **chapter 7** we carried out a review of the current evidence and potential of combination therapies for the treatment of VL. With no new compounds expected to come to the market in the near future and the risk of resistance to

new medicines, combination therapy is proposed as a possible alternative. Combining drugs from different chemical classes could also reduce treatment duration or total drug doses, resulting in fewer toxic effects, higher compliance, and less burden on the health system. Combination therapies may also reduce the overall costs (direct and indirect) and be more cost-effective. This assumption was investigated in *chapter 8* where we assessed the cost-effectiveness of all possible mono- and combination therapies for the treatment of visceral leishmaniasis in the Indian subcontinent using a decision analytical model based on a decision tree. The analysis was done from a societal perspective and included all costs irrespective of whether they were borne by the government or the patient/household (i.e. health provider and household perspective). Our results suggested that a combination of miltefosine and paromomycin provided the lowest cost-effectiveness ratio at US\$92 per death averted. The next best alternative was liposomal amphotericin B and paromomycin with an incremental cost-effectiveness ratio of US\$ 652. Due to a high drug efficacy and expected compliance, the most effective treatment option was a single dose of 10mg per kg of liposomal amphotericin B. This treatment is also recommended by WHO as the preferred monotherapy as it is safe and effective and only a single dose is needed allowing point-of-care diagnosis and treatment on the same day. Such a strategy would significantly reduce the economic burden to households and would relieve public health care facilities in VL endemic areas particularly during the peak transmission periods. However, L-AmB needs to be administered intravenously and stored below 25°C. In addition, the incremental cost-effectiveness ratio of liposomal amphotericin B was US\$ 8,224 due to the cost of the drug, which is its main barrier to widespread use. Although price negotiations between the WHO and the manufacturer reduced the price of liposomal amphotericin B (AmBisome®) to US\$ 20 per vial at the time of the study, as several vials are needed per treatment course, the total cost of liposomal amphotericin B was higher than any of the combination therapies. For liposomal amphotericin B to become the most cost-effective strategy, we showed in our study that the price would need to be reduced to under US\$ 10 per vial.

Although making strong recommendations about the best treatment options is beyond the scope of this thesis, we have shown that, from a societal perspective, more effective and cheaper alternatives are available. Combination therapies and liposomal amphotericin B monotherapy are possible alternatives for the current first-line treatment with miltefosine in the Indian subcontinent, while a preliminary cost analysis for East Africa presented in the discussion section suggests that a combination of sodium stibogluconate and paromomycin is the cheapest and has a similar efficacy to the current first line treatment. This strategy is already successfully implemented in East Africa by non-governmental organizations (NGO's) such as Médecins sans Frontières; similarly liposomal amphotericin B based strategies are also

used on a small scale by NGO's in India at the primary health care level and have overcome the cold chain constraints.

Lastly, in the discussion section we presented a case for investment in VL treatment. We showed that interventions for VL treatment are within the range of highly cost-effective public health interventions, with cost-effectiveness estimates (cost per DALY averted) similar to those for other neglected tropical diseases. However, this does not mean that VL treatment is affordable to national control programmes. Further efforts are therefore needed to: (i) reduce the delays between the onset of symptoms and correct VL diagnosis, (ii) change the first line treatment so as to reduce the length of treatment, and (iii) bring the price of liposomal amphotericin B further down.

SAMENVATTING



SAMENVATTING

Viscerale leishmaniose (VL) is een tropische ziekte veroorzaakt door de *Leishmania* parasiet en wordt overgedragen door de beet van een geïnfecteerde vrouwelijke zandvlieg. De ziekte is een belangrijk volksgezondheidsprobleem en oorzaak van sterfte en morbiditeit op het Indische subcontinent en in Oost-Afrika. Sinds 2005 zijn de regeringen van India, Nepal en Bangladesh partij in een regionaal initiatief om VL te bestrijden, en indien mogelijk te elimineren op het Indische subcontinent. Bestrijdingsstrategieën bestaan enerzijds uit het vroeger diagnosticeren van de ziekte en het verbeteren van behandeling en begeleiding van VL patiënten; anderzijds probeert men risicopopulaties minder bloot te stellen aan de vector (i.e. vector bestrijding) door het verstuiwen van insecticide binnenshuis en het gebruik van met insecticide geïmpregneerde bednetten. In Oost-Afrika bestaat een dergelijke geïntegreerde aanpak nog niet en bestrijding bestaat grotendeels uit het passief detecteren en behandelen van VL gevallen, hoewel geïmpregneerde bednetten worden uitgedeeld door niet-gouvernementele organisaties.

VL treft hoofdzakelijk arme en gemarginaliseerde bevolkingsgroepen in landen met lage inkomens. Bestrijding, en zeker eliminatie van VL, kan enkel succesvol zijn indien de inspanningen die worden geleverd over een langere periode worden volgehouden. Doeltreffende bestrijding van VL zal uiteindelijk afhangen van een combinatie van maatregelen, inclusief tijdige diagnose en behandeling, met behandelingsregimes die de last voor zowel de patiënt als het gezondheidssysteem zoveel mogelijk beperken. Het gebrek aan betrouwbare gegevens omtrent VL, zowel epidemiologisch als economisch, draagt bij tot de lage zichtbaarheid en prioriteit die aan VL wordt gegeven zowel in nationale ziektebestrijdingprogramma's als door internationale donoren.

Economische analyse kan een belangrijke bijdrage leveren tot “evidence-based” beleid rond VL. Gezien de beperkte financiële middelen voor VL bestrijding, en het toenemende aantal mogelijke interventies (vooral rond de behandeling van de ziekte) kan men door economische evaluatiemethoden de kosten en effecten/baten van verschillende bestrijdingsstrategieën van VL vergelijken; also kan worden bijgedragen tot een meer rationele besluitvorming en doelmatige verdeling van schaarse middelen. Economische analyse kan ook betere inzichten geven omtrent factoren die zowel het aanbod van VL interventies als het zoekgedrag van gezinnen naar VL-zorg bepalen. Nationale en internationale beleidsmakers beschouwen economische analyse in toenemende mate als een belangrijk instrument in de ontwikkeling en ondersteuning van een VL beleid; het gebrek aan betrouwbare gegevens, vooral rond kosten, belemmert in grote mate de inspanningen om de ziekte te voorkomen, bestrijden of elimineren.

De algemene doelstelling van deze doctoraatsthesis is inzicht verwerven in de economische aspecten van viscerale leishmaniose, bijdragen tot meer rationele besluitvorming rond diagnose, behandeling en bestrijding van VL, alsook het motiveren voor bijkomende (financiële) middelen voor VL onderzoek en bestrijding op het Indische subcontinent en in Oost-Afrika.

De thesis is onderverdeeld in drie delen gestructureerd rond de volgende specifieke doelstellingen: (i) inzicht verwerven in de ziektelast van VL in India, Nepal en Soedan; (ii) een betere kennis verwerven van het zoekgedrag en de kosten van VL-ziekte vanuit het perspectief van de zorgverstrekker en het gezin; en (iii) onderzoeken van de kosten en kosten-effectiviteit van alternatieve behandelingen van VL met een focus op combinatietherapieën.

Voor de eerste onderzoeksvraag geven we in **hoofdstuk 2** een overzicht van studies die de socio-economische aspecten van twee verwaarloosde tropische ziekten hebben onderzocht: viscerale leishmaniose en slaapziekte. Beide aandoeningen treffen de “armsten der armen” in endemische landen, veroorzaken aanzienlijke directe en indirecte kosten (ondanks gratis zorg door nationale bestrijdingsprogramma's) en duwen getroffen gezinnen verder in de armoede. In **hoofdstuk 3** hebben we het socio-economische profiel bestudeerd van gezinnen die wonen in gebieden van Bihar met een hoge VL endemiciteit. De gegevens komen van een community intervention trial (i.e. de KALANET studie) en van een nationale gezinsenquête (i.e. Third National Family Health Survey; NFHS-3). Drieëntachtig percent van de gezinnen in VL-endemische gebieden behoren tot de armste twee kwintielen in Bihar, wat aantoonde dat in gebieden waar VL voorkomt, de socio-economische status van gezinnen aanzienlijk lager is dan in de rest van Bihar, één van de armste provincies in India. Deze studie bevestigt de algemene indruk dat leishmaniose sterk geassocieerd is met armoede en de “armste der armen” treft.

De tweede onderzoeksvraag – beter inzicht verwerven in het zoekgedrag en de directe en indirecte kosten van VL vanuit het perspectief van de zorgverstrekker en het gezin – werden bestudeerd in **hoofdstuk 4** (India), **hoofdstuk 5** (Nepal) en **hoofdstuk 6** (Soedan). We vinden dat het gratis verstrekken van VL diagnose en behandeling een belangrijke beleidsmaatregel is geweest die de financiële barrières heeft verminderd en heeft geleid tot een betere toegang tot VL zorg. Zonder deze maatregelen zou de economische last van VL veel hoger zijn. Omdat vele gezinnen in extreme armoede leven, is een VL episode voor hen nog altijd katastrofaal. Eén van de belangrijkste kostenfactoren zowel voor openbare ziekenhuizen als gezinnen is de duur van de behandeling en de hospitalisatie. De financiële last voor gezinnen is ook zeer hoog door de uitgaven tijdens het zoekproces vóór de correcte VL diagnose, en door het mobiliseren van de nodige financiële middelen zoals het aangaan van een lening om de medische kosten te dekken.

Voor de derde onderzoeksvraag bestudeerden we in hoofdstukken 7 en 8 de kost en kosten-effectiviteit van alternatieve VL behandelingen. In **hoofdstuk 7** is een literatuurstudie gedaan naar de huidige kennis en potentieel van combinatietherapieën voor de behandeling van VL. Combinatietherapieën zijn belangrijk omdat er in de nabije toekomst geen nieuwe geneesmiddelen op de markt zullen komen, en de ontwikkeling van resistentie van de parasiet tegen de huidige geneesmiddelen een reëel risico is. Het combineren van geneesmiddelen kan ook leiden tot een kortere behandelingsduur, het gebruik van lagere dosissen waardoor er minder toxische bijwerkingen optreden, een betere therapietrouw, en in het algemeen minder belasting van het gezondheidssysteem. Combinatietherapieën kunnen ook leiden tot lagere (directe en indirecte) kosten en kosten-effectiever zijn. Dit laatste werd onderzocht in **hoofdstuk 8** waar we de kosten en effecten van alle mogelijke mono -en combinatietherapieën voor de behandeling van VL hebben vergeleken voor het Indische subcontinent. De analyse werd uitgevoerd vanuit een maatschappelijk perspectief. Onze resultaten tonen aan dat een combinatie van miltefosine en paromomycine de laagste kosten-effectiviteit ratio heeft met US\$ 92 per vermeden sterfgeval. Het tweede beste alternatief was een combinatie van liposomale amfotericine B met paromomycine dat een incrementele kosten-effectiviteit ratio heeft van US\$652. De meest doeltreffende behandeling was één dosis van 10mg liposomale amfotericine B omwille van de hoge efficaciteit en ook de verwachte therapietrouw. Deze behandeling wordt ook aanbevolen door de Wereldgezondheidsorganisatie (WGO) omdat ze veilig en doeltreffend is, en ze maakt “point-of-care” diagnose en behandeling op dezelfde dag mogelijk. Deze behandeling zou ook de economische last voor gezinnen sterk verminderen evenals de werkdruk in openbare ziekenhuizen verlichten, vooral tijdens de transmissieperiode van VL. Er zijn een aantal nadelen aan het gebruik van liposomale amfotericine B: het moet intraveneus worden toegediend en bewaard aan een temperatuur lager dan 25 °C. De grootste struikelblok voor het wijdverspreid gebruik is echter de prijs van het geneesmiddel; de incrementele kosten-effectiviteit ratio was US\$ 8,225. Hoewel de prijs van liposomale amfotericine B (AmBisome®) sterk is gedaald door onderhandelingen tussen de WGO en de producent, toch is het nog altijd aanzienlijk duurder dan eender welke combinatietherapie. We hebben aangetoond dat een strategie met liposomale amfotericine B kosten-effectief kan zijn indien de prijs per flacon lager is dan US\$ 10.

Hoewel specifieke aanbevelingen maken over de meest gepaste behandeling buiten het bestek van deze doctoraatsthesis valt, hebben we aangetoond dat er meer doeltreffende en goedkopere behandelingen bestaan: zowel combinatietherapieën als liposomale amfotericine B zijn mogelijke alternatieven voor de huidige eerstelijnsbehandeling met miltefosine op het Indische subcontinent. Een preliminaire kostenanalyse voor Oost-Afrika, vermeld in de discussie van de thesis, toont aan dat een combinatie van sodium stibogluconate met

paromomycine het goedkoopste alternatief is en een vergelijkbare doeltreffendheid heeft als de huidige eerstelijnsbehandeling met sodium stibogluconate. Deze combinatietherapie wordt met succes reeds gebruikt door niet-gouvernementele organisaties zoals Artsen zonder Grenzen. Liposomale amfotericine B wordt ook al gebruikt op kleine schaal in gezondheidscentra in India, ook door niet-gouvernementele organisaties.

Ten slotte hebben we in de discussie een uitgebreide argumentatie gegeven voor investeringen in VL behandeling. We hebben aangetoond dat de behandeling van VL zich binnen de criteria van zeer kosten-effectieve interventies situeert, met een kost per vermeden DALY die vergelijkbaar is met die van andere tropische verwaarloosde ziektes. Dit impliceert echter niet dat VL behandeling betaalbaar is voor nationale bestrijdingsprogramma's. Verdere inspanningen zijn daarom noodzakelijk om: (i) de tijd tussen de eerste symptomen van VL en correcte diagnose te verminderen, (ii) een andere eerstelijnsbehandeling gebruiken om de duur van de behandeling in te korten, en (iii) bijkomende prijsdalingen van liposomale amfotericine B te bekomen.

ACKNOWLEDGMENTS



ACKNOWLEDGMENTS

There are many people I would like to thank who contributed to the completion of this thesis, and with whom I had the chance to exchange ideas and work together over the past years.

I would like to express my deeply felt gratitude to Prof. Marleen Boelaert. Marleen was more than just a thesis promoter, but gave me the opportunities, support and resources to develop a research career in academia. She provided me my first field assignment abroad in 2005 in Muzaffarpur (famous for Shahi lychees and unfortunately visceral leishmaniasis (VL) as well), offered me a job in her Unit at the Institute of Tropical Medicine in Antwerp in 2008 and engaged me in a broad spectrum of interesting studies and projects. Under her guidance, we introduced and developed the need for economic analysis in research and policy-making for VL control and sleeping sickness, which ultimately led to the completion of this thesis. Marleen, thank you for your kind support and guidance; working with you has always been a pleasure (and still is!), and I hope I'll be able to reflect some of your enthusiasm, high scientific standards and hard work in my future endeavours.

A special word of thanks to my promoter and co-promoter at the Radboud University Nijmegen Medical Center: Prof. Koos van der Velden, for making it possible to pursue a PhD degree at the Radboud University Nijmegen Medical Center, for your guidance, scientific inputs and giving me that final push to (finally...) submit the thesis. To Dr. Rob Baltussen, who I first met as a teacher during my master studies in health economics at the Erasmus University Rotterdam, many thanks for supervising and mentoring my PhD, for the motivating discussions, your technical guidance and insightful comments that have greatly contributed to improving my skills in the field of cost and cost-effectiveness analysis. It was always a pleasure being at the NICHE (Nijmegen International Centre for Health Systems Research and Education) research centre.

I would like to thank all my co-authors with whom I had the privilege to collaborate during the past years. In particular Surendra Uranw and Abuzaid Abdallah for the enjoyable work experience on various studies in Nepal and Sudan, and Kamlesh Gidwani for his assistance at the Kala-Azar Medical Research Centre (KAMRC) in Muzaffarpur, India. I am very grateful to Prof. Suman Rijal from the BP Koirala Institute of Health Sciences (Nepal), Prof. Shyam Sundar from Banaras Hindu University (India), and Prof. Ahmed Musa from the Institute of Endemic Diseases (Sudan). The PhD thesis would not have been possible without their guidance and strong support during my fieldwork and stay at their research institutes. Many thanks to Dr. Manica Balasegaram, working at the time at the Drugs for Neglected Diseases Initiative (now

Executive Director of the Access campaign at Médecins sans Frontières), for his advice and inputs to the studies in Sudan.

Furthermore I would like to thank all my colleagues and friends at ITM in the *landschap* office and elsewhere (Professor Patrick van der Stuyft, Veerle Vanlerberghe, Bart Ostyn, Epco Hasker, David Hendrickx, Pol De Vos, Pierre Lefèvre, Séverine Thys, Albert Picado, Pascal Lutumba, and many others not mentioned here but certainly not forgotten) for their support and providing such a pleasant and productive working environment. Many thanks to the Unit secretariat: Anne Marie, Arabella and Greet for their patience with me and (still) handling my endless stream of requests and questions. I also want to thank all PhD researchers at ITM, present and past, for making ITM such a cosmopolitan and interesting place to work (Alonso, Lely (thank you for hosting my farewell party!), Larissa, Patrick, Elsa, Michel, Alberto and again so many others). And of course the NICHE collaborators (Noor, Sten, Marianne, Caroline, Ernst, Genevieve, Evelinn, Dragana). Finally, a special word of thanks to Veerle and Sten for standing beside me on this rather nerve-wrecking day.

I also would like to acknowledge the financial support I received over the past years which allowed the development and completion of this PhD. This PhD and my employment at ITM, would not have been possible without the generous financial support from a philanthropic donation to the Institute, referred to as the *VL Fund*. During the last 6 months of writing the thesis, I was financially supported by the ITM SOFI-A programme of the Flemish Government. I would also like to thank the Drugs for Neglected Diseases Initiative (DNDi) for providing financial support to the studies in Sudan, as well as the NIDIAG collaborative project supported by the European Commission 7th framework agreement programme (grant agreement 260260).

Lastly, “but not least” as is often said, I would like to thank my family and in particular my parents for all their love and encouragements given throughout the years. From a very young age, I had the incredible chance to travel around the globe with my parents, often for my fathers’ work as an academic. These experiences have undoubtedly shaped me both as a person, but also as a researcher/academic. It is hard to express my gratitude and thanks to you in words and the unconditional support you have given me to all the choices I made in both my personal and professional life (such as moving now to Cape Town)

Filip Meheus

Cape Town, South Africa

June 2013

CURRICULUM VITAE



CURRICULUM VITAE

Filip Meheus (born 7 September 1978, Ghent, Belgium) grew up in Antwerp (Belgium) and Ferney-Voltaire (France). In 2000 he graduated in Applied Economic Sciences at the University of Antwerp (RijksUniversitair Centrum Antwerpen) with a major in International Economics. Upon completion of his studies he was awarded the Price of the Minister of Foreign Trade from the former Belgian Foreign Trade Office (Ministry of Foreign Affairs, Foreign Trade, and Development Cooperation) and worked intermittently for one year in Morocco.

After a brief period in the private sector, Filip became in 2002 research assistant in the unit of Globalisation and Economic Development at the Institute of Development Policy and Management (University of Antwerp); he obtained a Master in Health Economics at the Erasmus University Rotterdam in 2006. From 2006 to 2008 he was a health economist at the Royal Tropical Institute in Amsterdam (The Netherlands) and joined in 2008 the Institute of Tropical Medicine in Antwerp (Belgium) to work on neglected tropical diseases with Prof. Marleen Boelaert (Unit of Epidemiology and Disease Control).

Since January 2013, Filip joined the Health Economics Unit (School of Public Health & Family Medicine) of the University of Cape Town in South Africa and remains visiting fellow at the Institute of Tropical Medicine in Antwerp.

PUBLICATIONS



PUBLICATIONS

Publications presented in this thesis

- Boelaert M, [Meheus E](#), Robays J, Lutumba P (2010) Socio-economic aspects of neglected diseases: sleeping sickness and visceral leishmaniasis. *Ann Trop Med Parasitol*. 104: 535–542
- Boelaert M, [Meheus E](#), Sanchez A, et al. (2009) The poorest of the poor: a poverty appraisal of households affected by visceral leishmaniasis in Bihar, India. *Trop Med Int Health* 14: 639-644
- [Meheus E](#), Boelaert M, Baltussen R, Sundar S (2006) The costs of patient management of visceral leishmaniasis in Muzaffarpur, India. *Trop Med Int Health* 11: 1715-1724
- Uranw S[§], [Meheus E](#)[§], Baltussen R, Rijal S, Boelaert M (2013) The economic burden of visceral leishmaniasis on households in eastern Nepal. *PLoS Negl Trop Dis*. 7: e2062 (§ shared first authorship)
- [Meheus E](#), Abuzaid AA, Baltussen R, et al. (2013) The economic burden of visceral leishmaniasis in Sudan: an assessment of hospital and household costs. *Submitted for publication*.
- Van Griensven J, Balasegaram M, [Meheus E](#), et al. (2010) Combination therapy for visceral leishmaniasis. *Lancet Infect Dis* 10: 184 -194
- [Meheus E](#), Balasegaram M, Oliaro P, et al. (2010) Cost-effectiveness analysis of combination therapies for visceral leishmaniasis in the Indian subcontinent. *PLoS Negl Trop Dis* 4: e818.

Other publications

- Phan V, Thai S, Koole O, Menten J, [Meheus E](#), et al. (2013) Validation of a clinical prediction score to target viral load testing in adults with suspected first line treatment failure in resource-constrained settings. *J Acquir Immune Defic Syndr* 62: 509-16.
- Uranw S, Hasker E, Roy L, [Meheus E](#), et al. (2013) An outbreak investigation of visceral leishmaniasis among residents of Dharan town, eastern Nepal: evidence for urban transmission of *L. donovani*. *BMC Infect Dis* 13: 21.
- Marcotty T, Thys E, Conrad P, Godfroid J, Craig P, Zinsstag J, [Meheus E](#), et al. (2013) Intersectoral collaboration between the medical and veterinary professions in low-resource societies: The role of research and training institutions. *Comp Immunol, Microbiol Infect Dis*. 36: 233-39
- Blecher M, [Meheus E](#), Kollipara A, et al. (2012) Financing vaccinations - the South African experience. *Vaccine* 30S: C79-C86.
- [Meheus E](#), Rijal S, Lutumba P, Hendrickx D, Boelaert M (2012) Neglected Tropical Disease control and health system strengthening: a continuing debate. *Lancet* 379: 2149-2150 [letter].
- Mills A, Ataguba J, Akazili J, Borghi J, Garshong B, Makawia S, Mtei G, Harris B, Macha J, [Meheus E](#), McIntyre D (2012) Equity in financing and use of health care in Ghana, South Africa and Tanzania: implications for paths to universal coverage. *Lancet*. DOI:10.1016/S0140-6736(12)60357-2
- Mtei G, Makawia S, Ally M, Kuwawenaruwa A, [Meheus E](#), Borghi J (2012) Who pays and who benefits from health care? An assessment of equity in health care financing and benefit distribution in Tanzania. *Health Policy Plan*, 27(S1):i23-i34

Bisoffi Z, Sirima SB, [Meheus E](#), et al. (2011) Strict adherence to malaria rapid test results might lead to a neglect of other dangerous diseases: a cost benefit analysis from Burkina Faso. *Malaria J* 10: 226.

[Meheus E](#), Boelaert M (2010) The burden of visceral leishmaniasis in South Asia. *Trop Med Int Health*, 15: 1-3

Rijal S, Uranw S, Chappuis F, Picado A, Khanal B, Paudel I, Andersen E, [Meheus E](#), et al. (2010) The epidemiology of Leishmania donovani infection in high-transmission foci in Nepal. *Trop Med Int Health*, 15(2): 21-28.

Singh SP, Picado A, Boelaert M, Gidwani K, Andersen E, Ostyn B, [Meheus E](#), et al. (2010) The epidemiology of Leishmania donovani infection in high-transmission foci in India. *Trop Med Int Health* 15: 12-20.

Vassall A, Seme A, Compennolle P, [Meheus E](#) (2010) The patient costs of accessing collaborative TB/HIV interventions in Ethiopia. *Int J Tuberc Lung Dis* 14:604–610.

Afriandi I, Siregar AY, [Meheus E](#), et al. (2010) Costs of hospital-based methadone maintenance treatment in HIV/AIDS control among injecting drug users in Indonesia. *Health Policy*. 95: 69-73.

Borghi J, Ataguba J, Mtei G, Akazili J, [Meheus E](#), Rehnberg C, McIntyre D (2009) Methodological challenges in evaluating health care financing equity in data-poor contexts: Lessons from Ghana, South Africa and Tanzania. *Adv Health Econ Health Serv Res* 21:133-56.

Tun Lin W, Lenhart A, Vu Sinh N, Rebollar-Téllez EA, Morrison AC, Barbazan P, Cote M, Midega J, Sanchez F, Manrique-Saide P, Kroeger A, Nathan MB, [Meheus E](#), Petzold M (2009) Reducing costs and operational constraints of dengue vector control by targeting productive larval/pupal habitats: a multi-country non-inferiority randomized trial. *Trop Med Int Health* 14: 1143-53.

McIntyre D, Garshong B, Mtei G, [Meheus E](#), et al. (2008) Beyond fragmentation and towards universal coverage: Insights from Ghana, South Africa and Tanzania. *Bull World Health Organ*, 86(11): 871-876.

Posse M, [Meheus E](#), Van Asten H, Van Der Ven A, Baltussen R (2008) Barriers to access to antiretroviral treatment: a review of recent studies in developing countries. *Trop Med Int Health* 13: 904-913.

[Meheus E](#), Van Doorslaer E (2008) Achieving better measles immunization coverage: do the poor benefit? *Soc Sci Med*, 66: 1709-18.

Lutumba P, [Meheus E](#), Robays J, et al. (2007) Cost-effectiveness of algorithms for confirmation test of Human African trypanosomiasis. *Emerg Infect Dis* 13: 1484-1490.

Vanlerberghe V, Diap G, Guerin P, [Meheus E](#), et al. (2007) Drug policy for visceral leishmaniasis: a cost-effectiveness analysis. *Trop Med Int Health* 12: 274-283.

Lutumba P, Makieya E, Shaw A, [Meheus E](#), Boelaert M (2007) Human African trypanosomiasis in a rural community, Democratic Republic of Congo. *Emerg Infect Dis* 13: 248-254.

Batson A, [Meheus E](#), Brooke S. (2006). Innovative financing mechanisms to accelerate the introduction of HPV vaccines in developing countries. *Vaccine*, S3: S219-25.

Lutumba P, Robays J, Miaka mia Bilenge C, Betu Ku Mesu VK, Molisho D, Declercq J, Van der Veken W, [Meheus E](#), Jannin J, Boelaert M (2005) Trypanosomiasis control, Democratic Republic of Congo, 1993-2003. *Emerg Infect Dis* 11: 1382-1388.

